SEVERE COLD AND COUGH RELIEF DAYTIME- acetaminohpen, dextromethorphan hbr, phenylephrine hcl liquid Good Sense (Geiss, Destin & Dunn, Inc.)

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold
 - minor aches and pains
 - headache
 - nasal and sinus congestion
 - sore throat
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product,

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with a rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for

children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- do not take more than 6 doses (180 mL) in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL= milliliter
- keep dosing cup with product
- adults and children 12 years and over:
 30 mL every 4 hours
- children under 12 years of age: do not use

Other information

- each 30 mL contains: sodium 16 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

acesulfame potassium, alcohol, citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

Principal Display Panel

For Ages 12 Years and Over

Adult

DayTime

Severe Cold & Cough Relief

Acetaminophen - Pain reliever/Fever reducer

Dextromethorphan HBr - Cough Suppressant

Phenylephrine HCl - Nasal Decongestant

- Nasal Congestion
- Sore Throat
- Headache
- Body Ache
- Cough
- Fever

Compare to the active ingredients in Theraflu® DayTime Severe Cold & Cough*

Alcohol Content: 10%

Cherry Flavor

FL OZ (mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by GSK Consumer Health, distributor of Theraflu® Daytime Severe Cold & Cough.

Distributed by: Geiss, Destin & Dunn, Inc.

Peachtree City, GA 30269

www.valuelabels.com

Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Product Label

GOODSENSE®

For Ages 12 Years and Over

Adult

Davtime

Severe Cold & Cough Relief

Acetaminophen - Pain Reliever/Fever Reducer Dextromethorphan HBr - Cough Suppressant Phenylephrine HCI - Nasal Decongestant

Nasal Congestion - Sore Throat - Headache
 Body Ache - Cough - Fever



nts? Call 1-877-753-3935 day 9AM-SPM EST 8.3 FL OZ (245 mL)

6036 00260 7 Alcohol Content: 10% **Cherry Flavor**

h

Drug Facts (continued) depression, psychiatric, or emotional conditions,

or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have liver disease thyroid disease diabetes high blood pressure trouble urinating due to an enlarged prostate gland cough that occurs with too much phlegm (mucus) epersistent or chronic cough such as occurs with smoking, asthma, or emphysema Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days redness or swelling is present

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Theraflu® Daytime Severe Cold & Cough.

PLD-I264D LB003544

NDC 50804-316-08

Drug Facts	
Active ingredients (in each 30 mL)	Purposes
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Phenylephrine HCl 10 mg	

nasal and sinus congestion sore throat

- cough due to minor throat and bronchial irritation
- temporarily reduces fever

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

skin reddening blisters rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

Drug Facts (continued)
 new symptoms occur cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 6 doses (180 mL) in any 24-hour period ■ keep dosing cup with product ■ measure only with dosing cup provided. Do not use any other dosing device. ■ mL = milliliter ►

Drug Facts (continued)

- adults and children 12 years and over
- 30 mL every 4 hours

children under 12 years of age: do not use

Other information

each 30 mL contains: sodium 16 mg store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients acesulfame potassium, alcohol, citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

PEEL CORNER FOR MORE DRUG FACTS

GOOD SENSE Adult DayTime Severe Cold & Cough Relief

SEVERE COLD AND COUGH RELIEF DAYTIME

acetaminohpen, dextromethorphan hbr, phenylephrine hcl liquid

Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:50804-316

Active Ingred	dient/Active	Moiety					
J		dient Name		Basis of Stre	ngth	Strength	
ACETAMINOPHEI	•	L9D) (ACETAMINOPHEN - UNI	I:36209ITL9D)	ACETAMINOPHEN 650 mg in 30 mL		650 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KY (DEXTROMETHORPHAN - UNII:7355X3ROTS))	DEXTROMETHORPHAN HYDROBROMIDE		20 mg in 30 mL		
HENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHI NII:1W5297W6MV)		NYLEPHRINE -	- PHENYLEPHRINE HYDROCHLORIDE		10 mg in 30 mL		
Inactive Ingr	adiants						
Inactive Ingredients Ingredient Name						Strength	
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)							
ALCOHOL (UNII: 3K9958V90M)							
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)							
EDETATE DISOD							
FD&C BLUE NO.							
FD&C RED NO. 4							
GLYCERIN (UNII: I	· · · · · · · · · · · · · · · · · · ·	•					
MALTITOL (UNII: I							
PROPYLENE GLY	COL (UNII: 6DCS	Q167V3)					
WATER (UNII: 059	QF0KO0R)						
SODIUM BENZO	ATE (UNII: 0]245	FE5EU)					
TRISODIUM CITR	ATE DIHYDRAT	Έ (UNII: Β22547Β95Κ)					
Product Cha	racteristics						
Color		Se	core				
Shape		Si	ze				
Flavor		CHERRY In	nprint Code				
Contains							
Contains							
Packaging			M	larketing Start	Mark	eting End	
Packaging # Item Code		ackage Description		larketing Start Date	Mark	eting End Date	
Packaging # Item Code		TTLE, PLASTIC; Type 0: Not	2	Date	Mark 07/31/2	Date	
Packaging # Item Code 1 NDC:50804- 316-08	245 mL in 1 BC Combination Pr	OTTLE, PLASTIC; Type 0: Not roduct	2	Date		Date	
Packaging # Item Code 1 NDC:50804- 316-08 Marketing	245 mL in 1 BC Combination Pr	DTTLE, PLASTIC; Type 0: Not roduct	a 07	Date /31/2016	07/31/2	Date 2025	
Packaging # Item Code 1 NDC:50804-	245 mL in 1 BC Combination Pr	OTTLE, PLASTIC; Type 0: Not roduct	a 07	Date	07/31/2 Mark		

Revised: 10/2022

Good Sense (Geiss, Destin & Dunn, Inc.)