NICOTINE- nicotine polacrilex lozenge Rite Aid Corporation

Drug Facts

Active ingredient (in each lozenge)

Nicotine polacrilex USP, 2 mg

Purpose

Stop smoking aid

Use

 reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding,

only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness,

- diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away

Directions

- if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2	1 lozenge every 2 to 4	1 lozenge every 4 to 8
hours	hours	hours

- nicotine lozenge is a medicine and must be used a certain way to get the best results
- place the lozenge in your mouth and allow the lozenge to slowly dissolve (about 20 to 30 minutes). Minimize swallowing. **Do not chew or swallow lozenge.**
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved (about 20 to 30 minutes)
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.
- It is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

Other information

- each lozenge contains: sodium 5 mg
- store at 20° to 25°C (68° to 77°F)
- keep bottle tightly closed and protect from light

Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, cherry flavor, hydroxypropyl cellulose, magnesium stearate, mannitol, potassium bicarbonate, povidone, sodium alginate, sodium carbonate, sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredient in Nicorette® mini Lozenge*

FREE | SUGAR

FROM | FREE

mini

Nicotine Lozenges

Nicotine Polacrilex Lozenge

Stop Smoking Aid

SUGAR FREE

2 mg

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette **MORE THAN 30 MINUTES** after waking up, use the 4 mg Nicotine Polacrilex Lozenge.

includes User's Guide

CHERRY FLAVOR

LOZENGES

TO INCREASE YOUR SUCCESS IN QUITTING:

- **1.** You must be motivated to quit.
- 2. Use Enough Use at least 9 Nicotine Polacrilex lozenges per day during the first six weeks.
- **3.** Use Long Enough Use Nicotine Polacrilex lozenges for the full 12 weeks.
- 4. Use With a Support Program as directed in the enclosed User's Guide.
- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

*This product is not affiliated, with manufactured by, or produced by the makers or owners of Nicorette® Lozenge.

TAMPER EVIDENT FEATURE: DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.

RETAIN OUTER CARTON FOR FULL PRODUCT USES, DIRECTIONS AND WARNINGS.

DISTRIBUTED BY:

RITE AID, 30 HUNTER LANE,

CAMP HILL. PA 17011

www.riteaid.com

Package Label



RITE AID Mini Nicotine Lozenges Cherry Flavor

NICOTINE nicotine polacrilex lozenge Product Information Product Type HUMAN OTC DRUG Item Code (Source) Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	2 mg	

Inactive Ingredients		
Ingredient Name	Strength	
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
MANNITOL (UNII: 30WL53L36A)		
POTASSIUM BICARBONATE (UNII: HM5Z15LEBN)		
POVIDONE (UNII: FZ989GH94E)		
SODIUM ALGINATE (UNII: C269C4G2ZQ)		
SODIUM CARBONATE (UNII: 45P3261C7T)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		
XANTHAN GUM (UNII: TTV12P4NEE)		

Product Characteristics			
Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor	CHERRY	Imprint Code	A501
Contains			

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:11822- 9902-6	3 in 1 BOX	07/30/2020	02/01/2027		
1		27 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product				
2	NDC:11822- 9902-8	1 in 1 BOX	07/30/2020	02/01/2027		
2		27 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product				

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210711	07/30/2020	02/01/2027

Revised: 11/2025 Rite Aid Corporation