

NAPROXEN SODIUM - naproxen sodium tablet, film coated
Camber Consumer Care Inc

Naproxen Sodium Tablets USP, 220 mg (NSAID)

ACTIVE INGREDIENT(S)

(in each tablet)

Naproxen sodium USP 220 mg (naproxen 200 mg)
(NSAID)*

*nonsteroidal anti-inflammatory drug

PURPOSE

Pain reliever/ fever reducer

USE(S)

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - headache
 - the common cold
 - menstrual cramps
 - toothache
- temporarily reduces fever

WARNINGS

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

DO NOT USE

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

ASK A DOCTOR BEFORE USE IF

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

WHEN USING THIS PRODUCT

- take with food or milk if stomach upset occurs

STOP USE AND ASK A DOCTOR IF

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

PREGNANCY/BREASTFEEDING

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless

definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center right away.

DIRECTIONS

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none">• take 1 tablet every 8 to 12 hours while symptoms last• for the first dose you may take 2 tablets within the first hour• do not exceed 2 tablets in any 8- to 12-hour period• do not exceed 3 tablets in a 24-hour period
Children under 12 years	<ul style="list-style-type: none">• ask a doctor

OTHER INFORMATION

- **each tablet contains:** sodium 21 mg
- Store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

INACTIVE INGREDIENTS

corn starch, FD&C blue #2, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, sodium starch glycolate, stearic acid, talc and titanium dioxide.

QUESTIONS OR COMMENTS?

1-888-588-1418 (Mon - Fri 9AM - 5PM EST)

Do not use if bottle is open or if foil seal imprinted with "SEALED for YOUR PROTECTION" on bottle opening is missing or broken.

Mfg. Lic. No.: 24/MD/TS/2016/F/G

Distributed by:
Camber Consumer Care, Inc.
Piscataway, NJ 08854, USA.

PRINCIPAL DISPLAY PANEL

Naproxen Sodium Round Shape 100's container Label

Top Layer

CAMBER
CONSUMER CARE

NDC 69230-329-01

Mfg. Lic. No.: 241MD/TS/2016/F16

3169230132901

2102082

**Naproxen Sodium Tablets USP,
220 mg (NSAID)**
Pain reliever/fever reducer
STRENGTH TO LAST 12 HOURS
Round Shaped Tablets 100 Tablets

Actual Size

Without glue dotted line area

Un varnish area for Batch coding details
40 x 30 mm

Distributed by: Camber Consumer Care, Inc.
Piscataway, NJ 08854, USA.

40 mm

115 mm

Top Layer Back

See Carton For Full Information

Warnings Allergy alert:
Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, or blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a non steroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer.

40 mm

115 mm

Second Layer

Ask a doctor or pharmacist before use if you are taking any other drug.

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better), have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling), or if pain or fever gets worse.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get

medical help or contact a Poison Control Center right away.

Directions Adults and children 12 years and older: take 1 tablet with water every 8 to 12 hours while symptoms last. For the first dose you may take 2 tablets within the first hour. Do not exceed 3 tablets in a 24 hour period. Children under 12 years: ask a doctor.


Questions or comments? 1-888-588-1418 (Mon - Fri 9AM - 5PM EST)

Do not use if bottle is open or if foil seal imprinted with "SEALED for YOUR PROTECTION" on bottle opening is missing or broken.

40 mm

115 mm

Naproxen Sodium Round Shape 100's container Carton Label



NAPROXEN SODIUM

naproxen sodium tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69230-329
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg	
Inactive Ingredients			
Ingredient Name	Strength		
STARCH, CORN (UNII: O8232NY3SJ)			
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)			
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)			
MICROCRYSTALLINE CELLULOSE 101 (UNII: 7T9FYH5QMK)			
POVIDONE K30 (UNII: U725QWY32X)			

SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	

Product Characteristics

Color	blue	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	H;N11
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69230-329-24	1 in 1 CARTON	10/28/2022	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:69230-329-50	1 in 1 CARTON	10/28/2022	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:69230-329-01	1 in 1 CARTON	10/28/2022	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:69230-329-02	1 in 1 CARTON	10/28/2022	
4		200 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:69230-329-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	10/28/2022	
6	NDC:69230-329-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	10/28/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211065	10/28/2022	

Labeler - Camber Consumer Care Inc (079539968)

Establishment

Name	Address	ID/FEI	Business Operations
Annora Pharma Private Limited		650980746	manufacture(69230-329)