

**ACETAMINOPHEN 500MG- acetaminophen tablet**  
**Pharmaceutica North America, Inc.**

-----

**Active Ingredient(in each tablet)**

Acetaminophen 500mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash
- If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if the user has** liver disease

**Ask a doctor or pharmacist before use if the user is** taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of the reach of children.**

**Overdose warning:** In the case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

• **do not take more than directed (see overdose warning)**

**adults and children 12 years and over:**

- take 2 tablets every 6 hours while symptoms last
- do not take more than 6 tablets in 24 hours, unless directed by a doctor
- do not take for more than 10 days unless directed by a doctor

**children under 12 years:**ask a doctor

### ***Other information***

- store between 20-25°C (68-77°F)

### ***Inactive Ingredients***

povidone, pregelatinized corn starch, sodium starch glycolate, stearic acid

***Questions or comments? (888)788-6472***

**Product label**



NDC 45861-0072-01

# Acetaminophen Extra Strength 500mg Tablet

Pain Reliever / Fever Reducer

100 Tablets

### Drug Facts

**Active Ingredient (in each tablet)**  
Acetaminophen 500mg,.....Pain reliever / fever reducer

**Uses**  
Temporarily relieves minor aches and pains due to:  
■ headache ■ muscular aches ■ backache  
■ minor pain of arthritis ■ the common cold ■ toothache  
■ premenstrual and menstrual cramps ■ temporarily reduces fever

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.

**Do not use:** with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist if you are allergic to acetaminophen or any of the inactive ingredients in this product.

**Drug Facts (continued under label)**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**



Questions or comments? (888)788-6472

Manufactured For:  
Pharmaceuticals North America, Inc.  
Glendale, CA 91205



**NO COPY  
ADHESIVE ZONE**

### Drug Facts (continued)

Ask a doctor before use if the user has liver disease. Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

**Stop use and ask doctor if** ■ pain gets worse or lasts more than 10 days in adults ■ pain gets worse or lasts more than 5 days in children under 12 years ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present  
These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.**

**Overdose warning:** In the case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

Do not take more than directed (see overdose warning)  
**Adults and children 12 years and over:** ■ take 2 tablets every 6 hours while symptoms last ■ do not take more than 6 tablets in 24 hours, unless directed by a doctor ■ do not take for more than 10 days unless directed by a doctor  
**Children under 12 years:** ask a doctor

**Other information**  
Store between 20-25°C (68-77°F)

**Inactive ingredients**  
povidone, pregelatinized corn starch, sodium starch glycolate, stearic acid

**Questions or comments? (888)788-6472**

Manufactured For:  
Pharmaceuticals North America, Inc.  
Glendale, CA 91205

## ACETAMINOPHEN 500MG

acetaminophen tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:45861-072
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

### Inactive Ingredients

Ingredient Name	Strength
POVIDONE K30 (UNII: U725QWY32X)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z 65AP)	

**Product Characteristics**

<b>Color</b>	white (Off White)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:45861-072-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2025	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/01/2025	

**Labeler** - Pharmaceutica North America, Inc. (962739699)

Revised: 3/2025

Pharmaceutica North America, Inc.