

**TOPCARE CHILDRENS COLD AND COUGH- brompheniramine maleate,
dextromethorphan hbr, phenylephrine hcl solution
Topco Associates LLC**

Topco Associates LLC. Children's Cold and Cough Drug Facts

Active ingredients (in each 10 mL)

Brompheniramine maleate, USP 2 mg

Dextromethorphan HBr, USP 10 mg

Phenylephrine HCl, USP 5 mg

Purposes

Antihistamine

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- heart disease

- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers

When using this product

- do not use more than directed
- may cause marked drowsiness
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- mL = milliliter

age	dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other information

- each 10 mL contains: sodium 4 mg
- store at 20°-25°C (68°-77°F)

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution

Questions or comments?

1-888-423-0139

Principal Display Panel

COMPARE TO CHILDREN'S DIMETAPP® COLD & COUGH ACTIVE INGREDIENTS

children's Cold & Cough

ANTIHISTAMINE – BROMPHENIRAMINE MALEATE

COUGH SUPPRESSANT – DEXTROMETHORPHAN HBr

NASAL DECONGESTANT – PHENYLEPHRINE HCl

Relieves Nasal Symptoms

Stuffy Nose

Runny Nose

Sneezing

OUR PHARMACISTS RECOMMEND

Plus Other Symptoms

Itchy, Watery Eyes

Cough

ALCOHOL FREE

Ages 6 Years and over

GRAPE FLAVOR

4 FL OZ (118mL)

NDC 36800-987-26



COMPARE TO CHILDREN'S DIMETAPP® COLD & COUGH ACTIVE INGREDIENTS*

children's Cold & Cough

ANTIHISTAMINE - BROMPHENIRAMINE MALEATE
COUGH SUPPRESSANT - DEXTROMETHORPHAN HBR
NASAL DECONGESTANT - PHENYLEPHRINE HCl

Relieves Nasal Symptoms

- Stuffy Nose
- Runny Nose
- Sneezing

Plus Other Symptoms

- Itchy, Watery Eyes
- Cough

ALCOHOL FREE

Ages 6 Years and over



NDC 36800-987-26



COMPARE TO CHILDREN'S DIMETAPP® COLD & COUGH ACTIVE INGREDIENTS*

children's Cold & Cough

ANTIHISTAMINE - BROMPHENIRAMINE MALEATE
COUGH SUPPRESSANT - DEXTROMETHORPHAN HBR
NASAL DECONGESTANT - PHENYLEPHRINE HCl

Relieves Nasal Symptoms

- Stuffy Nose
- Runny Nose
- Sneezing

Plus Other Symptoms

- Itchy, Watery Eyes
- Cough


ALCOHOL FREE

Ages 6 Years and over



4 FL OZ (118 mL)

4 FL OZ (118 mL)



DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

www.StopMedicineAbuse.org
Learn about teen medicine abuse

children's Cold & Cough

Drug Facts

Active Ingredients (in each 10 mL)
Brompheniramine maleate, USP 2 mg.....Antihistamine
Dextromethorphan HBr, USP 10 mg.....Cough suppressant
Phenylephrine HCl, USP 5 mg.....Nasal decongestant

Purposes

- temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily restores freer breathing through the nose

Uses

- temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- do not use more than directed
- may cause marked drowsiness
- avoid alcoholic beverages
- alcohol, alcoholic beverages, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Drug Facts (continued)

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- mL = milliliter

age	dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other Information

- each 10 mL contains: sodium 4 mg
- store at 20°-25°C (68°-77°F)

Inactive Ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution

Questions or comments?
1-888-423-0139

DISTRIBUTED BY TOPCO ASSOCIATES LLC
ELK GROVE VILLAGE, IL 60007
©TOPCO PER917 QUESTIONS? 1-888-423-0139
topcare@topco.com www.topcarebrand.com



*This product is not manufactured or distributed by Pfizer, distributor of Children's Dimetapp® Cold & Cough.



LOT NO.

EXP.

: 96726 88 C4

TOPCARE CHILDRENS COLD AND COUGH

brompheniramine maleate, dextromethorphan hbr, phenylephrine hcl solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-987
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
BROMPHENIRAMINE MALEATE (UNII: IXA7C9ZN03) (BROMPHENIRAMINE - UNII:H57G17P2FN)	BROMPHENIRAMINE MALEATE	2 mg in 10 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 10 mL

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics			
Color	PURPLE (Clear bluish-red)	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-987-26	1 in 1 CARTON	09/29/2006	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/29/2006	

Labeler - Topco Associates LLC (006935977)