

**VICKS NYQUIL SEVERE COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, and phenylephrine hydrochloride solution**  
**The Procter & Gamble Manufacturing Company**

-----

**Vicks<sup>®</sup> NyQuil<sup>™</sup> Severe Cold & Flu Liquid**

***Drug Facts***

**Active Ingredients (in each 30 mL)**

**Acetaminophen** 650 mg  
Dextromethorphan HBr 20 mg  
Doxylamine succinate 12.5 mg  
Phenylephrine HCl 10 mg

**Purpose**

Pain reliever/fever reducer  
Cough suppressant  
Antihistamine  
Nasal decongestant

**Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease • heart disease • high blood pressure
- thyroid disease • diabetes • glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present • new symptoms occur

- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

|                                 |                   |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 4 to under 12 yrs      | ask a doctor      |
| children under 4 yrs            | do not use        |

**Other information**

- **each 30 mL contains:**sodium 92 mg
- store at no greater than 25°C and do not refrigerate

**Inactive ingredients**

citric acid, FD&C Blue No.1, FD&C Red No. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

**Questions?**

**1-800-362-1683**

**TAMPER EVIDENT:**

**Do not use if printed shrinkband is broken or missing.**

**DIST. BY PROCTER & GAMBLE**

**CINCINNATI OH 45202**

# PRINCIPAL DISPLAY PANEL - 354 mL Bottle Label

VICKS®

NyQuil™

SEVERE

COLD & FLU

**Acetaminophen**, Phenylephrine HCl, Doxylamine Succinate, Dextromethorphan HBr  
Headache, Fever, Sore Throat, Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Sneezing, Runny Nose

Cough

12 FL OZ (354 mL)

Nighttime Relief



Headache, Fever, Sore Throat,  
Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Sneezing, Runny Nose

Cough



12 FL OZ (354 mL)

Nighttime Relief



**Drug Facts (continued)**

- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

|                                 |                   |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 4 to under 12 yrs      | ask a doctor      |
| children under 4 yrs            | do not use        |

**Other information**

- each 30 mL contains: sodium 92 mg
- store at no greater than 25°C and do not refrigerate

**Inactive ingredients** citric acid, FD&C Blue No. 1, FD&C Red No. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

**Questions?** 1-800-362-1683

**Drug Facts (continued)**

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- 3 or more alcoholic drinks every day while using this product
- with other drugs containing acetaminophen

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease • heart disease • high blood pressure
- thyroid disease • diabetes • glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- do not use more than directed
- excitability may occur, especially in children

**TAMPER EVIDENT:**  
Do not use if printed shrinkband is broken or missing.

← PEEL BACK FOR DRUG FACTS

**Drug Facts**

| Active ingredients (in each 30 mL) | Purpose                     |
|------------------------------------|-----------------------------|
| Acetaminophen 650 mg               | Pain reliever/fever reducer |
| Dextromethorphan HBr 20 mg         | Cough suppressant           |
| Doxylamine succinate 12.5 mg       | Antihistamine               |
| Phenylephrine HCl 10 mg            | Nasal decongestant          |

**Uses** temporarily relieves common cold/flu symptoms:

- nasal congestion • sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep • minor aches & pains
- headache • fever • sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

**PARENTS:**  
Learn more about this medicine at [www.DrugFacts.gov](http://www.DrugFacts.gov)

**DIST. BY PROCTER & GAMBLE, CINCINNATI OH 45202**  
Patents: [www.pg.com/patents](http://www.pg.com/patents)

**P&G**  
[www.pg.com](http://www.pg.com)

21521482

## VICKS NYQUIL SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, and phenylephrine hydrochloride solution

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:37000-812 |
| <b>Route of Administration</b> | ORAL           |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength                | Strength            |
|--|----------------------------------|---------------------|
| <b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)                    | ACETAMINOPHEN                    | 650 mg<br>in 30 mL  |
| <b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN<br>HYDROBROMIDE | 20 mg<br>in 30 mL   |
| <b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)                | DOXYLAMINE SUCCINATE             | 12.5 mg<br>in 30 mL |
| <b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)      | PHENYLEPHRINE<br>HYDROCHLORIDE   | 10 mg<br>in 30 mL   |

## Inactive Ingredients

| Ingredient Name                                   | Strength |
|---|----------|
| <b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP) |          |
| <b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)     |          |
| <b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)     |          |
| <b>GLYCERIN</b> (UNII: PDC6A3C0OX)                |          |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)        |          |
| <b>WATER</b> (UNII: 059QF0KO0R)                   |          |
| <b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)        |          |
| <b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)         |          |
| <b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)         |          |
| <b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)          |          |
| <b>SORBITOL</b> (UNII: 506T60A25R)                |          |
| <b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)               |          |
| <b>XANTHAN GUM</b> (UNII: TTV12P4NEE)             |          |

## Product Characteristics

|                 |       |                     |  |
|-----------------|-------|---------------------|--|
| <b>Color</b>    | red   | <b>Score</b>        |  |
| <b>Shape</b>    |       | <b>Size</b>         |  |
| <b>Flavor</b>   | BERRY | <b>Imprint Code</b> |  |
| <b>Contains</b> |       |                     |  |

## Packaging

| # | Item Code        | Package Description  | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:37000-812-08 | 236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 07/22/2013           |                    |
| 2 | NDC:37000-812-12 | 354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 07/22/2013           |                    |
| 3 | NDC:37000-812-24 | 2 in 1 PACKAGE   | 07/22/2013           |                    |
| 3 |                  | 354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product |                      |                    |
| 4 | NDC:37000-812-36 | 3 in 1 PACKAGE   | 07/22/2013           | 07/10/2019         |
| 4 |                  | 354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product |                      |                    |
| 5 | NDC:37000-812-01 | 30 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product  | 10/13/2020           |                    |
| 6 | NDC:37000-       | 4 in 1 PACKAGE   | 07/22/2013           |                    |

| 812-04                       | 4 III I PACKAGE   | 07/13/2021           |                    |
|------------------------------|---|----------------------|--------------------|
| 6                            | 30 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product |                      |                    |
| <b>Marketing Information</b> |   |                      |                    |
| Marketing Category           | Application Number or Monograph Citation                      | Marketing Start Date | Marketing End Date |
| OTC Monograph Drug           | M012  | 07/22/2013           |                    |

**Labeler** - The Procter & Gamble Manufacturing Company (004238200)

Revised: 2/2026

The Procter & Gamble Manufacturing Company