

ASPIRIN- aspirin tablet, coated
PD-Rx Pharmaceuticals, Inc.

Aspirin 325 mg

Active ingredient (in each tablet)

Aspirin 325 mg(NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains
- ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome:Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert:Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you are taking a diuretic
- you have a history of stomach problems, such as heartburn
- you have: -high blood pressure -heart disease -liver cirrhosis -kidney disease -asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: -feel faint -vomit blood -have bloody or black stools -have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- any new symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in

pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- swallow whole, do not chew or crush
- do not exceed recommended dose
- adults and children 12 years and older: take 1-2 tablets every 4 hours, as needed, not more than 12 tablets in 24 hours, or as directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 20°C - 25°C (68°F - 77°F); excursions permitted between 15°C - 30°C (59°F - 86°F)

Inactive ingredients

black iron oxide, cellulose, corn starch, D&C yellow #10 lake, FD&C yellow #6 lake, hypromellose, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, propylene glycol, shellac wax, silicon dioxide, simethicone, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate.

Questions or comments?

1-800-540-3765

package label

Drug Facts	Drug Facts (continued)	Drug Facts (continued)				
<p>Active Ingredient (in each tablet) Purpose Aspirin 325 mg (NSAID) Pain reliever *Nonsteroidal anti-inflammatory drug</p> <p>USES • for the temporary relief of minor aches and pains • ask your doctor about other uses for aspirin</p> <p>Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed</p>	<p>Do not use if you are allergic to aspirin or any other pain reliever/fever reducer • if you ever had an allergic reaction to this product or any of its ingredients</p> <p>Ask a doctor before use if • the stomach bleeding warning applies to you • you are taking a diuretic • you have a history of stomach problems such as heartburn • you have: • high blood pressure • heart disease • liver cirrhosis • kidney disease • asthma</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug for • diabetes • gout • arthritis</p> <p>Stop use and ask a doctor if • you experience any of the following signs of stomach bleeding - feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • an allergic reaction occurs. Seek medical help right away - pain gets worse or lasts more than 10 days • redness or swelling is present • fever gets worse or lasts more than 3 days • any new symptoms occur • ringing in the ears or loss of hearing occurs. These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p>	<p>Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Directions: • drink a full glass of water with each dose • swallow whole, do not chew or crush • do not exceed recommended dose</p> <table border="1"> <tr> <td>adults and children 12 years and older</td> <td>Take 1-2 tablets every 4 hours, as needed, not more than 12 tablets in 24 hours, or as directed by a doctor</td> </tr> <tr> <td>children under 12</td> <td>do not use unless directed by a doctor</td> </tr> </table> <p>Other information • store between 20°C-25°C (68°F-77°F); excursions permitted between 15°C-30°C (59°F-86°F)</p> <p>Inactive Ingredients: black iron oxide, cellulose, corn starch, DCC yellow #10 lake, FD&C yellow #6 lake, hypromellose, polydextrose, polyethylene glycol, polyvinyl acetate, povidone, propylene glycol, shellac wax, silica, stearic acid, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate.</p> <p>Questions or comments? 1-800-540-3765 GTIN: 00372789556079 SNO: C26F1500003 EXP: 07/2027 LOT: C26F15</p>	adults and children 12 years and older	Take 1-2 tablets every 4 hours, as needed, not more than 12 tablets in 24 hours, or as directed by a doctor	children under 12	do not use unless directed by a doctor
adults and children 12 years and older	Take 1-2 tablets every 4 hours, as needed, not more than 12 tablets in 24 hours, or as directed by a doctor					
children under 12	do not use unless directed by a doctor					

NDC 72789-556-07



Aspirin
325 mg

Regular Strength
Enteric Coated
Pain Reliever (NSAID)
Marketed and Packaged By:
PD-Rx Pharmaceuticals, Inc
Oklahoma City, OK 73127
1-405-942-3040

v.12.19.0

7 Tablets
TAMPER EVIDENT: DO NOT USE IF
SEAL IS BROKEN OR MISSING FROM BOTTLE.

ASPIRIN

aspirin tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72789-556(NDC:57896-921)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength

FERROSFERRIC OXIDE (UNII: XM0M87F357)
STARCH, CORN (UNII: O8232NY3SJ)
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)
HYPROMELLOSES (UNII: 3NXW29V3WO)
POLYVINYL ACETATE PHTHALATE (UNII: 58QVG85GW3)
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)
POLYDEXTROSE (UNII: VH2XOU12IE)
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
SHELLAC (UNII: 46N107B71O)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
STEARIC ACID (UNII: 4ELV7Z65AP)
SODIUM BICARBONATE (UNII: 8MDF5V39QO)
SODIUM ALGINATE (UNII: C269C4G2ZQ)
TALC (UNII: 7SEV7J4R1U)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
TRIACETIN (UNII: XHX3C3X673)
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)
DIMETHICONE (UNII: 92RU3N3Y1O)

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;227
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72789-556-07	7 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/26/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/01/2000	

Labeler - PD-Rx Pharmaceuticals, Inc. (156893695)

Registrant - PD-Rx Pharmaceuticals, Inc. (156893695)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Revised: 3/2026

PD-Rx Pharmaceuticals, Inc.