

# **ACETAMINOPHEN- acetaminophen tablet, film coated, extended release RITE AID**

-----  
**8 HR Arthritis Pain Relief**

## **Acetaminophen Extended-Release Tablets USP, 650mg**

### **Active ingredient (in each caplet)**

Acetaminophen USP, 650mg

### **Purpose**

Pain reliever/fever reducer

### **Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- premenstrual and menstrual cramps
- the common cold
- headache
- toothache
- temporarily reduces fever

### **Liver warning**

This product contains acetaminophen. Severe Liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

### **Allergy alert**

acetaminophen may cause severe skin reactions

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

### **Do not use**

- with any other drugs containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

### **Ask a doctor before use if you have**

liver disease

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

### **Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

### **If pregnant or breast-feeding**

ask a health professional before use

### **Keep out of reach of children**

Keep out of reach of children

### **Overdose warning**

In case of overdose, get medical help or contact a poison control center right away.(1-800 222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

### **Directions**

- **do not take more than directed (see overdose warning)**

adults

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

under 18 years of age

- ask a doctor

**Other information**

- store between 20-25°C (68-77°F)
- **do not use if foil inner seal is broken or missing.**

**Inactive ingredients**

hydroxy ethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid

**Questions or comments?**

contact **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST

**Principal Display Panel**



DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN  
NDC 11822-5091-5

Compare to the active ingredient in Tylenol® 8HR Arthritis Pain†

# 8 HR ARTHRITIS PAIN RELIEF ACETAMINOPHEN

EXTENDED-RELEASE TABLETS USP, 650 mg

PAIN RELIEVER / FEVER REDUCER  
For the temporary relief of minor arthritis pain



ACTUAL SIZE

**225 CAPLETS\***  
\*CAPSULE-SHAPED BI-LAYER TABLETS

**IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING**  
**DO NOT USE IF FOIL INNER SEAL IS BROKEN OR MISSING**

**Drug Facts**  
**Active ingredient (in each caplet)** Acetaminophen USP 650 mg, ... pain reliever/fever reducer

**Purpose**  
Pain relief/fever reducer

**Uses**  
Temporarily relieves minor aches and pains due to:  
 ■ minor pain of arthritis  
 ■ muscle aches  
 ■ backache  
 ■ premenstrual and menstrual cramps  
 ■ the common cold  
 ■ headache  
 ■ toothache  
 ■ temporarily reduces fever

**Warnings**  
Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take:  
 ■ more than 6 caplets in 24 hours, which is the maximum daily amount  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin redness  
 ■ hives  
 ■ rash  
 ■ a skin reaction occurs, stop use and seek medical help right away

**Do not use**  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease**  
 ■ Stop use and ask a doctor if:  
 ■ pain gets worse or lasts more than 10 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ new symptoms occur  
 ■ redness or swelling is present  
 ■ These could be signs of a serious condition

**CONTAINS NO ASPIRIN**  
DISTRIBUTED BY: RITE AID  
200 NELLBERRY COMMONS  
ETTERS, PA 17319  
www.riteaid.com

MADE IN INDIA

LOT: 1822509152  
EXP: 20200808

ESTABLISHMENT: QUAKERTON, PA  
By prescription only  
Keep out of reach of children

simulate a printed label, fold along dotted line.

Inside (adhesive side)

**Drug Facts (continued)**

**If pregnant or breast-feeding, ask a health professional before use.**  
Keep out of reach of children.  
Overdose warning: In cases of overdose, get medical help or contact a Poison Control Center right away; (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions:**  
 ■ do not take more than directed (see overdose warning)  
 ■ take 2 caplets every 8 hours with water  
 ■ swallow whole; do not crush, chew, split, or dissolve  
 ■ do not take more than 6 caplets in 24 hours  
 ■ do not use for more than 10 days unless directed by a doctor

**under 18 years of age** ■ ask a doctor  
 ■ store between 20°-25°C (68°-77°F)  
 ■ do not use if foil inner seal is broken or missing

**Other information**  
 ■ do not use if foil inner seal is broken or missing

**Inactive ingredients**  
hydroxy ethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, pregelatinized starch, sodium starch glycolate, stearic acid

**Questions or comments?**  
call 1-877-776-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST

\*This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark "Kenvue".

## ACETAMINOPHEN

acetaminophen tablet, film coated, extended release

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-5091
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

### Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	

<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	
<b>HYDROXYETHYL CELLULOSE (140 CPS AT 5%)</b> (UNII: 8136Y38GY5)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

**Product Characteristics**

<b>Color</b>	white (WHITE TO OFF WHITE COLORED)	<b>Score</b>	no score
<b>Shape</b>	OVAL (CAPSULE SHAPED, BICONVEX INTACT FILM COATED TABLETS)	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	G;650
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-5091-6	50 in 1 BOTTLE; Type 0: Not a Combination Product	11/01/2024	
2	NDC:11822-5091-5	225 in 1 BOTTLE; Type 0: Not a Combination Product	11/01/2024	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211544	11/01/2024	

**Labeler - RITE AID (014578892)**