

ADULT LOW DOSE ENTERIC COATED ASPIRIN- aspirin tablet, coated
Geri-Care Pharmaceutical Corp

gc 983

Active ingredient (in each tablet)

Aspirin 81 mg(NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains
- ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if your are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- the stomach bleeding warning applies to you

- you are taking a diuretic
- you have a history of stomach problems, such as heartburn
- you have: -high blood pressure -heart disease -liver cirrhosis -kidney disease -asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: -feel faint -vomit blood -have bloody or black stools -have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- any new symptoms occur
- ringing in the ears or loss of hearing occurs

These can be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose , get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- swallow whole, do not chew or crush
- do not exceed recommended dose
- adults and children 12 years and older: take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor
- children under 12 years: ask a doctor

Other information

- store at 20-25°C (68-77°F); excursions permitted between 15°C-30°C (59°F-86°F)

Inactive ingredients

cellulose, croscarmellose sodium, D&C Yellow #10 Lake, lactose monohydrate, methacrylic acid co-polymer, PEG, polacrillin potassium, polyvinyl alcohol, silica, sodium lauryl sulfate, soya powder, talc, titanium dioxide, triethyl citrate

Questions or comments?

1-800-540-3765

package label

NDC 57896-983-50

GERI CARE®

**ADULT LOW DOSE
ENTERIC COATED
ASPIRIN 81 mg
PAIN RELIEVER (NSAID)**

500 Tablets

Compare to active ingredient
in **BAYER®** Low Dose Aspirin®

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each tablet) *Purpose*
Aspirin 81 mg (NSAID)* Pain reliever
*nonsteroidal anti-inflammatory drug

Uses ■ for the temporary relief of minor aches and pains
■ ask your doctor about other uses for aspirin

Warnings
Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy Alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

*This product is not manufactured or distributed by the owner of the registered trademark Bayer®.
Product of India
Dist. by: Geri-Care PHARMACEUTICALS CORP.
1295 Towbin Ave, Lakewood, NJ 08701

REV 0422B



3 57896 98150 1

PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you are taking a diuretic ■ you have a history of stomach problems, such as heartburn ■ you have: ■ high blood pressure ■ heart disease ■ liver cirrhosis ■ kidney disease ■ asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ an allergic reaction occurs. Seek medical help right away. ■ pain gets worse or lasts more than 10 days ■ redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ any new symptoms occur ■ ringing in the ears or loss of hearing occurs. These can be signs of a serious condition.

if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not exceed recommended dose ■ drink a full glass of water with each dose ■ swallow whole, do not chew or crush

adults and children take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor

children under 12 ask a doctor

Other information ■ store at 20-25°C (68-77°F); excursions permitted between 15°C-30°C (59°F-86°F)

Inactive ingredients: cellulose, croscarmellose sodium, D&C Yellow #10 Lake, lactose monohydrate, methacrylic acid co-polymer, PEG, polacrillin potassium, polyvinyl alcohol, silica, sodium lauryl sulfate, soya powder, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-540-3765

STOP PEELING

ADULT LOW DOSE ENTERIC COATED ASPIRIN			
aspirin tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57896-983
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg	
Inactive Ingredients			
Ingredient Name	Strength		
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)			

D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)
POLACRILIN POTASSIUM (UNII: 0BZ5A00FQU)
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)
BUTYL ACRYLATE/METHYL METHACRYLATE/METHACRYLIC ACID COPOLYMER (18000 MW) (UNII: JZ1374NL9E)
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)
SOYBEAN (UNII: L7HT8F1ZOD)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
SODIUM LAURYL SULFATE (UNII: 368GB5141J)
TALC (UNII: 7SEV7J4R1U)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	P
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-983-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2021	11/30/2024
2	NDC:57896-983-12	120 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2021	04/30/2026
3	NDC:57896-983-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2022	11/30/2025

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/01/2021	04/30/2026

Labeler - Geri-Care Pharmaceutical Corp (611196254)

Registrant - Geri-Care Pharmaceutical Corp (611196254)