

OATHOLOGY INZINCABLE SUNSCREEN SPF 50- homosalate, octocrylene, zinc oxide lotion
SKINOATH LLC

Oathology Inzincable Sunscreen SPF 50

Drug Facts

Active Ingredients

Homosalate 10.00 %
Octocrylene 10.00 %
Zinc Oxide 5.00 %

Purposes

Sunscreen

Uses:

- Helps prevent sunburn.
- higher SPF gives more sunburn protection
- If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings:

For external use only

Do not use

- on damaged or broken skin.

When using this product

- keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if

- rash or irritation develops and lasts.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast-feeding,

ask a health professional before use

Directions:

- Apply liberally 15 minutes before sun exposure and as needed.
- Children under 6 months of age: ask a doctor.
- Reapply at least every 2 hours
- Use a water-resistant sunscreen if swimming or sweating
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease the risk, regularly use a sunscreen with broad spectrum SPF of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 a.m. - 2 p.m.
 - Wear long-sleeved shirts, pants, hats, and sunglasses.

Other Information:

- Protect the product in this container from excessive heat and direct sun.

Inactive Ingredients:

Acetic Acid, Acetyl Hexapeptide-8, Allantoin, Aloe Barbadensis Leaf Juice, Azelaic Acid, Butyloctyl Salicylate, Caprylyl Glycol, Caprylyl Methicone, Citric Acid, Citrullus Lanatus (Watermelon) Fruit Extract, Cyclopentasiloxane, Glycerin, Glyceryl Laurate, Glyceryl Stearate SE, Glyceryl Undecylenate, Hydroxyethyl Acrylate/Sodium Acryloyldimethyl Taurate Copolymer, Isohexadecane, Niacinamide, PEG-10 Dimethicone, Pentylene Glycol, Phenoxyethanol, Polysorbate 20, Polysorbate 60, Polyhydroxystearic Acid, Potassium Sorbate, Sodium Polyacrylate, Sorbitan Isostearate, Sorbitan Sesquioleate, Tocopheryl Acetate, Tribehenin PEG-20 Esters, Water (Aqua), Zingiber Officinale (Ginger) Root Oil.

Questions?

(513) 650-OATH

Package Labeling:

SUNSCREEN + SKINCARE:

AZELAIC ACID: BRIGHTENS SKIN TONE AND IMPROVES REDNESS

GINGER: BRIGHTENS SKIN TONE AND HAS ANTI-AGING PROPERTIES

NIACINAMIDE: PROTECTS AGAINST SUN DAMAGE AND IMPROVES HYPERPIGMENTATION

PEPTIDES: PROMOTE SKIN FIRMNESS AND IMPROVE FINE LINES AND WRINKLES

WATERMELON EXTRACT: HELPS AGAINST SUN DAMAGE AND IS SOOTHING

ZINC: REDUCES OIL AND DECREASES INFLAMMATION



SUITABLE FOR ALL SKIN TONES



DERMATOLOGIST DEVELOPED



VEGAN



FRAGRANCE FREE

QUESTIONS OR COMMENTS?
DROP US A LINE AT:
SUPPORT@OATHOLOGY.COM

OATHOLOGY.COM
#OATHOLOGY

OATHOLOGY®

INZINCABLE®
SUNSCREEN

BROAD SPECTRUM SPF 50
UVA/UVB SUNSCREEN



3.4 FL OZ | 100ML

USES:

HELPS PREVENT SUNBURN.
HIGHER SPF GIVES MORE
SUNBURN PROTECTION. IF USED
AS DIRECTED WITH OTHER SUN
PROTECTION MEASURES (SEE
DIRECTIONS), DECREASES THE
RISK OF SKIN CANCER AND EARLY
SKIN AGING CAUSED BY THE SUN.
WATER RESISTANT (80 MINUTES).

DIRECTIONS:

APPLY GENEROUSLY 15 MINUTES
BEFORE SUN EXPOSURE.
DURING SUN EXPOSURE,
REAPPLY AFTER 80 MINUTES
AFTER SWIMMING OR SWEATING
IMMEDIATELY AFTER TOWEL
DRYING AT LEAST EVERY TWO
HOURS FOR OPTIMAL
PROTECTION. CHILDREN UNDER
SIX MONTHS OF AGE: ASK A
DOCTOR. SUN PROTECTION
MEASURES: SPENDING TIME IN
THE SUN INCREASES YOUR RISK
OF SKIN CANCER AND EARLY
SKIN AGING. TO DECREASE THIS
RISK, REGULARLY USE A
SUNSCREEN WITH A BROAD
SPECTRUM SPF VALUE OF 15 OR
HIGHER AND OTHER SUN
PROTECTION MEASURES
INCLUDING LIMIT TIME IN THE
SUN, ESPECIALLY FROM 10 A.M. -
2 P.M. WEAR LONG-SLEEVED
SHIRTS, PANTS, HATS, AND
SUNGLASSES.

OTHER INFORMATION:

PROTECT THE PRODUCT IN THIS
CONTAINER FROM EXCESSIVE
HEAT AND DIRECT SUN.

USE ONLY AS DIRECTED

DISTRIBUTED BY: SKINOATH LLC
7357 E. KEMPER RD STE. B
CINCINNATI, OH 45249

Drug Facts

Active Ingredients	Purposes
Homosalate 10.00 %	Sunscreen
Octocrylene 10.00 %	Sunscreen
Zinc Oxide 5.00 %	Sunscreen
Uses: <ul style="list-style-type: none">■ Helps prevent sunburn■ higher SPF gives more sunburn protection■ If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun	
Warnings: <p>For external use only</p> <p>Do not use ■ on damaged or broken skin.</p> <p>When using this product: ■ keep out of eyes. Rinse with water to remove.</p> <p>Stop use and ask a doctor if: ■ rash or irritation develops and lasts.</p> <p>Keep out of reach of children.</p> <p>If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>If pregnant or breast-feeding, ask a health professional before use</p>	
Directions: <ul style="list-style-type: none">■ Apply liberally 15 minutes before sun exposure and as needed.■ Children under 6 months of age: ask a doctor.■ Reapply at least every 2 hours■ Use a water-resistant sunscreen if swimming or sweating■ Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease the risk, regularly use a sunscreen with broad spectrum SPF of 15 or higher and other sun protection measures including:<ul style="list-style-type: none">■ Limit time in the sun, especially from 10 a.m. - 2 p.m.■ Wear long-sleeved shirts, pants, hats, and sunglasses.	
Other Information: <ul style="list-style-type: none">■ Protect the product in this container from excessive heat and direct sun.	
Inactive Ingredients: <p>Acetic Acid, Acetyl Hexapeptide-8, Allantoin, Aloe Barbadensis Leaf Juice, Azelaic Acid, Butylcetyl Salicylate, Caprylyl Glycol, Caprylyl Methicone, Citric Acid, Citrullus Lanatus (Watermelon) Fruit Extract, Cyclopentasiloxane, Glycerin, Glyceryl Laurate, Glyceryl Stearate SE, Glyceryl Undecylenate, Hydroxyethyl Acrylate/Sodium Acryloyldimethyl Taurate Copolymer, Isohexadecane, Niacinamide, PEG-10 Dimethicone, Pentaerythritol, Phenoxylphenol, Polysorbate 20, Polysorbate 60, Polyhydroxystearic Acid, Potassium Sorbate, Sodium Polyacrylate, Sorbitan Isostearate, Sorbitan Sesquioleate, Tocopheryl Acetate, Tribehenin PEG-20 Esters, Water (Aqua), Zingiber Officinale (Ginger) Root Oil.</p>	
Questions? (513) 650-OATH	



OATHOLOGY[®]

INZINCABLE[®] SUNSCREEN

BROAD SPECTRUM SPF 50
UVA/UVB SUNSCREEN



NET 3.4 FL OZ | 100ML

USES

HELPS PREVENT SUNBURN. HIGHER SPF GIVES MORE SUNBURN PROTECTION. IF USED AS DIRECTED WITH OTHER SUN PROTECTION MEASURES (SEE DIRECTIONS), DECREASES THE RISK OF SKIN CANCER AND EARLY SKIN AGING CAUSED BY THE SUN. WATER RESISTANT (80 MINUTES).

DIRECTIONS

APPLY GENEROUSLY 15 MINUTES BEFORE SUN EXPOSURE. DURING SUN EXPOSURE, REAPPLY 80 MINUTES AFTER SWIMMING OR SWEATING, IMMEDIATELY AFTER TOWEL DRYING, AND AT LEAST EVERY TWO HOURS FOR OPTIMAL PROTECTION. CHILDREN UNDER SIX MONTHS OF AGE: ASK A DOCTOR. SUN PROTECTION MEASURES: SPENDING TIME IN THE SUN INCREASES YOUR RISK OF SKIN CANCER AND EARLY SKIN AGING. TO DECREASE THIS RISK, REGULARLY USE A SUNSCREEN WITH A BROAD SPECTRUM SPF VALUE OF 15 OR HIGHER AND OTHER SUN PROTECTION MEASURES INCLUDING LIMITING TIME IN THE SUN, ESPECIALLY FROM 10 A.M. - 2 P.M. WEAR LONG-SLEEVED SHIRTS, PANTS, HATS, AND SUNGLASSES.

WARNINGS

FOR EXTERNAL USE ONLY. DO NOT USE ON DAMAGED OR BROKEN SKIN. WHEN USING THIS PRODUCT, KEEP OUT OF EYES. RINSE WITH WATER TO REMOVE. DISCONTINUE USE IF IRRITATION OR REDNESS OCCURS. STOP USE AND ASK A DOCTOR IF RASH OCCURS. KEEP OUT OF REACH OF CHILDREN. IF SWALLOWED, GET MEDICAL HELP OR CONTACT A POISON CONTROL CENTER RIGHT AWAY.

DRUG FACTS

HOMOSALATE 10.00 %, OCTOCRYLENE 10.00 %, ZINC OXIDE 5.00 %

DISTRIBUTED BY: SKINOATH LLC
MASON, OH
OATHOLOGY.COM
#OATHOLOGY

OATHOLOGY INZINCABLE SUNSCREEN SPF 50

homosalate, octocrylene, zinc oxide lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:85562-594
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HOMOSALATE (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	100 mg in 1 mL
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	100 mg in 1 mL
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC CATION - UNII:13S1S8SF37)	ZINC CATION	50 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
ACETYL HEXAPEPTIDE-8 AMIDE (UNII: L4EL31FWML)	
ALLANTOIN (UNII: 344S277G0Z)	
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
AZELAIC ACID (UNII: F2VW3D43YT)	
BUTYLOCTYL SALICYLATE (UNII: 2EH13UN8D3)	
CAPRYLYL GLYCOL (UNII: 00YIU5438U)	
CAPRYLYL TRISILOXANE (UNII: Q95M2P1KJL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
WATERMELON (UNII: 231473QB6R)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
GLYCERYL LAURATE (UNII: Y98611C087)	
GLYCERYL STEARATE SE (UNII: FCZ5MH785I)	
ISOHEXADECANE (UNII: 918X1OUF1E)	
NIACINAMIDE (UNII: 25X51I8RD4)	
PENTYLENE GLYCOL (UNII: 50C1307PZG)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
POLYSORBATE 20 (UNII: 7T1F30V5YH)	
POLYSORBATE 60 (UNII: CAL22UVI4M)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
SORBITAN ISOSTEARATE (UNII: 01S2G2C1E4)	
SORBITAN SESQUIOLEATE (UNII: 0W8RRI5W5A)	
.ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)	
TRIBEHENIN PEG-20 ESTERS (UNII: 84K9EH29Y9)	
WATER (UNII: 059QF0KO0R)	
GINGER OIL (UNII: SAS9Z1SVUK)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85562-594-00	1 in 1 BOX	03/04/2025	
1		100 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	03/04/2025	

Labeler - SKINOATH LLC (090820399)

Registrant - Pure Source, LLC (080354456)