

STRONGID- pyrantel pamoate paste
Zoetis Inc.

Strongid®
(pyrantel pamoate) paste

Horse Dewormer


Read Entire Package Insert Carefully Before Using This Product

DESCRIPTION

Strongid Paste is a pale yellow to buff paste containing 43.9% w/w pyrantel pamoate in an inert vehicle. Each syringe contains 3.6 grams of pyrantel base in 23.6 grams (20 mL) paste. Each mL contains 180 mg pyrantel base as pyrantel pamoate.

Composition

Pyrantel pamoate is a compound belonging to a family classified chemically as tetrahydropyrimidines. It is a yellow, water-insoluble crystalline salt of the tetrahydropyrimidine base and pamoic acid containing 34.7% base activity. The chemical structure and name are given below.



Chemical name: (E)-1,4,5,6-tetrahydro-1-methyl-2-[2-(2-thienyl)-vinyl]-pyrimidine 4,4'-methylenebis [3-hydroxy-2-naphtholate] (1:1)

INDICATIONS FOR USE

For the removal and control of mature infections of large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*); small strongyles; pinworms (*Oxyuris equi*); and large roundworms (*Parascaris equorum*) in horses and ponies.

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

DOSAGE AND TREATMENT

Strongid Paste is to be administered as a single oral dose of 3 mg pyrantel base per lb of body weight. The syringe has 4 weight mark increments. Each weight mark indicates the recommended dose for 300 lb of body weight.

Dosage		
Body Weight Range	Volume	mg Pyrantel Base
up to 300 lb	1/4 syringe (5 mL)	900 mg
301-600 lb	1/2 syringe (10mL)	1800 mg
601-900 lb	3/4 syringe (15 mL)	2700 mg
901-1200 lb	1 full syringe (20 mL)	3600 mg

Note: Position screw-gauge over appropriate mark on plunger. Each mL contains 180 mg of pyrantel base as pyrantel pamoate.

Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

For maximum control of parasitism, it is recommended that foals (2–8 months of age) be dosed every 4 weeks. To minimize the potential source of infection that the mare may pose to the foal, the mare should be treated 1 month prior to anticipated foaling date followed by retreatment 10 days to 2 weeks after birth of foal. Horses and ponies over 8 months of age should be routinely dosed every 6 weeks.

ADMINISTRATION

After removing the cap, the paste should be deposited on the dorsum of the tongue. Introduce the nozzle end of the syringe at the corner of the mouth. Direct the syringe backwards and depress the plunger to deposit the paste onto the tongue. Given in this manner, it is unlikely that rejection of the paste will occur. Raising the horse's head sometimes assists in the swallowing process. When only part of the paste has been used, replace the cap on the syringe nozzle.

EFFICACY

Critical (worm count) studies in horses demonstrated that Strongid Paste administered at the recommended dosage was efficacious against mature infections of *Strongylus vulgaris* (>90%), *S. edentatus* (69%), *S. equinus* (>90%), *Oxyuris equi* (81%), *Parascaris equorum* (>90%), and small strongyles (>90%).

WARNINGS

Do not use in horses intended for human consumption. Keep out of reach of children.

It is recommended that severely debilitated animals not be treated with this preparation.

OTHER WARNINGS

Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers. Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance. Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method). A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

RECOMMENDED STORAGE

Store at controlled room temperature 15°-25°C (59°-77°F).

Approved by FDA under NADA # 129-831

zoetis

Distributed by:
Zoetis Inc.
Kalamazoo, MI 490

52477600
Revised: November 2025

PRINCIPAL DISPLAY PANEL - 20 mL Syringe Carton



52477800

STRONGID				
pyrantel pamoate paste				
Product Information				
Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:54771-2410	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	PYRANTEL PAMOATE (UNII: 81BK194Z5M) (PYRANTEL - UNII:4QIH0N49E7)	PYRANTEL	180 mg in 1 mL	
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54771-2410-2	1 in 1 CARTON		
1		20 mL in 1 SYRINGE, PLASTIC		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NADA	NADA129831	10/26/1982		

