

**GOOD SENSE COMBO PACK DAYTIME AND NIGHTTIME- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride**  
**L. Perrigo Company**

-----

**Perrigo Combo Pack Daytime & Nighttime Drug Facts**

**Nighttime Cold & Flu**

**Active ingredients (in each 30 mL)**

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

**Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

### **Stop use and ask doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

|                                 |                   |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 6 hrs |
| children 4 to under 12 yrs      | ask a doctor      |
| children under 4 yrs            | do not use        |

**Other information**

- **each 30 mL contains:** sodium 36 mg
- store at 20-25°C (68-77°F)

**Inactive ingredients**

alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

**Questions or comments?**

**1-800-719-9260**

**Daytime Cold & Flu****Active ingredients (in each 15 mL)**

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

## Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

## Ask a doctor before use if you have

- liver disease

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**When using this product**

**do not use more than directed**

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

|                                 |                   |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs      | 15 mL every 4 hrs |
| children 4 to under 6 yrs       | ask a doctor      |
| children under 4 yrs            | do not use        |

**Other information**

- **each 15 mL contains:** sodium 15 mg
- store at 20-25°C (68-77°F)

### **Inactive ingredients**

anhydrous citric acid, FD&C yellow no. 6, flavor, glycerin, menthol, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

### **Questions or comments?**

**1-800-719-9260**

### **Package/Label Principal Display Panel**

GOODSENSE®

COMBO PACK DAYTIME & NIGHTTIME

Powerful Relief Day or Night

GOODSENSE®

Multi-Symptom Relief

Non-Drowsy

Pain Reliever, Fever Reducer

Cough Suppressant

Nasal Decongestant

Day Time

Cold & Flu

Acetaminophen

Dextromethorphan HBr

Phenylephrine HCl

- Headache, Fever, Sore Throat, Minor Aches & Pains

- Nasal Congestion

- Cough

Alcohol Free

Compare to active ingredients of Vicks® DayQuil® Cold & Flu

100% SATISFACTION GUARANTEED

GOODSENSE®

Multi-Symptom Relief

Pain Reliever, Fever Reducer

Cough Suppressant

Antihistamine

Night Time

Cold & Flu

Acetaminophen

Dextromethorphan HBr

Doxylamine Succinate

- Headache, Fever, Sore Throat, Minor Aches & Pains
- Sneezing, Runny Nose
- Cough

Cherry Flavor

ALCOHOL 10%

Compare to active ingredients of Vicks® NyQuil® Cold & Flu

100% SATISFACTION GUARANTEED

2 BOTTLES - 1 DAYTIME/1NIGHTTIME

12 FL OZ (354 mL) EACH, TOTAL 24 FL OZ (1.5 pt) (708 mL)

GOODSENSE®

**DayTime**

Cold & Flu

**NightTime**

Cold & Flu

|  |   |  |
|--|---|--|
| PAPER<br>BOX<br><br>Widely Recyclable | PLASTIC<br>BOTTLE<br><br>Widely Recyclable | PLASTIC<br>CUP<br><br>Do Not Recycle |
| Discard Seal,<br>Empty &<br>Replace Cap  |   |  |

how2recycle.info

GOODSENSE®

NDC 0113-6219-02

**COMBO PACK DAYTIME & NIGHTTIME**  
Powerful Relief Day or Night



- Headache, Fever, Sore Throat, Minor Aches & Pains
- Nasal Congestion
- Cough

Alcohol Free

Compare to active ingredients of Vicks® DayQuil® Cold & Flu

100% SATISFACTION GUARANTEED

- Headache, Fever, Sore Throat, Minor Aches & Pains
- Sneezing, Runny Nose
- Cough

ALCOHOL 10%

**Cherry Flavor**

Compare to active ingredients of Vicks® NyQuil® Cold & Flu

100% SATISFACTION GUARANTEED

2 BOTTLES – 1 DAYTIME/1 NIGHTTIME  
12 FL OZ (354 mL) EACH, TOTAL 24 FL OZ (1.5 pt) (708 mL)



73302 C2 C1

**Nighttime Cold & Flu**

| <b>Drug Facts</b>                         |                             |
|---|-----------------------------|
| <b>Active ingredients (in each 30 mL)</b> | <b>Purpose</b>              |
| Acetaminophen 650 mg                      | Pain reliever/fever reducer |
| Dextromethorphan HBr 30 mg                | Cough suppressant           |
| Doxylamine succinate 12.5 mg              | Antihistamine               |

**Uses** temporarily relieves common cold/flu symptoms:  
 ■ cough due to minor throat and bronchial irritation  
 ■ sore throat ■ headache ■ minor aches and pains  
 ■ fever ■ runny nose and sneezing

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
 ■ more than 4,000 mg of acetaminophen in 24 hours  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin reddening ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**  
 ■ liver disease ■ glaucoma  
 ■ cough that occurs with too much phlegm (mucus)  
 ■ a breathing problem such as emphysema or chronic bronchitis  
 ■ trouble urinating due to an enlarged prostate gland  
 ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema  
 ■ a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**  
 ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

**When using this product**  
 ■ excitability may occur, especially in children  
 ■ marked drowsiness may occur ■ avoid alcoholic drinks  
 ■ be careful when driving a motor vehicle or operating machinery  
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**  
 ■ pain or cough gets worse or lasts more than 7 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ redness or swelling is present  
 ■ new symptoms occur  
 ■ cough comes back or occurs with rash or headache that lasts.  
 These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

|   |                   |
|---|-------------------|
| ■ take only as directed – see Overdose warning                      |                   |
| ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs |                   |
| adults & children 12 yrs & over                                     | 30 mL every 6 hrs |
| children 4 to under 12 yrs  | ask a doctor      |
| children under 4 yrs  | do not use        |

**Other information**  
 ■ each 30 mL contains: sodium 36 mg  
 ■ store at 20-25°C (68-77°F)

**Inactive ingredients** alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

**Questions or comments?** 1-800-719-9260

**Daytime Cold & Flu**

| <b>Drug Facts</b>                         |                             |
|---|-----------------------------|
| <b>Active ingredients (in each 15 mL)</b> | <b>Purpose</b>              |
| Acetaminophen 325 mg                      | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg                | Cough suppressant           |
| Phenylephrine HCl 5 mg                    | Nasal decongestant          |

**Uses** temporarily relieves common cold/flu symptoms:  
 ■ nasal congestion ■ cough due to minor throat and bronchial irritation  
 ■ sore throat ■ headache ■ minor aches and pains ■ fever

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if  
 ■ adult takes more than 4,000 mg of acetaminophen in 24 hours  
 ■ child takes more than 5 doses in 24 hours  
 ■ taken with other drugs containing acetaminophen  
 ■ adult has 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin reddening ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**  
 ■ liver disease ■ heart disease ■ high blood pressure  
 ■ thyroid disease ■ diabetes  
 ■ trouble urinating due to an enlarged prostate gland  
 ■ cough that occurs with too much phlegm (mucus)  
 ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product do not use more than directed**

**Stop use and ask a doctor if**  
 ■ you get nervous, dizzy or sleepless  
 ■ pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)  
 ■ fever gets worse or lasts more than 3 days  
 ■ redness or swelling is present  
 ■ new symptoms occur  
 ■ cough comes back or occurs with rash or headache that lasts.  
 These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

|   |                   |
|---|-------------------|
| ■ take only as directed – see Overdose warning                      |                   |
| ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs |                   |
| adults & children 12 yrs & over                                     | 30 mL every 4 hrs |
| children 6 to under 12 yrs  | 15 mL every 4 hrs |
| children 4 to under 6 yrs   | ask a doctor      |
| children under 4 yrs  | do not use        |

**Other information**  
 ■ each 15 mL contains: sodium 15 mg ■ store at 20-25°C (68-77°F)

**Inactive ingredients** anhydrous citric acid, FD&C yellow no. 6, flavor, glycerin, menthol, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

**Questions or comments?** 1-800-719-9260

**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

Gluten Free

Distributed By

**Perrigo®**

Allergan, MI 49010



## GOOD SENSE COMBO PACK DAYTIME AND NIGHTTIME

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride kit

### Product Information

|                     |                |                           |               |
|---------------------|----------------|---------------------------|---------------|
| <b>Product Type</b> | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:0113-6219 |
|---------------------|----------------|---------------------------|---------------|

### Packaging

| # | Item Code        | Package Description                              | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:0113-6219-02 | 1 in 1 CARTON; Type 0: Not a Combination Product | 01/08/2026           |                    |

### Quantity of Parts

| Part # | Package Quantity | Total Product Quantity |
|--------|------------------|------------------------|
| Part 1 | 1 BOTTLE         | 354 mL                 |
| Part 2 | 1 BOTTLE         | 354 mL                 |

## Part 1 of 2

### GOOD SENSE NIGHTTIME

acetaminophen, dextromethorphan hbr, doxylamine succinate solution

### Product Information

|                           |               |
|---------------------------|---------------|
| <b>Item Code (Source)</b> | NDC:0113-0459 |
|---------------------------|---------------|

|                                |      |
|--------------------------------|------|
| <b>Route of Administration</b> | ORAL |
|--------------------------------|------|

### Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|-----------------|-------------------|----------|
|                 |                   | 650 mg   |

|   |                                  |                     |
|---|----------------------------------|---------------------|
| <b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)                       | ACETAMINOPHEN                    | 650 mg<br>in 30 mL  |
| <b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH)<br>(DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN<br>HYDROBROMIDE | 30 mg<br>in 30 mL   |
| <b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)                   | DOXYLAMINE SUCCINATE             | 12.5 mg<br>in 30 mL |

## Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| <b>ALCOHOL</b> (UNII: 3K9958V90M)                          |          |
| <b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)            |          |
| <b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)              |          |
| <b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)              |          |
| <b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)         |          |
| <b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A) |          |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)                 |          |
| <b>WATER</b> (UNII: 059QF0KO0R)                            |          |
| <b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)                 |          |
| <b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR) |          |

## Product Characteristics

|                 |                        |                     |  |
|-----------------|------------------------|---------------------|--|
| <b>Color</b>    | RED (Clear/Dark Red)   | <b>Score</b>        |  |
| <b>Shape</b>    |                        | <b>Size</b>         |  |
| <b>Flavor</b>   | CHERRY (Menthol Aroma) | <b>Imprint Code</b> |  |
| <b>Contains</b> |                        |                     |  |

## Packaging

| # | Item Code        | Package Description                                   | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:0113-0459-40 | 354 mL in 1 BOTTLE; Type 0: Not a Combination Product |                      |                    |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012                                     | 08/22/2011           |                    |

## Part 2 of 2

### GOOD SENSE DAYTIME NON DROWSY COLD AND FLU

acetaminophen, dextromethorphan hbr, phenylephrine hcl solution

## Product Information

|                                |               |
|--------------------------------|---------------|
| <b>Item Code (Source)</b>      | NDC:0113-0656 |
| <b>Route of Administration</b> | ORAL          |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength                | Strength           |
|--|----------------------------------|--------------------|
| <b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)                    | ACETAMINOPHEN                    | 325 mg<br>in 15 mL |
| <b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN<br>HYDROBROMIDE | 10 mg<br>in 15 mL  |
| <b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)      | PHENYLEPHRINE<br>HYDROCHLORIDE   | 5 mg<br>in 15 mL   |

### Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| <b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)            |          |
| <b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)            |          |
| <b>GLYCERIN</b> (UNII: PDC6A3C0OX)                         |          |
| <b>MENTHOL, UNSPECIFIED FORM</b> (UNII: L7T10EIP3A)        |          |
| <b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A) |          |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)                 |          |
| <b>WATER</b> (UNII: 059QF0KO0R)                            |          |
| <b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)                 |          |
| <b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)                  |          |
| <b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)                  |          |
| <b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR) |          |
| <b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)                |          |
| <b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)                        |          |
| <b>XANTHAN GUM</b> (UNII: TTV12P4NEE)                      |          |

### Product Characteristics

|                 |                      |                     |  |
|-----------------|----------------------|---------------------|--|
| <b>Color</b>    | ORANGE (clear)       | <b>Score</b>        |  |
| <b>Shape</b>    |                      | <b>Size</b>         |  |
| <b>Flavor</b>   | MENTHOL (with fruit) | <b>Imprint Code</b> |  |
| <b>Contains</b> |                      |                     |  |

### Packaging

| # | Item Code        | Package Description                                   | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:0113-0656-40 | 354 mL in 1 BOTTLE; Type 0: Not a Combination Product |                      |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
|                    |  |                      |                    |

|                              |   |                             |                           |
|------------------------------|---|-----------------------------|---------------------------|
| OTC Monograph Drug           | M012  | 07/25/2006                  |                           |
| <b>Marketing Information</b> |   |                             |                           |
| <b>Marketing Category</b>    | <b>Application Number or Monograph Citation</b> | <b>Marketing Start Date</b> | <b>Marketing End Date</b> |
| OTC Monograph Drug           | M012  | 01/08/2026                  |                           |

**Labeler** - L. Perrigo Company (006013346)

Revised: 5/2026

L. Perrigo Company