

**LIQUID WORMER 2X- pyrantel pamoate suspension**  
**Durvet Inc**

-----  
**Liquid Wormer 2X**

**Extended Content Label Panel 1**

IMPORTANT: SHAKE WELL BEFORE USING

DOSAGE AND ADMINISTRATION: Administer 1 teaspoon (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment. For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation.

RECOMMENDED STORAGE: STORE BELOW 86°F (30°C).

[Durvet logo]

WWW.DURVET.COM

MANUFACTURED FOR DURVET, INC.

100 S.E. MAELLAN DRIVE

BLUE SPRINGS, MISSOURI 64014

1-800-821-5570 / ISS25XB07

NDC 30798-760-22

DURVET

Liquid Wormer (TM) 2X

(pyrantel pamoate oral suspension)

Canine Anthelmintic Suspension

Active Ingredient: 4.54 mg of Pyrantel Base as pyrantel pamoate per mL

NOT FOR USE IN HUMANS

Approved by FDA under ANADA #200-007

NET CONTENTS: 2 FL OZ (60 mL)

KEEP OUT OF REACH OF CHILDREN:

Keep Liquid Wormer 2X in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

INDICATIONS AND USAGE: For the removal of large roundworms (*Toxocara canis* and

*Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Liquid Wormer 2X suspension may also be used to prevent reinfestation of *T. canis* in puppies and adult dogs and in lactating bitches after whelping. Read accompanying professional information under this label.

Take Time

(Eye Symbol)

Observe Label Directions

PRODUCT LOT NUMBER AND EXPIRATION DATE PRINTED ON BOTTOM OF CONTAINER.  
FOR FULL PRESCRIBING INFORMATION PEEL HERE

## Extended Content Label Panel 2


Liquid Wormer 2X(TM) (pyrantel pamoate oral suspension)

Canine Anthelmintic Suspension

NOT FOR USE IN HUMANS

DESCRIPTION: Liquid Wormer 2X is a suspension of pyrantel pamoate in a palatable butterscotch flavored vehicle. Each mL contains 4.54 mg of pyrantel base as pyrantel pamoate.

Pyrantel pamoate is a compound belonging to a family classified chemically as tetrahydropyrimidines. It is a yellow, water-insoluble crystalline salt of the tetrahydropyrimidine base and pamoic acid containing 34.7% base activity. The chemical structure and name are given to the right:



(E)-1,4,5,6-Tetrahydro-1-methyl-2-[2-(2-thienyl) vinyl] pyrimidine 4,4' methylenebis [3-hydroxy-2-naphthoate](1 :1)

INDICATIONS AND USAGE: Liquid Wormer 2X suspension is a highly palatable formulation intended as a single treatment for the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

### **Extended Content Label Panel 3**

Liquid Wormer 2X suspension may also be used to prevent reinfestation of *T. canis* in puppies and adult dogs and in lactating bitches after whelping.

PRECAUTIONS: THIS PRODUCT IS A SUSPENSION AND AS SUCH WILL SEPARATE. TO ENSURE UNIFORM RESUSPENSION AND TO ACHIEVE PROPER DOSAGE, IT IS EXTREMELY IMPORTANT THAT THE PRODUCT BE SHAKEN THOROUGHLY BEFORE EVERY USE.

KEEP OUT OF REACH OF CHILDREN

DOSAGE AND ADMINISTRATION: Administer 1 teaspoon (5 mL) for each 10 lb of bodyweight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment.

For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation.

SAFETY AND EFFICACY STUDIES: Critical (worm count) studies in dogs demonstrated that pyrantel pamoate at the recommended dosage is highly efficacious against *T. leonina* (99%), *T. canis* (85%), *A. caninum* (97%), and *U. stenocephala* (94%).

One of the most outstanding and significant features of pyrantel pamoate is its wide margin of therapeutic safety in dogs. The acute oral LD(50) of pyrantel pamoate administered in gelatin capsules to female and male dogs is greater than 314 mg base per lb of body weight, which indicates a therapeutic index in excess of 138 times

PROFESSIONAL INFORMATION CONTINUED PEEL HERE

### **Extended Content Label Panel 4**

the recommended dosage. In subacute and chronic studies, no significant morphological abnormalities could be attributed to pyrantel pamoate when administered to dogs at daily dose rates up to 94 mg base per pound of body weight (40x) for periods of 19, 30, and 90 days. Clinical studies conducted in a wide variety of geographic locations using more than 40 different breeds of dogs showed no drug-induced toxic effects. Included in these studies were nursing pups, weaned pups, adults, pregnant bitches, and males at stud.

Take Time

(Eye Symbol)

Observe Label Directions

Additional data has demonstrated the safe use of pyrantel pamoate in dogs having

heartworm infestations and/or receiving medication for heartworms, dogs exposed to organophosphate flea collars or flea/tick-dip treatments, and dogs undergoing concurrent treatment or medication at the time of worming such as immunization and antibacterial treatment.

**RECOMMENDED STORAGE:** STORE BELOW 86° F (30° C). Keep Liquid Wormer 2X in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

**HOW SUPPLIED:** Liquid Wormer 2X is supplied in 2 fl. oz. (60 mL), 8 fl. oz. (236 mL), and 16 fl. oz. (474 mL) bottles.

Approved by FDA under ANADA # 200-007

**Contact Information:** To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Durvet, Inc. at 1-800-821-5570 or [www.durvet.com](http://www.durvet.com).

For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

(Durvet logo)

[WWW.DURVET.COM](http://WWW.DURVET.COM)

### **Extended Content Label Panel 5 (Bottom Ply)**

**IMPORTANT: SHAKE WELL BEFORE USING**

**DOSAGE AND ADMINISTRATION:** Administer 1 teaspoon (5 ml) for each 10 lb of body weight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment. For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation.

**RECOMMENDED STORAGE:** STORE BELOW 86°F (30°C)

(Durvet Logo)

[WWW.DURVET.COM](http://WWW.DURVET.COM)

MANUFACTURED FOR DURVET, INC.

100 S.E. MAGELLAN DRIVE

BLUE SPRINGS, MISSOURI 64014

1-800-821-5570 / ISS25XB07

NDC 30798-760-22

DURVET

Liquid Wormer (TM) 2X

(pyrantel pamoate oral suspension)

Canine Anthelmintic Suspension

Active Ingredient: 4.54 mg of Pyrantel Base as

Pyrantel Pamoate per mL

NOT FOR USE IN HUMANS

Approved by FDA under ANADA #200-007

NET CONTENTS - 2 fl. oz. (60 mL)

KEEP OUT OF REACH OF CHILDREN. Keep Liquid Wormer 2X in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

INDICATIONS AND USAGE: For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Liquid Wormer 2X suspension may also be used to prevent reinfestation of *T. canis* in puppies and adult dogs and in lactating bitches after whelping.

Take Time

(Eye Symbol)

Observe Label

Directions

PRODUCT LOT & EXPIRATION DATE PRINTED ON BOTTOM OF CONTAINER

**PRINCIPAL DISPLAY PANEL - 2 mL Bottle Label**

**NDC 30798-760-22**

EACH 2 OZ BOTTLE TREATS 120 LBS

Liquid Wormer (TM) 2X

(pyrantel pamoate oral suspension)

Canine Anthelmintic Suspension

Removes Large Roundworms & Hookworms in Puppies & Dogs

(Durvet Logo)

**Active Ingredient:**

4.54 mg of Pyrantel Base  
as Pyrantel Pamoate per ml.

MADE IN USA

NOT FOR USE IN HUMANS

KEEP OUT OF REACH OF CHILDREN

**APPROVED BY FDA UNDER ANADA # 200-007**

**NET CONTENTS: 2 FL OZ (60ML)**

**IMPORTANT: SHAKE WELL BEFORE USING**

**DOSAGE AND ADMINISTRATION:** Administer 1 teaspoon (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment. For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation.

**RECOMMENDED STORAGE:** STORE BELOW 86°F (30°C)



MANUFACTURED FOR DURVET, INC.  
100 S.E. MAGELLAN DRIVE  
BLUE SPRINGS, MISSOURI 64014  
1-800-821-5570 / ISS25XB07

NDC 30798-760-22

**durvet.**

# Liquid Wormer™

(pyrantel pamoate oral suspension) **2X**

**Canine Anthelmintic Suspension**

**Active Ingredient:** 4.54 mg of pyrantel base as pyrantel pamoate per mL

**NOT FOR USE IN HUMANS**

Approved by FDA under ANADA # 200-007

**NET CONTENTS: 2 FL OZ (60 mL)**

**KEEP OUT OF REACH OF CHILDREN:**

Keep Liquid Wormer 2X in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

**INDICATIONS AND USAGE:** For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Liquid Wormer 2X suspension may also be used to prevent reinfestation of *T. canis* in puppies and adult dogs and in lactating bitches after whelping. Read accompanying professional information under this label.



PRODUCT LOT NUMBER AND EXPIRATION DATE PRINTED ON BOTTOM OF CONTAINER.


FOR FULL PRESCRIBING INFORMATION **PEEL HERE** ▶

## Liquid Wormer 2X™ (pyrantel pamoate oral suspension) Canine Anthelmintic Suspension NOT FOR USE IN HUMANS

**DESCRIPTION:** Liquid Wormer 2X is a suspension of pyrantel pamoate in a palatable butterscotch flavored vehicle. Each mL contains 4.54 mg of pyrantel base as pyrantel pamoate.

Pyrantel pamoate is a compound belonging to a family classified chemically as tetrahydropyrimidines. It is a yellow, water-insoluble crystalline salt of the tetrahydropyrimidine base and pamoic acid containing 34.7% base activity. The chemical structure and name are given to the right.

**INDICATIONS AND USAGE:** Liquid Wormer 2X suspension is a highly palatable formulation intended as a single treatment for the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies.



The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

BINDING EDGE

BINDING EDGE

Liquid Wormer 2X suspension may also be used to prevent reinfestation of *T. canis* in puppies and adult dogs and in lactating bitches after whelping.

**PRECAUTIONS:** THIS PRODUCT IS A SUSPENSION AND AS SUCH WILL SEPARATE. TO ENSURE UNIFORM RESUSPENSION AND TO ACHIEVE PROPER DOSAGE, IT IS EXTREMELY IMPORTANT THAT THE PRODUCT BE SHAKEN THOROUGHLY BEFORE EVERY USE.

**KEEP OUT OF REACH OF CHILDREN**

**DOSAGE & ADMINISTRATION:** Administer 1 teaspoon (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment.

For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation.

**SAFETY AND EFFICACY STUDIES:** Critical (worm count) studies in dogs demonstrated that pyrantel pamoate at the recommended dosage is highly efficacious against *T. leonina* (99%), *T. canis* (85%), *A. caninum* (97%), and *U. stenocephala* (94%).

One of the most outstanding and significant features of pyrantel pamoate is its wide margin of therapeutic safety in dogs. The acute oral LD(50) of pyrantel pamoate administered in gelatin capsules to female and male dogs is greater than 314 mg base per lb of body weight, which indicates a therapeutic index in excess of 138 times

PROFESSIONAL INFORMATION CONTINUED **PEEL HERE** ▶

the recommended dosage. In subacute and chronic studies, no significant morphological abnormalities could be attributed to pyrantel pamoate when administered to dogs at daily dose rates up to 94 mg base per pound of body weight (40x) for periods of 19, 30, and 90 days. Clinical studies conducted in a wide variety of geographic locations using more than 40 different breeds of dogs showed no drug-induced toxic effects. Included in these studies were nursing pups, weaned pups, adults, pregnant bitches and males at stud.



Additional data has demonstrated the safe use of pyrantel pamoate in dogs having heartworm infestations and/or receiving medication for heartworms, dogs exposed to organophosphate flea collars or flea/tick-dip treatments, and dogs undergoing concurrent treatment or medication at the time of worming such as immunization and antibacterial treatment.

**RECOMMENDED STORAGE: STORE BELOW 86°F (30°C).**

Keep Liquid Wormer 2X in a secure location out of reach of dogs, cats and other animals to prevent accidental ingestion or overdose.

**HOW SUPPLIED:** Liquid Wormer 2X is supplied in 2 fl. oz (60 mL), 8 fl. oz (236 mL), and 16 fl. oz. (474 mL) bottles.

Approved by FDA under ANADA # 200-007

**CONTACT INFORMATION:** To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Durvet, Inc. at 1-800-821-5570 or [www.durvet.com](http://www.durvet.com).

For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>



[WWW.DURVET.COM](http://WWW.DURVET.COM)

BINDING EDGE

GE

**IMPORTANT: SHAKE WELL BEFORE USING**

**DOSAGE AND ADMINISTRATION:** Administer 1 teaspoon (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to or after treatment. Dogs usually find

NDC 30798-760-22

**durvet.**

# Liquid

**KEEP OUT OF EACH OF CHILDREN:**

Keep Liquid Wormer 2X in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

BINDING ED

...this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment. For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation.

**RECOMMENDED STORAGE: STORE BELOW 86°F (30°C)**



MANUFACTURED FOR DURVET, INC.  
100 S.E. MAGELLAN DRIVE  
BLUE SPRINGS, MISSOURI 64014  
1-800-821-5570 / ISS25X807

**Liquid Wormer 2X**  
(pyrantel pamoate oral suspension)

**Canine Anthelmintic Suspension**

**Active Ingredient:** 4.54 mg of pyrantel base as pyrantel pamoate per mL  
**NOT FOR USE IN HUMANS**  
Approved by FDA under ANADA # 200-007  
**NET CONTENTS: 2 FL OZ (60 mL)**

**INDICATIONS AND USAGE:** For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Liquid Wormer 2X suspension may also be used to prevent reinfestation of *T. canis* in puppies and adult dogs and in lactating bitches after whelping.

**Take Time**  
 **Observe Label Directions**

PRODUCT LOT NUMBER AND EXPIRATION DATE PRINTED ON BOTTOM OF CONTAINER.

## LIQUID WORMER 2X

pyrantel pamoate suspension

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:30798-760
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
pyrantel pamoate (UNII: 81BK194Z5M) (pyrantel - UNII:4QIH0N49E7)	pyrantel	4.54 mg in 1 mL

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30798-760-22	60 mL in 1 BOTTLE, PLASTIC		
2	NDC:30798-760-28	237 mL in 1 BOTTLE, PLASTIC		
3	NDC:30798-760-31	473 mL in 1 BOTTLE, PLASTIC		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200007	08/24/2012	

**Labeler** - Durvet Inc (056387798)

**Registrant** - Happy Jack Inc (003189354)

### Establishment

Name	Address	ID/FEI	Business Operations
ABS Corporation		048022727	MANUFACTURE

## Establishment

Name	Address	ID/FEI	Business Operations
COSMA Spa		428655732	API MANUFACTURE

Revised: 10/2025

Durvet Inc