

**PAIN RELIEF- acetaminophen tablet**  
**Meijer Distribution Inc**

-----  
**Meijer Distribution, Inc. Pain Relief Drug Facts**

**Active ingredient (in each caplet)**

Acetaminophen 500 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold
  - headache
  - minor pain of arthritis
  - backache
  - muscular aches
  - toothache
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

liver disease

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"><li>• take 2 caplets every 6 hours while symptoms last</li><li>• do not take more than 6 caplets in 24 hours, unless directed by a doctor</li><li>• do not use for more than 10 days unless directed by a doctor</li></ul>
children under 12 years	ask a doctor

**Other information**

- store at 20-25°C (68-77°F)

**Inactive ingredients**

carnauba wax, corn starch\*, croscarmellose sodium\*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate\*, stearic acid

\*may contain one or more of these ingredients

## Questions or comments?

1-800-719-9260

## Principal Display Panel

meijer®

EASY OPEN CAP

EXTRA STRENGTH

Pain Relief

COMPARE TO EXTRA STRENGTH TYLENOL® ACTIVE INGREDIENT

ACETAMINOPHEN

PAIN RELIEVER | FEVER REDUCER FOR ADULTS

PACKAGE NOT CHILD-RESISTANT

Actual Size

150 CAPLETS

500 mg each

**meijer.**  
EXTRA STRENGTH  
**Pain Relief**  
ACETAMINOPHEN  
PAIN RELIEVER | FEVER REDUCER  
FOR ADULTS

**COMPARE TO EXTRA STRENGTH TYLENOL® ACTIVE INGREDIENT\*\***

**PACKAGE NOT CHILD-RESISTANT**

**150 CAPLETS**  
500 mg each

Actual Size

EASY OPEN CAP

NDC 41250-975-47

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

**GLUE AREA**

DIRTY SEAL  
QUALITY GUARANTEE  
The Perfect Prescription  
www.meijer.com/  
advertising

Disrupt Seal  
Reduces Cap  
Removal Force  
**PLASTIC BOTTLE**  
how2recycle.info

DIST. BY MEIJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com

**Drug Facts**

**Active ingredient (in each caplet):** Acetaminophen 500 mg.....Pain reliever/fever reducer

**Purpose:** Pain reliever/fever reducer

**Uses:** Temporarily relieves minor aches and pains due to:
 

- minor aches and pains
- minor pain of arthritis
- muscular aches
- menstrual and neuralgic cramps
- toothache
- temporarily reduces fever

**Warnings:** Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours. Do not take other drugs containing acetaminophen. If you have ever had an allergic reaction to this product. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 

- skin redness
- blisters
- rash
- skin peeling
- skin sores
- skin swelling
- skin itching

**Directions:** Take 1 or 2 caplets every 6 hours as needed. Do not take more than 10 caplets in 24 hours. Do not take more than 4 caplets at one time. Do not take if you have ever had an allergic reaction to this product or any of its ingredients. Ask a doctor before use if you have liver disease or if you are taking the blood thinning drug warfarin. Do not use and ask a doctor if:
 

- you are pregnant or breastfeeding
- you are taking other drugs containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- you have ever had an allergic reaction to this product or any of its ingredients.

**Other information:** (N) 41250-975-47 (4)

**Inactive ingredients:** carnauba wax, croscarmellose sodium, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid. \*may contain one or more of these ingredients.

**Questions or comments?** 1-800-719-9260

**Directions (continued):**

- Do not take more than directed.
- Do not take more than 10 caplets in 24 hours.
- Do not take more than 4 caplets at one time.
- Do not take if you have ever had an allergic reaction to this product or any of its ingredients.
- Do not take if you are pregnant or breastfeeding.
- Do not take if you are taking other drugs containing acetaminophen (prescription or nonprescription).
- Do not take if you have ever had an allergic reaction to this product or any of its ingredients.

ADHESIVE AREA  
NO COPY

# PAIN RELIEF

acetaminophen tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-975
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

## Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	L484
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-975-47	150 in 1 BOTTLE; Type 0: Not a Combination Product	04/26/1995	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	04/26/1995	

**Labeler** - Meijer Distribution Inc (006959555)

