LORATADINE AND PSEUDOEPHEDRINE SULFATE- loratadine and pseudoephedrine sulfate tablet, film coated, extended release Topco Associates LLC

Drug Facts

ACTIVE INGREDIENTS (IN EACH TABLET)

Loratadine, USP 10 mg

Pseudoephedrine sulfate, USP 240 mg

PURPOSE

Antihistamine

Nasal decongestant

USES

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - sneezing
 - itchy, watery eyes
 - runny nose
 - itching of the nose or throat
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily restores freer breathing through the nose

WARNINGS

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed.

Taking more than directed may cause drowsiness.

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

• do not divide, crush, chew or dissolve the tablet

adults and children 12 years and over	1 tablet daily with a full glass of water; not more than 1 tablet in 24 hours
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

OTHER INFORMATION

- **sodium:** contains 10 mg/tablet
- **calcium:** contains 25 mg/tablet
- TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.
- store between 20° C to 25° C (68° F to 77° F).
- protect from light and store in a dry place

INACTIVE INGREDIENTS

Calcium carbonate, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, iron oxide black, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, sodium alginate, sodium citrate, talc and titanium dioxide

QUESTIONS?

call 1-888-423-0139

PRINCIPAL DISPLAY PANEL

TopCare®

NDC 36800-724-69

NON-DROWSY*

24 HOUR

Original Prescription Strength

Allergy Relief - D

PSEUDOEPHEDRINE SULFATE, USP 240 mg/NASAL DECONGESTANT LORATADINE, USP 10 mg/ANTIHISTAMINE EXTENDED RELEASE TABLETS

Indoor & Outdoor Allergies

Relief of:

Nasal and Sinus Congestion Due to Cold or Allergies

Sneezing; Runny Nose; Itchy, Watery Eyes

Itchy Throat or Nose Due to Allergies

10 EXTENDED RELEASE TABLETS

COMPARE TO CLARITIN-D® 24 HOUR active ingredients†

*When taken as directed. See Drug Facts Panel.

DIST.BY TOPCO ASSOCIATES LLC

5095191/R0412

^	:alcium: contains 25 mg/tablet TER UNITS ARETORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.	
	ask a doctor	

ask a doctor children under 12 years of age

adults and children 12 years and over | 1 tablet daily with a full glass of water, not more than 1 tablet in 24 hours do not divide, crush, chew or dissolve the tablet

Directions

Keep out of reach of children, In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222); if pregnant or breast-feeding, ask a health professional before use.

■ uervousness, dizziness or sieepiessness occurs

■ symptoms do not improve within 7 days or are accompanied by a fever Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

When using this product do not take more than directed. Taking more than directed may cause drowsiness. ■ liver or kidney disease. Your doctor should determine if you need a different dose

■ trouble urinating due to an enlarged prostate gland ■ µiôµ ploog biessnie ■ diabetes Ask a doctor before use if you have ■ thyroid disease ■ heart disease

do not know it your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you


if you have ever had an allergic reaction to this product or any of its ingredients
 if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression.

Marnings

9

- temporarily restores freer breathing through the nose
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies ■ reduces swelling of nasal passages ■ temporarily relieves sinus congestion and pressure

Drug Facts (continued)



Nassal decongestant Antinistamine

Pseudoephedrine sulfate, USP 240 mg.

Purpose

Active ingredients (in each tablet)
Loratadine, USP 10 mg

Drug Facts

NDC 36800-724-69

NON-DROWSY* 24 HOUR

Original Prescription Strength

lergy Relief-D

PSEUDOEPHEDRINE SULFATE, USP 240 mg/NASAL DECONGESTANT LORATADINE, USP 10 mg/ANTIHISTAMINE EXTENDED RELEASE TABLETS **Indoor & Outdoor Allergies**

Nasal and Sinus Congestion Due to Cold or Allergies Sneezing; Runny Nose; Itchy, Watery Eyes Itchy Throat or Nose Due to Allergies



COMPARE TO CLARITIN-D® 24 HOUR active ingredients

[†]The product is not manufactured or distributed by Schering-Plough Healthcare Products, CLARTIN-D® 24 HOUR is a registered trademark of Schering Corporation.

Non Varnish Area

*When taken as directed. See Drug Facts Panel

oxide black, lactose monohydrate, magnesium stearate, microcrystalline celiulose, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, sodium alginate, sodium citrate, talc and titanium dioxide Inactive ingredients calcium carbonate, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, iron

■ store between 20° C to 25° C (68° F to 77° F)

Drug Facts (continued)

Keep the carton. It contains important information. See end panel for expiration date. 6

50951

R0412

QUESTIONS? 1-888-423-0139

DIST. BY TOPCO ASSOCIATES LLC 7711 GROSS POINT ROAD SKOKIE, IL 60077 @TOPCO OHMR412

10 EXTENDED RELEASE TABLETS

Questions? call 1-888-423-0139

■ brotect from light and store in a dry place

LORATADINE AND PSEUDOEPHEDRINE SULFATE

loratadine and pseudoephedrine sulfate tablet, film coated, extended release

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-724
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
LORATADINE (UNII: 7AJO3BO7QN) (LORATADINE - UNII:7AJO3BO7QN)	LORATADINE	10 mg		
PSEUDO EPHEDRINE SULFATE (UNII: Y9 DL7QPE6B) (PSEUDO EPHEDRINE - UNII:7CUC9 DDI9F)	PSEUDOEPHEDRINE SULFATE	240 mg		

Inactive Ingredients			
Ingredient Name	Strength		
CALCIUM CARBONATE (UNII: H0 G9 379 FGK)			
COLLOIDAL SILICON DIO XIDE (UNII: ETJ7Z6XBU4)			
HYDROXYPROPYL CELLULOSE (UNII: RFW2ET671P)			
HYPROMELLOSES (UNII: 3NXW29V3WO)			
FERROSOFERRIC OXIDE (UNII: XM0M87F357)			
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
CELLULO SE, MICRO CRYSTALLINE (UNII: OP1R32D61U)			
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)			
PO VIDO NE (UNII: FZ989 GH94E)			
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
SHELLAC (UNII: 46 N10 7B710)			
SODIUM ALGINATE (UNII: C269C4G2ZQ)			
SODIUM CITRATE (UNII: 1Q73Q2JULR)			
TALC (UNII: 7SEV7J4R1U)			
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)			

Product Characteristics				
Color	white (White to Off-white)	Score	no score	
Shape	CAPSULE	Size	17mm	
Flavor		Imprint Code	RX724	
Contains				

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-724-69	10 in 1 BLISTER PACK		
2	NDC:36800-724-15	15 in 1 BLISTER PACK		

Marketing Information				
plication Number or Monograph Citation	Marketing Start Date	Marketing End Date		
A076557	11/17/2004			

Labeler - Topco Associates LLC (006935977)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment				
Name	Address	ID/FEI	Business Operations	
Ohm Laboratories Inc.		051565745	manufacture(36800-724)	

Revised: 12/2012 Topco Associates LLC