

TRIAMCINOLONE ACETONIDE- triamcinolone acetonide ointment
RPK Pharmaceuticals, Inc.

Triamcinolone Acetonide Ointment USP, 0.025%, 0.1%, 0.5%

For Dermatologic Use Only

Not For Ophthalmic Use

Rx Only

DESCRIPTION

The topical corticosteroids constitute a class of primarily synthetic steroids used as anti-inflammatory and anti-pruritic agents. Triamcinolone acetonide is designated chemically as pregna-1,4-diene-3,20-dione, 9-fluoro-11,21-dihydroxy-16,17-[(1-methylethylidene)bis(oxy)]-, (11 β ,16 α)-. C₂₄H₃₁FO₆, M.W. 434.51; CAS Reg. No. 76-25-5.

Each gram of Triamcinolone Acetonide Ointment USP, 0.025%, 0.1% or 0.5% contains 0.25 mg, 1 mg or 5 mg triamcinolone acetonide, respectively, in an ointment base of light mineral oil and white petrolatum.

CLINICAL PHARMACOLOGY

Topical corticosteroids share anti-inflammatory, anti-pruritic and vasoconstrictive actions. The mechanism of anti-inflammatory activity of the topical corticosteroids is unclear. Various laboratory methods, including vasoconstrictor assays, are used to compare and predict potencies and/or clinical efficacies of the topical corticosteroids. There is some evidence to suggest that a recognizable correlation exists between vasoconstrictor potency and therapeutic efficacy in man.

Pharmacokinetics -

The extent of percutaneous absorption of topical corticosteroids is determined by many factors including the vehicle, the integrity of the epidermal barrier, and the use of occlusive dressings. Topical corticosteroids can be absorbed from normal intact skin. Inflammation and/or other disease processes in the skin increase percutaneous absorption. Occlusive dressings substantially increase the percutaneous absorption of topical corticosteroids. Thus, occlusive dressings may be a valuable therapeutic adjunct for treatment of resistant dermatoses (see **DOSAGE AND ADMINISTRATION**). Once absorbed through the skin, topical corticosteroids are handled through pharmacokinetic pathways similar to systemically administered corticosteroids. Corticosteroids are bound to plasma proteins in varying degrees. Corticosteroids are metabolized primarily in the liver and are then excreted by the kidneys. Some of the topical corticosteroids and their metabolites are also excreted into the bile.

INDICATIONS AND USAGE

Topical corticosteroids are indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses.

CONTRAINDICATIONS

Topical corticosteroids are contraindicated in those patients with a history of hypersensitivity to any of

the components of the preparations.

PRECAUTIONS

General -

Systemic absorption of topical corticosteroids has produced reversible hypothalamic-pituitary-adrenal (HPA) axis suppression, manifestations of Cushing's syndrome, hyperglycemia, and glucosuria in some patients.

Conditions which augment systemic absorption include the application of the more potent steroids, use over large surface areas, prolonged use, and the addition of occlusive dressings. Therefore, patients receiving a large dose of a potent topical steroid applied to a large surface area or under an occlusive dressing should be evaluated periodically for evidence of HPA axis suppression by using the urinary free cortisol and ACTH stimulation tests. If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent steroid.

Recovery of HPA axis function is generally prompt and complete upon discontinuation of the drug. Infrequently, signs and symptoms of steroid withdrawal may occur, requiring supplemental systemic corticosteroids. Children may absorb proportionally larger amounts of topical corticosteroids and thus be more susceptible to systemic toxicity (see **PRECAUTIONS-Pediatric Use**).

If irritation develops, topical corticosteroids should be discontinued and appropriate therapy instituted. In the presence of dermatological infections, the use of an appropriate antifungal or antibacterial agent should be instituted. If a favorable response does not occur promptly, the corticosteroid should be discontinued until the infection has been adequately controlled.

Information for Patients

Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes.
2. Patients should be advised not to use this medication for any disorder other than for which it was prescribed.
3. The treated skin area should not be bandaged or otherwise covered or wrapped as to be occlusive unless directed by the physician.
4. Patients should report any signs of local adverse reactions especially under occlusive dressing.
5. Parents of pediatric patients should be advised not to use tight-fitting diapers or plastic pants on a child being treated in the diaper area, as these garments may constitute occlusive dressings.

Laboratory Tests

The following tests may be helpful in evaluating the HPA axis suppression:

urinary free cortisol test and ACTH stimulation test.

Carcinogenesis, Mutagenesis, Impairment of Fertility -

Long-term animal studies have not been performed to evaluate the carcinogenic potential or the effect on fertility of topical corticosteroids. Studies to determine mutagenicity with prednisolone and hydrocortisone have revealed negative results.

Pregnancy:

Teratogenic Effects:

Pregnancy Category C -

Corticosteroids are generally teratogenic in laboratory animals when administered systemically at relatively low dosage levels. The more potent corticosteroids have been shown to be teratogenic after dermal application in laboratory animals. There are no adequate and well-controlled studies in pregnant women on the teratogenic effects from topically applied corticosteroids. Therefore, topical corticosteroids should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Drugs of this class should not be used extensively on pregnant patients, in large amounts, or for prolonged periods of time.

Nursing Mothers -

It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in breast milk. Systemically administered corticosteroids are secreted into breast milk in quantities not likely to have a deleterious effect on the infant. Nevertheless, caution should be exercised when topical corticosteroids are administered to a nursing woman.

Pediatric Use -

Pediatric patients may demonstrate greater susceptibility to topical corticosteroid-induced HPA axis suppression and Cushing's syndrome than mature patients because of a larger skin surface area to body weight ratio.

Hypothalamic-pituitary-adrenal (HPA) axis suppression, Cushing's syndrome and intracranial hypertension have been reported in children receiving topical corticosteroids. Manifestations of adrenal suppression in children include linear growth retardation, delayed weight gain, low plasma cortisol levels, and absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

Administration of topical corticosteroids to children should be limited to the least amount compatible with an effective therapeutic regimen. Chronic corticosteroid therapy may interfere with the growth and development of children.

ADVERSE REACTIONS

The following local adverse reactions are reported infrequently with topical corticosteroids, but may occur more frequently with the use of occlusive dressings. These reactions are listed in an approximate decreasing order of occurrence: burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, maceration of the skin, secondary infection, skin atrophy, striae, and miliaria.

OVERDOSAGE

Topically applied corticosteroids can be absorbed in sufficient amounts to produce systemic effects (see **PRECAUTIONS**).

DOSAGE AND ADMINISTRATION

Topical corticosteroids are generally applied to the affected area as a thin film from two to four times daily depending on the severity of the condition.

Occlusive dressing may be used for the management of psoriasis or recalcitrant conditions. If an infection develops, the use of occlusive dressing should be discontinued and appropriate antimicrobial therapy instituted.

HOW SUPPLIED

Product: 53002-0150

NDC: 53002-0150-1 15 g in a TUBE

Product: 53002-9121

NDC: 53002-9121-1 15 g in a TUBE

NDC: 53002-9121-2 80 g in a TUBE

STORAGE

Store at 20-25°C (68-77°F) [see USP Controlled Room Temperature].

Manufactured By Perrigo

Bronx, NY 10457

Distributed By Perrigo

Allegan, MI 49010 • www.perrigo.com

Rev 05-15

: 5H100 RC JX1

Triamcinolone Acetonide Ointment 0.5%

Product Label for Triamcinolone Acetonide Ointment 0.5%:

- Top Left:** NDC# 53002-0150-1, LIST# 45802-048-35, **TRIAMCINOLONE ACETONIDE OINTMENT 0.5%**, 15 Gm Tube, LOT# 13719-412, EXP DATE: 05-31-2019, Shape & Color, Markings, Rx only.
- Top Center:** 15 Gm Tube, PATIENT NAME: _____, **TRIAMCINOLONE 0.5% OINTMENT**, PERRIGO, RxF# 173650591000 ITEM# 0150 015-43.
- Center:** **APPLY TO AFFECTED AREA 2-4 TIMES A DAY OR AS DIRECTED.**
- Bottom Left:** CLINIC NAME GOES HERE, IMPORTANT: THIS DRUG IS A CORTICOSTEROID FOR EXTERNAL USE ONLY. USE ONLY AS DIRECTED BY YOUR PRESCRIBER.
- Bottom Center:** Prescriber Name: _____, Date Dispensed: _____, **173650591000** (vertical barcode).
- Bottom Right:** APP AFF AREA 2-4 TIMES/DAY OR LO, LOT# 17279-012, EXP 06-31-2019, RxF# 173650591-000 FCA-0150, 15 Gm TRIAMCINOLONE 0.5% OINTMENT, **TRIAMCINOLONE ACETONIDE OINTMENT 0.5%**, 15 Gm Tube, DISCARD BY 05-31-2019, NDC# 53002-0150-1, RxF# 173650591-000, BILLING NDC# 45802-048-35, RxF# 173650591-000, 15 Gm TRIAMCINOLONE 0.5% OINTMENT, **15 Gm TRIAMCINOLONE 0.5% OINTMENT** (vertical text), RxF# 173650591000 ITEM# 0150.

Triamcinolone Acetonide Ointment 0.1%, USP

Product Label for Triamcinolone Acetonide Ointment 0.1%, USP:

- Top Left:** NDC# 53002-9121-1, LIST# 45802-055-35, **TRIAMCINOLONE ACETONIDE OINTMENT 0.1%, USP**, 15 Gm Tube, LOT# 18912-444, EXP DATE: 09-30-2019, Shape & Color, Markings, Rx only.
- Top Center:** 15 Gm Tube, PATIENT NAME: _____, **TRIAMCINOLONE 0.1% OINTMENT**, PERRIGO, RxF# 182961041000 ITEM# 9121 912-43.
- Center:** **APPLY TO AFFECTED AREA 2-4 TIMES A DAY OR AS DIRECTED.**
- Bottom Left:** CLINIC NAME GOES HERE, IMPORTANT: THIS DRUG IS A CORTICOSTEROID FOR EXTERNAL USE ONLY. USE ONLY AS DIRECTED BY YOUR PRESCRIBER.
- Bottom Center:** Prescriber Name: _____, Date Dispensed: _____, **182961041000** (vertical barcode).
- Bottom Right:** APP AFF AREA 2-4 X'S/DAY OR LO, LOT# 18062-080, EXP 09-30-2019, RxF# 182961041-000 FCA-9121, 15 Gm TRIAMCINOLONE 0.1% OINTMENT, **TRIAMCINOLONE ACETONIDE OINTMENT 0.1%, USP**, 15 Gm Tube, DISCARD BY 09-30-2019, NDC# 53002-9121-1, RxF# 182961041-000, BILLING NDC# 45802-055-35, RxF# 182961041-000, 15 Gm TRIAMCINOLONE 0.1% OINTMENT, **15 Gm TRIAMCINOLONE 0.1% OINTMENT** (vertical text), RxF# 182961041000 ITEM# 9121.

TRIAMCINOLONE ACETONIDE

triamcinolone acetonide ointment

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:53002-0150(NDC:45802-049)
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TRIAMCINOLONE ACETONIDE (UNII: F446C597KA) (TRIAMCINOLONE ACETONIDE - UNII:F446C597KA)	TRIAMCINOLONE ACETONIDE	5 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
PETROLATUM (UNII: 4T6H12BN9U)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53002-0150-1	15 g in 1 TUBE; Type 0: Not a Combination Product	10/01/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA087385	11/10/2006	

TRIAMCINOLONE ACETONIDE

triamcinolone acetonide ointment

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:53002-9121(NDC:45802-055)
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TRIAMCINOLONE ACETONIDE (UNII: F446C597KA) (TRIAMCINOLONE ACETONIDE - UNII:F446C597KA)	TRIAMCINOLONE ACETONIDE	1 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

LIGHT MINERAL OIL (UNII: N6K5787QVP)

PETROLATUM (UNII: 4T6H12BN9U)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53002-9121-1	15 g in 1 TUBE; Type 0: Not a Combination Product	10/01/2018	
2	NDC:53002-9121-2	80 g in 1 TUBE; Type 0: Not a Combination Product	10/01/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA087385	08/04/2006	

Labeler - RPK Pharmaceuticals, Inc. (147096275)

Establishment

Name	Address	ID/FEI	Business Operations
RPK Pharmaceuticals, Inc.		147096275	RELABEL(53002-9121, 53002-0150) , REPACK(53002-0150, 53002-9121)

Revised: 2/2020

RPK Pharmaceuticals, Inc.