

**LEADER DAYTIME SEVERE COLD AND FLU- acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution
Cardinal Health 110, LLC. dba Leader**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Cardinal Health Daytime Severe Cold & Flu Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Maximum Strength | Non-Drowsy

Daytime Severe Cold & Flu

Acetaminophen | Dextromethorphan HBr

Guaifenesin | Phenylephrine HCl

Pain Reliever / Fever Reducer

Cough Suppressant | Expectorant

Nasal Decongestant

Original Flavor

Relief of:

Aches, Fever, Cough, Sore Throat, Nasal Congestion, Chest Congestion

Alcohol-Free

COMPARE TO VICKS® DAYQUIL® SEVERE active ingredients

SEE NEW WARNINGS

8 FL OZ (237 mL)

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

© 2016 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.



DISTRIBUTED BY CARDINAL HEALTH
DUBLIN, OHIO 43017
www.myleader.com 1-800-200-6316
Essential to Care™ since 1979



Return to place of purchase if not satisfied.



LEADER²

NDC 70000-0192-1

Maximum Strength | Non-Drowsy

Daytime Severe Cold & Flu

Acetaminophen | Dextromethorphan HBr
Guaifenesin | Phenylephrine HCl

Pain Reliever / Fever Reducer
Cough Suppressant | Expectorant
Nasal Decongestant

Original Flavor

Relief of:
Aches, Fever, Cough,
Sore Throat,
Nasal Congestion,
Chest Congestion
Alcohol-Free

COMPARE TO
VICKS[®] DAYQUIL[®]
SEVERE
active ingredients*

100% Money
Back Guarantee

SEE NEW WARNINGS

8 FL OZ (237 mL)

Drug Facts

Active ingredients (in each 15 mL)

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- sinus congestion and pressure
 - minor aches and pains
 - reduces swelling of nasal passages
 - promotes nasal and/or sinus drainage
 - cough due to minor throat and bronchial irritation
 - temporarily restores freer breathing through the nose
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
 - nasal congestion
 - headache
 - sore throat
 - fever

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if
- adult takes more than 4,000 mg of acetaminophen in 24 hours
 - child takes more than 5 doses in 24 hours
 - taken with other drugs containing acetaminophen
 - adult has 3 or more alcoholic drinks every day while using this product

PEEL BACK AT CORNER FOR MORE INFORMATION

Drug Facts (continued)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)

Drug Facts (continued)

- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments? 1-800-719-9260

Gluten Free

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks[®] and DayQuil[®].

LEADER DAYTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0192
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL

DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A)	
SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM (UNII: 3980JIH2SW)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	ORANGE (clear)	Score	
Shape		Size	
Flavor	FRUIT, MENTHOL	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0192-1	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/23/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	11/23/2016	

Labeler - Cardinal Health 110, LLC. dba Leader (063997360)