

MOORE MEDICAL ANTACID- calcium carbonate tablet, chewable
Moore Medical LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Moore Medical Antacid

Drug Facts

Active ingredient (in each tablet)

Calcium Carbonate 420 mg

Purpose

Antacid

Uses

For the relief of the following symptoms

- acid indigestion
- sour stomach
- heartburn
- upset stomach associated with these symptoms

Warnings

Do not use

- the maximum dosage of this product for more than 2 weeks except under the advice and supervision of a physician or take more than 19 tablets in a 24 hour period.

Ask a doctor or pharmacist if you are

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if

- symptoms last more than 2 weeks

If pregnant or breast feeding,

ask a health professional before use.

Keep out of reach of children.

Directions

- do not use more than directed
- chew or crush tablets completely before swallowing
- do not swallow tablets whole

Adults and children: (12 years and older)

Chew 2 tablets every 2 or 3 hours as symptoms occur or as directed by a physician. Do not take more than 19 tablets in 24 hours.

Children under 12 years:

Do not give to children under 12 years of age.

Other information

- Phenylketonurics: contains phenylalanine 1.5 mg per tablet
- each tablet contains 168 mg of elemental calcium
- store at room temperature 59-86°F (15-30°C) in a dry place
- tamper-evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

aspartame*, croscarmellose sodium*, gum acacia*, magnesium stearate, maltodextrin, mineral oil*, mint flavor, sorbitol*, sucrose*

**may contain*

Questions or comments?

Call 1-800.234.1464

Moore Medical Antacid Label

moore medical

Supporting Health &Care

Reorder No.: 87675

UPN# 0-0607415-87675-8

Moore Brand®

Antacid Tablets

250 Packets/2 Tablets

Tablets for acid indigestion, gas, heartburn and acid stomach

Chew or crush tablets completely before swallowing.

Do not swallow tablets whole.

To Reorder:

Call: 1.800.234.1464

Fax: 1.800.944.6667

Online: www.mooremedical.com

 Supporting Health & Care	 Supporting Health & Care	Retain carton for complete product information <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th colspan="2" style="text-align: left;">Drug Facts</th> <th style="text-align: right;">Purpose</th> </tr> <tr> <td colspan="2">Active ingredient (in each tablet)</td> <td style="text-align: right;">Antacid</td> </tr> <tr> <td colspan="3">Caution: Contains 429 mg.</td> </tr> <tr> <td colspan="3">Uses</td> </tr> <tr> <td colspan="3">For the relief of the following symptoms</td> </tr> <tr> <td> <ul style="list-style-type: none"> ■ acid indigestion ■ heartburn ■ upset stomach associated with these symptoms </td> <td colspan="2"></td> </tr> <tr> <td colspan="3">Warnings</td> </tr> <tr> <td colspan="3">Do not use</td> </tr> <tr> <td colspan="3"> <ul style="list-style-type: none"> ■ if the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a physician, or take more than 10 tablets in a 24-hour period. </td> <td colspan="2"></td> </tr> <tr> <td colspan="3"> Ask a doctor or pharmacist before use if you are <ul style="list-style-type: none"> ■ already using a prescription drug. Antacids may interact with certain prescription drugs. </td> <td colspan="2"></td> </tr> <tr> <td colspan="3"> Stop use and ask a doctor if <ul style="list-style-type: none"> ■ symptoms last more than 2 weeks. ■ if pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. </td> <td colspan="2"></td> </tr> <tr> <td colspan="3">Directions</td> <td colspan="2"></td> </tr> <tr> <td colspan="3"> <ul style="list-style-type: none"> ■ do not use more than directed ■ chew or crush tablets completely before swallowing ■ do not swallow tablets whole </td> <td colspan="2"></td> </tr> <tr> <td colspan="3"> Adults and children: Chew 2 tablets every 2 or 3 hours as symptoms occur or as directed by a physician. Do not exceed 10 (12 years and older) tablets in 24 hours. </td> <td colspan="2"></td> </tr> <tr> <td colspan="3"> Children under 12 years: Do not give to children under 12 years of age. </td> <td colspan="2"></td> </tr> <tr> <td colspan="3">Other information</td> <td colspan="2"></td> </tr> <tr> <td colspan="3"> <ul style="list-style-type: none"> ■ Each tablet contains 150 mg of elemental calcium ■ Store at room temperature (20-25°C) (68-77°F) in a dry place ■ Do not use any opened or torn packets </td> <td colspan="2"> <ul style="list-style-type: none"> ■ each tablet contains 168 mg of elemental calcium ■ tamper-evident sealed packets </td> </tr> <tr> <td colspan="3">Inactive ingredients</td> <td colspan="2"></td> </tr> <tr> <td colspan="3">aspartame, croscarmellose sodium, gum arabic, magnesium stearate, maltodextrin, mineral oil, microcrystalline cellulose, sorbitol, xanthan gum</td> <td colspan="2"></td> </tr> <tr> <td colspan="3">*may contain</td> <td colspan="2"></td> </tr> <tr> <td colspan="3">Questions or comments: Call 800.234.1464.</td> <td colspan="2"></td> </tr> </table>	Drug Facts		Purpose	Active ingredient (in each tablet)		Antacid	Caution: Contains 429 mg.			Uses			For the relief of the following symptoms			<ul style="list-style-type: none"> ■ acid indigestion ■ heartburn ■ upset stomach associated with these symptoms 			Warnings			Do not use			<ul style="list-style-type: none"> ■ if the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a physician, or take more than 10 tablets in a 24-hour period. 					Ask a doctor or pharmacist before use if you are <ul style="list-style-type: none"> ■ already using a prescription drug. Antacids may interact with certain prescription drugs. 					Stop use and ask a doctor if <ul style="list-style-type: none"> ■ symptoms last more than 2 weeks. ■ if pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. 					Directions					<ul style="list-style-type: none"> ■ do not use more than directed ■ chew or crush tablets completely before swallowing ■ do not swallow tablets whole 					Adults and children: Chew 2 tablets every 2 or 3 hours as symptoms occur or as directed by a physician. Do not exceed 10 (12 years and older) tablets in 24 hours.					Children under 12 years: Do not give to children under 12 years of age.					Other information					<ul style="list-style-type: none"> ■ Each tablet contains 150 mg of elemental calcium ■ Store at room temperature (20-25°C) (68-77°F) in a dry place ■ Do not use any opened or torn packets 			<ul style="list-style-type: none"> ■ each tablet contains 168 mg of elemental calcium ■ tamper-evident sealed packets 		Inactive ingredients					aspartame, croscarmellose sodium, gum arabic, magnesium stearate, maltodextrin, mineral oil, microcrystalline cellulose, sorbitol, xanthan gum					*may contain					Questions or comments: Call 800.234.1464.					 Supporting Health & Care
Drug Facts		Purpose																																																																																										
Active ingredient (in each tablet)		Antacid																																																																																										
Caution: Contains 429 mg.																																																																																												
Uses																																																																																												
For the relief of the following symptoms																																																																																												
<ul style="list-style-type: none"> ■ acid indigestion ■ heartburn ■ upset stomach associated with these symptoms 																																																																																												
Warnings																																																																																												
Do not use																																																																																												
<ul style="list-style-type: none"> ■ if the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a physician, or take more than 10 tablets in a 24-hour period. 																																																																																												
Ask a doctor or pharmacist before use if you are <ul style="list-style-type: none"> ■ already using a prescription drug. Antacids may interact with certain prescription drugs. 																																																																																												
Stop use and ask a doctor if <ul style="list-style-type: none"> ■ symptoms last more than 2 weeks. ■ if pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. 																																																																																												
Directions																																																																																												
<ul style="list-style-type: none"> ■ do not use more than directed ■ chew or crush tablets completely before swallowing ■ do not swallow tablets whole 																																																																																												
Adults and children: Chew 2 tablets every 2 or 3 hours as symptoms occur or as directed by a physician. Do not exceed 10 (12 years and older) tablets in 24 hours.																																																																																												
Children under 12 years: Do not give to children under 12 years of age.																																																																																												
Other information																																																																																												
<ul style="list-style-type: none"> ■ Each tablet contains 150 mg of elemental calcium ■ Store at room temperature (20-25°C) (68-77°F) in a dry place ■ Do not use any opened or torn packets 			<ul style="list-style-type: none"> ■ each tablet contains 168 mg of elemental calcium ■ tamper-evident sealed packets 																																																																																									
Inactive ingredients																																																																																												
aspartame, croscarmellose sodium, gum arabic, magnesium stearate, maltodextrin, mineral oil, microcrystalline cellulose, sorbitol, xanthan gum																																																																																												
*may contain																																																																																												
Questions or comments: Call 800.234.1464.																																																																																												
Reorder No. 87675 UPN# 0-0607415-87675-8 MooreBrand® Antacid Tablets 250 Packets/2 Tablets Tablets for acid indigestion, gas, heartburn and acid stomach. Chew or crush tablets completely before swallowing. Do not swallow tablets whole.	Reorder No. 87675 UPN# 0-0607415-87675-8 MooreBrand® Antacid Tablets 250 Packets/2 Tablets Tablets for acid indigestion, gas, heartburn and acid stomach. Chew or crush tablets completely before swallowing. Do not swallow tablets whole.	Reorder No. 87675 UPN# 0-0607415-87675-8 MooreBrand® Antacid Tablets 250 Packets/2 Tablets Tablets for acid indigestion, gas, heartburn and acid stomach. Chew or crush tablets completely before swallowing. Do not swallow tablets whole.																																																																																										
Distributed by: Moore Medical LLC Farmington, CT 06032 Manufacturer: 21305178 Made in USA	To Reorder: Call: 800.234.1464 Fax: 800.344.5667 Online: www.mooremedical.com	Distributed by: Moore Medical LLC Farmington, CT 06032 Manufacturer: 21305178 Made in USA	To Reorder: Call: 800.234.1464 Fax: 800.344.5667 Online: www.mooremedical.com																																																																																									

MOORE MEDICAL ANTACID

calcium carbonate tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55670-141
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK) (CALCIUM CATION - UNII:2M83C4R6ZB, CARBONATE ION - UNII:7UJQ50PE7D)	CALCIUM CARBONATE	420 mg

Inactive Ingredients

Ingredient Name	Strength
ASPARTAME (UNII: Z0H242BBR1)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MINERAL OIL (UNII: T5L8T28FGP)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	white (white)	Score	no score
Shape	ROUND (ROUND)	Size	11mm
Flavor		Imprint Code	AZ;036
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date

1	NDC:55670-141-13	250 in 1 BOX	07/11/2014	
1	NDC:55670-141-99	2 in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC monograph final	part331		07/11/2014	

Labeler - Moore Medical LLC (051420107)

Revised: 4/2019

Moore Medical LLC