

**MOTION SICKNESS LESS DROWSY FORMULA- meclizine hcl tablet
MEIJER, INC.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each tablet)

Meclizine HCl 25 mg

Purpose

Antiemetic

Uses

for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.

Warnings

Do not use

for children under 12 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to an enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- to prevent motion sickness, the first dose should be taken 1 hour before starting activity
- adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor
- do not exceed 2 tablets in 24 hours

Other information

- store at room temperature 20°-25°C (68°-77°F)

Inactive ingredients

anhydrous lactose, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminium lake, magnesium stearate, microcrystalline cellulose, sodium starch glycolate

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

Compare to Dramamine® Less Drowsy Formula active ingredient*

Less Drowsy Formula

MotionSicknessTabs

Meclizine HCl 25 mg/ Antiemetic

For Nausea And Dizziness From Travel

Tablets

*This product is not manufactured or distributed by MedTech Products Inc., owner of the registered trademark Dramamine® Less Drowsy Formula.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DIST. BY MEIJER DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

www.meijer.com

Product Label

MotionSickness Tabs

meijer

Less Drowsy Formula

NDC 41250-620-08

meijer

Compare to Dramamine®
Less Drowsy Formula
active ingredient*

Less Drowsy Formula

MotionSickness Tabs

Meclizine HCl 25 mg/ Antiemetic

For Nausea And Dizziness From Travel



8 Tablets

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Lot No.:
Exp. Date:

*This product is not manufactured or distributed by MedTech Products Inc., owner of the registered trademark Dramamine® Less Drowsy Formula.

DIST. BY MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com



PLD-D
FC001822

Drug Facts	Meclizine HCl 25 mg.....Antiemetic
Purpose	
Uses	for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.
Warnings	Do not use for children under 12 years of age unless directed by a doctor. Ask a doctor before use if you have ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ difficulty in urination due to an enlargement of the prostate gland Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. When using this product ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ use caution when driving a motor vehicle or operating machinery If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions	■ to prevent motion sickness, the first dose should be taken 1 hour before starting activity ■ adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor ■ do not exceed 2 tablets in 24 hours
Other information	■ store at room temperature 20°-25°C (68°-77°F)
Inactive ingredients	magnesium stearate, anhydrous lactose, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, microcrystalline cellulose, sodium starch glycolate
Questions or comments?	Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Meijer Less Drowsy Formula Motion Sickness Tablets

MOTION SICKNESS LESS DROWSY FORMULA

meclizine hcl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-620
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
ALUMINUM OXIDE (UNII: LMI26O6933)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	TCL086
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-620-08	8 in 1 CARTON	03/28/2014	03/28/2025
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part336	03/28/2014	03/28/2025

Labeler - MEIJER, INC. (006959555)

