

GUAIFENESIN- guaifenesin liquid
Innovida Pharmaceutique Corporation

Active Ingredient

Guaifenesin, USP 200 mg

Purpose

Expectorant

Uses

- helps loosen phlegm(mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

ASK A DOCTOR BEFORE USE IF

- cough that occurs with too much phlegm(mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

STOP USE AND ASK DOCTOR IF cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache.

IF PREGNANT OR BREAST-FEEDING, ask a health professional before use. **KEEP OUT OF REACH OF CHILDREN.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- shake well before using
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- ml=milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Other Information

Store between 20° to 25°C (68° to 77°F)

Inactive Ingredients

caramel color, citric acid anhydrous, FD&C red# 40 dye, flavor, glycerin, high fructose corn syrup, menthol, propylene glycol, purified water, saccharine sodium, sodium benzoate.

Questions or Comments? 1-888-462-4166

Product label

Drug Facts	
Active ingredient Guaifenesin, USP 200 mg	Purpose Expectorant
Uses(s) • helps loosen phlegm(mucus) and thin bronchial secretions to drain bronchial tubes	
Warnings ASK A DOCTOR BEFORE USE IF • cough that occurs with too much phlegm(mucus) • cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema	
STOP USE AND ASK DOCTOR IF cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache.	
IF PREGNANT OR BREAST-FEEDING , ask a health professional before use. KEEP OUT OF REACH OF CHILDREN. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions • shake well before using • do not take more than 6 doses in any 24-hour period • measure only with dosing cup provided • keep dosing cup with product • mL=milliliter • this adult product is not intended for use in children under 12 years of age	
age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Drug Facts (continued)

Other Information

- each 10 mL contains: sodium 12 mg
- store at 20-25°C(68-77°F)
- protect from light

Inactive ingredients
caramel color, citric acid anhydrous, FD&C red# 40 dye, flavor, glycerin, high fructose corn syrup, menthol, propylene glycol, purified water, saccharine sodium, sodium benzoate.

Questions or Comments? 1-888-462-4166

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.

Mfd. for: Innovida Pharmaceutique Corporation, Phoenix, AZ 85040
Tel: 888-462-4166



INN^oVIDA
PHARMACEUTIQUE CORPORATION

Guaifenesin
200 mg per 10 mL

Expectorant Loosens
and Relieves Chest
Congestion

NDC: 71800-051-05

INN^oVIDA
PHARMACEUTIQUE CORPORATION

Guaifenesin
200 mg per 10 mL

Expectorant Loosens
and Relieves Chest
Congestion

NDC: 71800-051-05

INN^oVIDA
PHARMACEUTIQUE CORPORATION

Guaifenesin
200 mg per 10 mL


Expectorant Loosens
and Relieves Chest
Congestion

NDC: 71800-051-05

LIQUID

4 FL OZ (118 mL)


Drug Facts
Active ingredient (in each 10 mL) **Purpose**
Guaifenesin, USP 200 m.....Expectorant
Uses(s) • helps loosen phlegm(mucus) and thin bronchial secretions to drain bronchial tubes
Warnings
ASK A DOCTOR BEFORE USE IF • cough that occurs with too much phlegm(mucus) • cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema
STOP USE AND ASK DOCTOR IF cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache.
IF PREGNANT OR BREAST-FEEDING, ask a health professional before use. **KEEP OUT OF REACH OF CHILDREN.** In case of overdose, get medical help or contact a Poison Control Center right away.

NDC: 71800-051-05

Guaifenesin
200 mg per 10 mL
Expectorant Loosens and Relieves Chest Congestion
LIQUID
4 FL OZ (118 mL)

Drug Facts (continued)
Directions
• shake well before using • do not take more than 6 doses in any 24-hour period • measure only with dosing cup provided • keep dosing cup with product • ml=milliliter • this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Other Information
• each 10 mL contains: sodium 12 mg
• store at 20-25oC(68-77oF)
• protect from light
Inactive ingredients
caramel color, citric acid anhydrous, FD&C red# 40 dye, flavor, glycerin, high fructose corn syrup, menthol, propylene glycol, purified water, saccharine sodium, sodium benzoate.
Questions or Comments? 1-888-462-4166



DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.
Mfd. for: Innovida Pharmaceutique Corporation, Phoenix, AZ 85040
Tel: 888-462-4166

GUAIFENESIN

guaifenesin liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71800-051
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL (UNII: L7T10EIP3A)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
CARAMEL (UNII: T9D99G2B1R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:71800-051-05	118 mL in 1 TUBE; Type 0: Not a Combination Product	02/10/2025	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		02/10/2025	

Labeler - Innovida Pharmaeutique Corporation (080892908)

Revised: 3/2025

Innovida Pharmaeutique Corporation