

**BERLEY AND JENSEN PAIN RELIEF- acetaminophen tablet, film coated
BJWC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

BJWC Pain Relief Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 caplets every 6 hours while symptoms last• do not take more than 6 caplets in 24 hours, unless directed by a doctor• do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store at 20-25°C (68-77°F)
- do not use if printed seal under cap is broken or missing

Inactive ingredients

croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, mica-based pearlescent pigment, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, stearic acid

Questions or comments?

1-800-934-1204

Package/Label Principal Display Panel

Compare to the active ingredient in Tylenol® Extra Strength Rapid Release Gels®

CONTAINS NO ASPIRIN

EXTRA STRENGTH

RAPID RELEASE

PAIN RELIEF

ACETAMINOPHEN

PAIN RELIEVER/FEVER REDUCER

GLUTEN FREE

FOR ADULTS

400 CAPLETS | 500 mg EACH

ACTUAL SIZE

Drug Facts (continued)

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warnings: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions do not take more than directed (see overdose warning)

adults and children 12 years and over	take 2 caplets every 6 hours while symptoms last
	do not take more than 6 caplets in 24 hours, unless directed by a doctor
children under 12 years	do not use for more than 10 days unless directed by a doctor
	ask a doctor

Other information store at 20-25°C (68-77°F) do not use if printed seal under cap is broken or missing

Inactive ingredients croscarmellose sodium, croscopolidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, mica-based pearlescent pigment, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, stearic acid

Questions or comments? 1-800-934-1204

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Tylenol® Extra Strength Rapid Release Gels.

Distributed by: BJ's Wholesale Club, 25 Research Drive, Westborough, MA 01581 Questions? 1-800-934-1204

100% MONEY-BACK GUARANTEE Every Berley Jensen product comes with our 100% Money-Back Guarantee.

Compare to the active ingredient in Tylenol® Extra Strength Rapid Release Gels®

berley jensen

CONTAINS NO ASPIRIN

EXTRA STRENGTH RAPID RELEASE PAIN RELIEF ACETAMINOPHEN PAIN RELIEVER/FEVER REDUCER FOR ADULTS 400 | 500 CAPLETS | mg EACH

ACTUAL SIZE

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg	Pain reliever/fever reducer
----------------------	-----------------------------

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - minor pain of arthritis
 - headache
 - backache
 - toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours or with other drugs containing acetaminophen or 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- new symptoms occur
- pain gets worse or lasts more than 10 days

CODE AREA

8867003184

35074 D3 F2

BERLEY AND JENSEN PAIN RELIEF

acetaminophen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:6839 1-004
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients	
Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (15 MPAS AT 5%) (UNII: 68401960MK)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Product Characteristics

Color	RED	Score	no score
Shape	OVAL	Size	18mm
Flavor		Imprint Code	3S0
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68391-004-79	400 in 1 BOTTLE; Type 0: Not a Combination Product	11/30/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	11/30/2015	

Labeler - BJWC (159082692)