

**NITETIME DAYTIME COLD AND FLU- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl**  
**Meijer Distribution Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Meijer Distribution, Inc. NiteTime DayTime Cold & Flu Drug Facts**

**Nighttime Severe Cold & Flu**  
**Active ingredients (in each 30 mL)**

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Doxylamine succinate 12.5 mg

Phenylephrine HCl 10 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

**Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and sneezing
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed - see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- **each 30 mL contains:** sodium 41 mg
- store at 20-25°C (68-77°F)

**Inactive ingredients**

anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

## Questions or comments?

**1-800-719-9260**

### **Daytime Severe Cold & Flu**

#### **Active ingredients (in each 15 mL)**

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

### **Purpose**

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

### **Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

### **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

### **When using this product**

**do not use more than directed**

### **Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present

- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed - see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- **each 15 mL contains:** sodium 6 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients**

butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

**Questions or comments?**

**1-800-719-9260**

**Principal Display Panel - Nighttime**

DAYTIME & NITETIME COMBO PACK

Compare to Vicks® NyQuil® Severe and DayQuil® Severe active ingredients

meijer®

Compare to Vicks® DayQuil® Severe active ingredients

meijer®

MAXIMUM STRENGTH

NON-DROWSY

daytime cold & flu

Acetaminophen

Pain Reliever | Fever Reducer

Dextromethorphan HBr

Cough Suppressant

Guaifenesin | Expectorant

Phenylephrine HCl

Nasal Decongestant

SEVERE

Alcohol Free

Antihistamine Free

Compare to Vicks® NyQuil® Severe active ingredients

meijer®

MAXIMUM STRENGTH

nighttime cold & flu

Acetaminophen

Pain Reliever | Fever Reducer

Dextromethorphan HBr

Cough Suppressant

Doxylamine Succinate

Antihistamine

Phenylephrine HCl

Nasal Decongestant

SEVERE

Mixed Berry Flavor

Sugar Free | Alcohol Free

Multi-Symptom Relief

2 - 12 FL OZ (355 mL) BOTTLES

TOTAL 24 FL OZ (1.5 pt) (710 mL)





meijer

MAXIMUM STRENGTH  
NON-DROWSY

daytime  
cold & flu

SEVERE

Multi-Symptom Relief

meijer

MAXIMUM STRENGTH

nitetime  
cold & flu

SEVERE

Multi-Symptom Relief

Mixed Berry Flavor

DAYTIME & NITETIME  
COMBO PACK

NDC 41250-597-02

meijer

Compare to Vicks<sup>®</sup>  
NyQuil<sup>®</sup> Severe and  
DayQuil<sup>®</sup> Severe  
active ingredients\*

meijer

Compare to Vicks<sup>®</sup>  
NyQuil<sup>®</sup> Severe  
active ingredients\*

MAXIMUM  
STRENGTH

nitetime  
cold & flu

SEVERE

Multi-Symptom  
Relief

Mixed Berry Flavor  
Sugar Free | Alcohol Free



DO NOT USE IF PRINTED NECKBAND  
IS BROKEN OR MISSING

DIST. BY MEIJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com



meijer

Compare to Vicks<sup>®</sup>  
DayQuil<sup>®</sup> Severe  
active ingredients\*

MAXIMUM  
STRENGTH  
NON-DROWSY

daytime  
cold & flu

Acetaminophen  
Pain Reliever | Fever Reducer  
Dextromethorphan HBr  
Cough Suppressant  
Guaifenesin | Expectorant  
Phenylephrine HCl  
Nasal Decongestant

SEVERE

Alcohol Free  
Antihistamine Free



meijer

Compare to Vicks<sup>®</sup>  
NyQuil<sup>®</sup> Severe  
active ingredients\*

MAXIMUM  
STRENGTH

nitetime  
cold & flu

Acetaminophen  
Pain Reliever | Fever Reducer  
Dextromethorphan HBr  
Cough Suppressant  
Doxylamine Succinate  
Antihistamine  
Phenylephrine HCl  
Nasal Decongestant

SEVERE

Mixed Berry Flavor  
Sugar Free | Alcohol Free

Multi-Symptom Relief



2-12 FL OZ (355 mL) BOTTLES  
TOTAL 24 FL OZ (1.5 pt) (710 mL)

PD 3416660



7 0882041630 7

59702 6E C5



Nighttime Severe Cold & Flu

Drug Facts	Purpose
<b>Active ingredients (in each 30 mL)</b>	
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Doxymetamine succinate 12.5 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal decongestant

**Uses:** temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and an itchy
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores their breathing through the nose

**Warnings:**  
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:  
 • more than 4,000 mg of acetaminophen in 24 hours  
 • with other drugs containing acetaminophen  
 • 3 or more alcoholic drinks every day while using this product  
 Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:  
 • skin reddening • blisters • rash  
 If a skin reaction occurs, stop use and seek medical help right away.  
 Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use:**  
 • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 • if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have:**  
 • liver disease • heart disease • high blood pressure  
 • thyroid disease • diabetes • glaucoma  
 • cough that occurs with too much phlegm (mucus)  
 • a breathing problem such as emphysema or chronic bronchitis  
 • trouble urinating due to an enlarged prostate gland  
 • persistent or chronic cough such as occurs with smoking, asthma, or emphysema  
 • a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are:**  
 • taking sedatives or tranquilizers • taking the blood thinning drug warfarin

**When using this product:**  
 • do not use more than directed • excitability may occur, especially in children  
 • marked drowsiness may occur • avoid alcoholic drinks  
 • be careful when driving a motor vehicle or operating machinery  
 • alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if:**  
 • you get nervous, dizzy or sleepless  
 • pain, nasal congestion, or cough gets worse or lasts more than 7 days  
 • fever gets worse or lasts more than 3 days  
 • redness or swelling is present • new symptoms occur  
 • cough comes back or occurs with rash or headache that lasts.  
 These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

take only as directed - see Overdose warning	
only use the dose cup provided	do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**  
 • each 30 mL contains sodium 41 mg  
 • store at 20-25°C (68-77°F)

**Inactive ingredients:** anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

**Questions or comments?** 1-800-719-9280

Daytime Severe Cold & Flu

Drug Facts	Purpose
<b>Active ingredients (in each 15 mL)</b>	
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses:** temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores their breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make cough more productive

**Warnings:**  
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if:  
 • adult takes more than 4,000 mg of acetaminophen in 24 hours  
 • child takes more than 5 doses in 24 hours  
 • taken with other drugs containing acetaminophen  
 • adult has 3 or more alcoholic drinks every day while using this product  
 Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:  
 • skin reddening • blisters • rash  
 If a skin reaction occurs, stop use and seek medical help right away.  
 Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use:**  
 • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 • if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have:**  
 • liver disease • heart disease • thyroid disease  
 • diabetes • high blood pressure  
 • trouble urinating due to an enlarged prostate gland  
 • cough that occurs with too much phlegm (mucus)  
 • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product do not use more than directed**

**Stop use and ask a doctor if:**  
 • you get nervous, dizzy or sleepless  
 • pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)  
 • fever gets worse or lasts more than 3 days  
 • redness or swelling is present  
 • new symptoms occur  
 • cough comes back or occurs with rash or headache that lasts.  
 These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

take only as directed - see Overdose warning	
only use the dose cup provided	do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**  
 • each 15 mL contains sodium 6 mg  
 • store at 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients:** butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, inositol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

**Questions or comments?** 1-800-719-9280



Compare to Vicks® DayQuil® Severe active ingredients\*

**MAXIMUM STRENGTH NON-DROWSY**

**daytime cold & flu**

**SEVERE**


**Multi-Symptom Relief**

**Alcohol Free Antihistamine Free**

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)



how2recycle.info

\*These products are not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks® NyQuil® and DayQuil®.

## NITETIME DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-597
---------------------	----------------	---------------------------	---------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-597-02	1 in 1 KIT; Type 0: Not a Combination Product	12/30/2013	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	355 mL
Part 2	1 BOTTLE	355 mL

## Part 1 of 2

### NITETIME COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl solution

### Product Information

<b>Item Code (Source)</b>	NDC:41250-763
---------------------------	---------------

<b>Route of Administration</b>	ORAL
--------------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>	RED (clear, dark)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-763-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/11/2013	

## Part 2 of 2

### DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution

### Product Information

<b>Item Code (Source)</b>	NDC:41250-603
<b>Route of Administration</b>	ORAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>BUTYLATED HYDROXYANISOLE</b> (UNII: REK4960K2U)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MENTHOL, UNSPECIFIED FORM</b> (UNII: L7T10EIP3A)	
<b>SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM</b> (UNII: 3980JIH2SW)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

**Product Characteristics**

<b>Color</b>	ORANGE (clear)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	FRUIT, MENTHOL	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-603-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/11/2013	

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final	part341	12/30/2013	

**Labeler** - Meijer Distribution Inc (006959555)

Revised: 9/2022

Meijer Distribution Inc