

**CHILDRENS IBUPROFEN- ibuprofen tablet, chewable**  
**Meijer Distribution Inc**

-----  
**Meijer Distribution, Inc. Children's Ibuprofen Drug Facts**

**Active ingredient (in each chewable tablet)**

Ibuprofen 100 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more

than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

### **Do not use**

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

*Ask a doctor or pharmacist before use if the child is*

- under a doctor's care for any serious condition
- taking any other drug

*When using this product*

- mouth or throat burning may occur; give with food or water
- take with food or milk if stomach upset occurs

*Stop use and ask a doctor if*

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
  - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days

- redness or swelling is present in the painful area
- any new symptoms appear

*Keep out of reach of children.*

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- **chew or crush tablets completely before swallowing**
- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**

<b>Dosing Chart</b>		
<b>Weight (lb)</b>	<b>Age (yr)</b>	<b>Tablets</b>
under 24	under 2	ask a doctor
24-35	2-3	1
36-47	4-5	1 ½
48-59	6-8	2
60-71	9-10	2 ½
72-95	11	3

## Other information

- phenylketonurics: contains phenylalanine 6 mg per tablet
- store between 20-25°C (68-77°F)
- **do not use if printed seal under cap is broken or missing**

## Inactive ingredients

acesulfame potassium, ammonium glycyrrhizin, aspartame, carnauba wax, croscarmellose sodium, D&C red no. 27 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, mannitol, natural and artificial flavors, silicon dioxide, sodium lauryl sulfate, soybean oil, succinic acid

## Questions or comments?

**1-800-719-9260**

## Package/Label Principal Display Panel

meijer®

Compare to Children's Motrin® active ingredient

For Ages 2 to 11

CHILDREN'S

ibuprofen

Chewable Tablets, 100 mg

CHEWABLES LASTS UP TO 8 HOURS

Pain Reliever | Fever Reducer (NSAID)

24 Tablets

Grape Flavored Chewables

Chew or Crush Tablets Completely Before Swallowing



PID 26282 19

KEEP CARTON FOR REFERENCE

\*\*This product is not manufactured or distributed by Johnson & Johnson, owner of the registered trademark, Motrin®.

**Important:** Read all product information before using. Keep this box for important information. This product is for children only.

**Drug Facts**

**Active ingredient (in each chewable tablet)** **Purposes**  
 ibuprofen 100 mg (NSAID)\* Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**Uses temporarily:** ■ reduce fever  
 ■ relieve minor aches and pains due to the common cold, flu, sore throat, headaches, and toothaches

**Warnings**

Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 ■ hives ■ facial swelling ■ asthma-like wheezing  
 ■ shock ■ skin redness ■ rash ■ hives  
 If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you child:  
 ■ has had stomach ulcers or bleeding problems  
 ■ takes a blood thinning (anticoagulant) or steroid drug  
 ■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 ■ takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Some throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 24 days or a maximum of 3 years of age unless directed by doctor.

**Drug Facts (continued)**

**Do not use**  
 ■ if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer  
 ■ right before or after heart surgery

**Ask a doctor before use if**

■ stomach bleeding warning applies to your child  
 ■ child has a history of stomach problems, such as heartburn  
 ■ child has problems or serious side effects from taking pain relievers or fever reducers  
 ■ child has not been drinking fluids  
 ■ child has lost a lot of fluid due to vomiting or diarrhea  
 ■ child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke  
 ■ child has asthma ■ child is taking a diuretic  
 ■ Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition  
 ■ taking any other drug

**When using this product:**

■ mouth or throat burning may occur: give with food or water  
 ■ take with food or milk if stomach upset occurs  
 ■ Stop use and ask a doctor if  
 ■ child experiences any of the following signs of stomach bleeding:  
 ■ tests faint ■ vomits blood ■ has bloody or black stools  
 ■ has stomach pain that does not get better  
 ■ child has symptoms of heart problems or stroke:  
 ■ chest pain ■ trouble breathing  
 ■ weakness in one part of side of body  
 ■ slurred speech ■ leg swelling

**Drug Facts (continued)**

■ the child does not get any relief within first day (24 hours) of treatment  
 ■ fever or pain gets worse or lasts more than 3 days  
 ■ redness or swelling is present in the part of ear  
 ■ any new symptoms appear

**Directions**

Chew or crush tablets completely before swallowing  
 ■ this product does not contain directions or complete warnings for adult use  
 ■ do not give more than directed  
 ■ first night dose on chart below. If possible, use weight to dose; otherwise use age.  
 ■ if needed, repeat doses every 6-8 hours  
 ■ do not use more than 4 times a day

**Dosing Chart**

Weight (lb)	Age (yr)	Tablets
under 24	under 2	ask a doctor
24-35	2-3	1
36-47	4-5	1½
48-59	6-8	2
60-71	9-10	2½
72-95	11	3

**Other information**

■ contains 24 tablets in a blister pack  
 ■ contains 24 tablets in a blister pack  
 ■ contains 24 tablets in a blister pack  
 ■ contains 24 tablets in a blister pack  
 ■ contains 24 tablets in a blister pack

**Questions or comments? 1-800-719-2600**

ibuprofen is a nonsteroidal anti-inflammatory drug (NSAID). It is used to relieve pain and reduce fever. It is also used to treat inflammation. It is not recommended for children under 2 years of age. It is not recommended for children 2 to 11 years of age unless directed by a doctor. It is not recommended for children 12 years of age and older unless directed by a doctor. It is not recommended for children with certain medical conditions. It is not recommended for children taking certain medications. It is not recommended for children with certain allergies. It is not recommended for children with certain bleeding disorders. It is not recommended for children with certain kidney or liver problems. It is not recommended for children with certain heart problems. It is not recommended for children with certain stomach problems. It is not recommended for children with certain blood pressure problems. It is not recommended for children with certain asthma problems. It is not recommended for children with certain seizure problems. It is not recommended for children with certain diabetes problems. It is not recommended for children with certain thyroid problems. It is not recommended for children with certain hormone problems. It is not recommended for children with certain cancer problems. It is not recommended for children with certain infection problems. It is not recommended for children with certain other medical conditions.

DIST. BY MEIJER DISTRIBUTION, INC.  
 GRAND RAPIDS, MI 49544  
 www.meijer.com



NDC 41250-726-62

Compare to Children's Motrin® active ingredient\*\*

**meijer**

For Ages 2 to 11

**CHILDREN'S ibuprofen**

Cheewable Tablets, 100 mg

**CHEWABLES**

**LASTS UP TO 8 HOURS**

**24 Tablets**

**Pain Reliever | Fever Reducer (NSAID)**

Grape Flavored Chewables

Chew or Crush Tablets Completely Before Swallowing

ibuprofen tablet, chewable

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-726
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>AMMONIUM GLYCYRRHIZATE</b> (UNII: 3VRD35U26C)	
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>SOYBEAN OIL</b> (UNII: 241ATL177A)	
<b>SUCCINIC ACID</b> (UNII: AB6MNQ6J6L)	

## Product Characteristics

<b>Color</b>	PURPLE (Lavender)	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	12mm
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	L521
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:41250-726-62	1 in 1 CARTON	07/18/2019	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing</b>	<b>Application Number or Monograph</b>	<b>Marketing Start</b>	<b>Marketing End</b>
------------------	--	------------------------	----------------------

Category	Citation	Date	Date
ANDA	ANDA076359	07/18/2019	

**Labeler** - Meijer Distribution Inc (006959555)

Revised: 10/2022

Meijer Distribution Inc