

**WINCO FOODS EFFERVESCENT COLD RELIEF- aspirin, chlorpheniramine maleate, and phenylephrine bitartrate tablet, effervescent**  
**WinCo Foods, LLC**

-----  
**Effervescent Cold Relief Tablets**

**Active ingredients**

*Aspirin 325 mg (NSAID)\**

*Chlorpheniramine maleate 2 mg*

*Phenylephrine Bitartrate 7.8 mg*

*\*Nonsteroidal anti-inflammatory drug*

**Purpose**

**Uses**

temporarily relieves these symptoms of the common cold:

- fever ■ sneezing ■ nasal congestion ■ sore throat ■ headache
- minor aches and pains ■ runny nose
- sinus congestion and pressure

**Uses**

temporarily relieves these symptoms of the common cold

- fever ■ sneezing ■ nasal congestion ■ sore throat ■ headache
- minor aches and pains ■ runny nose
- sinus congestion and pressure

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more

alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you are allergic to aspirin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if**

- stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic. ■ you are taking sedatives or tranquilizers.
- You have** ■ glaucoma ■ diabetes ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ been placed on a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

- presently taking a prescription drug ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

### **When using this product**

do not take more than 8 tablets (adults and children 12 years and over) in a 24-hour period or as directed by doctor.

- do not use more than directed ■ you may get drowsy ■ avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding
- feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better

■ an allergic reaction occurs. Seek medical help right away. ■ pain or nasal congestion gets worse or lasts more than 7 days

■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present

■ ringing in the ears or a loss of hearing occurs ■ nervousness, dizziness or sleeplessness occurs

### **If pregnant or breast-feeding,**

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.**

**Keep out of reach of children.**

### **In case of overdose,**

get medical help or contact a Poison Control Center right away.

### **Directions**

■ do not use more than directed (see overdose warning)

■ adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4-6 hours

■ do not take more than 8 tablets in 24 hours

■ children under 12 years: ask a doctor

### **Other information**

#### **each tablet contains:**

sodium 464 mg

■ phenylketonurics: contains phenylalanine 9 mg per tablet

■ store at room temperature (59°-86°F)

### **Inactive ingredients**

acesulfame potassium, aspartame, citric acid, docusate sodium, flavors, mannitol, povidone, sodium

benzoate, sodium bicarbonate

**Questions?**

Or to report an adverse event call

800-824-1706 Monday - Friday

9am - 4pm MST

**PACKAGE LABEL. PRINCIPAL DISPLAY PANEL**



# WINCO FOODS EFFERVESCENT COLD RELIEF

aspirin, chlorpheniramine maleate, and phenylephrine bitartrate tablet, effervescent

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67091-109
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>PHENYLEPHRINE BITARTRATE</b> (UNII: 27O3Q5ML57) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>DOCUSATE SODIUM</b> (UNII: F05Q2T2JA0)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	25mm
<b>Flavor</b>	LEMON	<b>Imprint Code</b>	CF
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-109-20	10 in 1 CARTON	07/02/2014	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug M012

07/02/2014

**Labeler** - WinCo Foods, LLC (056098817)

Revised: 11/2024

WinCo Foods, LLC