

**COLD AND FLU RELIEF- acetaminophen, dextromethorphan hydrobromide,  
phenylephrine hydrochloride liquid  
Gobrand, Inc. (Goodnow)**

-----

**Drug Facts**

**Active ingredients (in each 15 mL)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Purpose**

**Pain reliever/fever reducer**

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves common cold and flu symptoms:
  - minor aches and pains
  - headache
  - fever
  - sore throat
  - nasal congestion
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- adult take more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 4 doses (15 mL each) in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blistere
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- a sodium-restricted diet
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if user is**

taking the blood thinning drug warfarin

**When using this product**

**do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause serious liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-

800-222-1222) right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms

### Directions

- **do not take more than directed (see overdose warning)**
- do not take more than 4 doses in 24-hours period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

### Other information

- each 15 mL contains: sodium 12 mg
- store between 20°-25°C (68°-77°F). Do not refrigerate.

### Inactive ingredients

citric acid, FD&C Yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol sucralose, xanthan gum

### Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

### Principal Display Panel

Compare to the active ingredients in Vicks® DayQuil® Cold & Flu†

NON-DROWSY

### Cold & Flu

### Day Relief

Acetaminophen

Dextromethorphan HBr

Phenylephrine HCl

Pain Reliever  
 Fever Reducer  
 Cough Suppressant  
 Nasal Decongestant  
 Alcohol-free / Antihistamine-free  
 Multi-Symptom Relief  
 Headache & Fever  
 Sore Throat  
 Minor Aches & Pains  
 Nasal Congestion  
 Cough  
 FL OZ (mL)

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

**DISTRIBUTED BY: GOBRANDS, INC.**

**PHILADELPHIA, PA 19123**

**Product Label**

<p><b>Drug Facts (continued)</b></p> <p>When using this product, do not exceed recommended dosage.</p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>● nervousness, dizziness, or sleeplessness occur</li> <li>● pain, nasal congestion, or cough gets worse, or lasts more than 5 days (children) or 7 days (adult)</li> <li>● new symptoms occur</li> <li>● fever gets worse, or lasts more than 3 days</li> <li>● redness or swelling is present</li> <li>● cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.</li> </ul> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. <b>Overdose warning:</b> Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>● do not take more than directed (see Overdose warning)</li> <li>● do not take more than 4 doses in any 24-hour period</li> <li>● measure only with dosing cup provided. Do not use any other dosing device.</li> <li>● mL = milliliter</li> <li>● keep dosing cup with product</li> </ul> <table border="1"> <tr> <td>adults and children 12 years and over</td> <td>30 mL every 4 hours</td> </tr> <tr> <td>children 6 to under 12 years</td> <td>15 mL every 4 hours</td> </tr> <tr> <td>children 4 to under 6 years</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 years</td> <td>do not use</td> </tr> </table> <p>● when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing</p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>● each 15 mL contains: sodium 12 mg</li> <li>● store between 20°-25°C (68°-77°F). Do not refrigerate.</li> </ul> <p><b>Inactive ingredients</b> citric acid, FD&amp;C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum</p> <p><b>Questions or comments?</b> Call 1-877-763-3935 Monday-Friday 9AM-5PM EST</p> <p><small>*This product is not manufactured or distributed by The Procter &amp; Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter &amp; Gamble Company.</small></p> <p>DISTRIBUTED BY: GOBRANDS, INC.    PHILADELPHIA, PA 19123    ©2022 GOBRANDS, INC.</p> <p>PLD-A380D LB008799</p> <p><b>PARENTS:</b>    Learn about teen medicine abuse  <a href="http://www.StopMedicineAbuse.org">www.StopMedicineAbuse.org</a></p> <p>TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.</p>	adults and children 12 years and over	30 mL every 4 hours	children 6 to under 12 years	15 mL every 4 hours	children 4 to under 6 years	ask a doctor	children under 4 years	do not use	<p>Compare to active ingredients in Vicks® DayQuil® Cold &amp; Flu†</p> <p>NDC 82501-4661-2</p> <p><b>good now™</b></p> <p>NON-DROWSY</p> <p><b>Cold + Flu Day Relief</b></p> <p>Acetaminophen    Dextromethorphan HBr    Phenylephrine HCl</p> <p>Pain Reliever    Fever Reducer    Cough Suppressant    Nasal Decongestant</p> <p>Alcohol-free / Antihistamine-free</p> <p>Multi-Symptom Relief</p> <p>Headache &amp; Fever    Sore Throat    Minor Aches &amp; Pains    Nasal Congestion    Cough</p> <p>12 FL OZ (355 mL)</p>	<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each 15 mL)</b></p> <p>Acetaminophen 325 mg.....Pain reliever/fever reducer    Dextromethorphan HBr 10 mg.....Cough suppressant    Phenylephrine HCl 5 mg.....Nasal decongestant</p> <p><b>Purposes</b></p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>● temporarily relieves common cold and flu symptoms</li> <li>● minor aches and pains</li> <li>● headache</li> <li>● fever</li> <li>● sore throat</li> <li>● nasal congestion</li> <li>● cough due to minor throat and bronchial irritation</li> </ul> <p><b>Warnings</b></p> <p><b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if:</p> <ul style="list-style-type: none"> <li>● adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount</li> <li>● child takes more than 4 doses (15 mL each) in 24 hours</li> <li>● taken with other drugs containing acetaminophen</li> <li>● adult has 3 or more alcoholic drinks every day while using this product.</li> </ul> <p><b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: ● skin reddening ● blisters ● rash. If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>● with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>● if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> </ul> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>● liver disease</li> <li>● heart disease</li> <li>● high blood pressure</li> <li>● thyroid disease</li> <li>● diabetes</li> <li>● a sodium-restricted diet</li> <li>● trouble urinating due to an enlarged prostate gland</li> <li>● persistent or chronic cough such as occurs with smoking, asthma, or emphysema</li> <li>● cough that occurs with too much phlegm (mucus)</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</b></p>
adults and children 12 years and over	30 mL every 4 hours									
children 6 to under 12 years	15 mL every 4 hours									
children 4 to under 6 years	ask a doctor									
children under 4 years	do not use									

**GOOD NOW Non-Drowsy Cold + Flu Day Relief**

# COLD AND FLU RELIEF

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:82501-4661
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82501-4661-2	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/29/2022	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/29/2022	

Revised: 6/2024

Gobrand, Inc. (Goodnow)