

**COLD AND FLU RELIEF- acetaminophen, dextromethorphan hydrobromide,
phenylephrine hydrochloride liquid
Gobrand, Inc. (Goodnow)**

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - minor aches and pains
 - headache
 - fever
 - sore throat
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- adult take more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 4 doses (15 mL each) in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- a sodium-restricted diet
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if user is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause serious liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-

800-222-1222) right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms

Directions

- **do not take more than directed (see overdose warning)**
- do not take more than 4 doses in 24-hours period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

Other information

- each 15 mL contains: sodium 12 mg
- store between 20°-25°C (68°-77°F). Do not refrigerate.

Inactive ingredients

citric acid, FD&C Yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredients in Vicks® DayQuil® Cold & Flu†

NON-DROWSY

Cold & Flu

Day Relief

Acetaminophen

Dextromethorphan HBr

Phenylephrine HCl

Pain Reliever
 Fever Reducer
 Cough Suppressant
 Nasal Decongestant
 Alcohol-free / Antihistamine-free
 Multi-Symptom Relief
 Headache & Fever
 Sore Throat
 Minor Aches & Pains
 Nasal Congestion
 Cough
 FL OZ (mL)

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: GOBRANDS, INC.

PHILADELPHIA, PA 19123

Product Label

<p>Drug Facts (continued)</p> <p>When using this product, do not exceed recommended dosage.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ● nervousness, dizziness, or sleeplessness occur ● pain, nasal congestion, or cough gets worse, or lasts more than 5 days (children) or 7 days (adult) ● new symptoms occur ● fever gets worse, or lasts more than 3 days ● redness or swelling is present ● cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition. <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <ul style="list-style-type: none"> ● do not take more than directed (see Overdose warning) ● do not take more than 4 doses in any 24-hour period ● measure only with dosing cup provided. Do not use any other dosing device. ● mL = milliliter ● keep dosing cup with product <table border="1"> <tr> <td>adults and children 12 years and over</td> <td>30 mL every 4 hours</td> </tr> <tr> <td>children 6 to under 12 years</td> <td>15 mL every 4 hours</td> </tr> <tr> <td>children 4 to under 6 years</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 years</td> <td>do not use</td> </tr> </table> <p>● when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing</p> <p>Other information</p> <ul style="list-style-type: none"> ● each 15 mL contains sodium 12 mg ● store between 20°-25°C (68°-77°F). Do not refrigerate. <p>Inactive ingredients citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum</p> <p>Questions or comments? Call 1-877-763-3935 Monday-Friday 9AM-5PM EST</p> <p><small>*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.</small></p> <p>DISTRIBUTED BY: GOBRANDS, INC. PHILADELPHIA, PA 19123 ©2022 GOBRANDS, INC.</p> <p>TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.</p>	adults and children 12 years and over	30 mL every 4 hours	children 6 to under 12 years	15 mL every 4 hours	children 4 to under 6 years	ask a doctor	children under 4 years	do not use	<p>Compare to active ingredients in Vicks® DayQuil® Cold & Flu†</p> <p>NDC 82501-4661-2</p> <p>good now™</p> <p>NON-DROWSY</p> <p>Cold + Flu Day Relief</p> <p>Acetaminophen Dextromethorphan HBr Phenylephrine HCl</p> <p>Pain Reliever Fever Reducer Cough Suppressant Nasal Decongestant</p> <p>Alcohol-free / Antihistamine-free</p> <p>12 FL OZ (355 mL)</p> <p>Multi-Symptom Relief</p> <p>Headache & Fever Sore Throat Minor Aches & Pains Nasal Congestion Cough</p>	<p>Drug Facts</p> <p>Active ingredients (in each 15 mL)</p> <p>Acetaminophen 325 mg.....Pain reliever/fever reducer Dextromethorphan HBr 10 mg.....Cough suppressant Phenylephrine HCl 5 mg.....Nasal decongestant</p> <p>Uses</p> <ul style="list-style-type: none"> ● temporarily relieves common cold and flu symptoms ● minor aches and pains ● headache ● fever ● sore throat ● nasal congestion ● cough due to minor throat and bronchial irritation <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if:</p> <ul style="list-style-type: none"> ● adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount ● child takes more than 4 doses (15 mL each) in 24 hours ● taken with other drugs containing acetaminophen ● adult has 3 or more alcoholic drinks every day while using this product. <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ● skin reddening ● blisters ● rash. If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> ● with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ● if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ● liver disease ● heart disease ● high blood pressure ● thyroid disease ● diabetes ● a sodium-restricted diet ● trouble urinating due to an enlarged prostate gland ● persistent or chronic cough such as occurs with smoking, asthma, or emphysema ● cough that occurs with too much phlegm (mucus) <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p>
adults and children 12 years and over	30 mL every 4 hours									
children 6 to under 12 years	15 mL every 4 hours									
children 4 to under 6 years	ask a doctor									
children under 4 years	do not use									

GOOD NOW Non-Drowsy Cold + Flu Day Relief

COLD AND FLU RELIEF

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82501-4661
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SORBITOL (UNII: 506T60A25R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
XANTHAN GUM (UNII: TTV12P4NEE)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82501-4661-2	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/29/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/29/2022	

Revised: 4/2024

Gobrand, Inc. (Goodnow)