

VITAMINS A, C, D AND FLUORIDE DROPS- vitamin a palmitate, ascorbic acid, cholecalciferol, and sodium fluoride solution/ drops
H2-Pharma, LLC

Vitamins A, C, D and Fluoride Drops

Supplement Facts		
Serving Size 1.0 mL		
Servings Per Container 50		
	Amount Per Serving	% Daily Value
Children over 4 & Adults		
Vitamin A	450 mcg	50%
Vitamin C	35 mg	39%
Vitamin D	10 mcg (400 IU)	50%
Fluoride	0.5 mg	*

* Daily Value not established

*U.S. Recommended Daily Allowance not established.


Consult your physician for use by infants and young children 6 months to 3 years of age.

Active ingredient for caries prophylaxis: Fluoride as sodium fluoride.

Other ingredients: Ascorbic acid, caramel color, cherry flavor, cholecalciferol, citric acid, ferrous sulfate, glycerin, methyl paraben, polysorbate 80, purified water, sodium benzoate, sodium fluoride, sodium hydroxide, sucralose, vitamin A palmitate.

CLINICAL PHARMACOLOGY

It is well established that fluoridation of the water supply (0.7 mg/L fluoride) during the period of tooth development leads to a significant decrease in the incidence of dental caries. Hydroxyapatite is the principal crystal for all calcified tissue in the human body. The fluoride ion reacts with the hydroxyapatite in the tooth as it is formed to produce the more caries-resistant crystal, fluorapatite. The reaction may be expressed by the equation:



Three stages of fluoride deposition in tooth enamel can be distinguished:

1. Small amounts (reflecting the low levels of fluoride in tissue fluids) are incorporated into the enamel crystals while they are being formed.
2. After enamel has been laid down, fluoride deposition continues in the surface enamel.

Diffusion of fluoride from the surface inward is apparently restricted.

3. After eruption, the surface enamel acquires fluoride from water, food, supplementary fluoride and small amounts from saliva.

Vitamins A, C, D and Fluoride Drops 0.5 mg provide supplementation of the diet with three essential vitamins, and with fluoride for caries prophylaxis. The American Academy of Pediatrics recommends that children up to age 16, in areas where drinking water contains less than optimal levels of fluoride, receive daily fluoride supplementation. Vitamins A, C, D and Fluoride Drops 0.5 mg provide fluoride in drop form for infants and young children 6 months to 3 years of age in areas where the drinking water contains less than 0.3 ppm of fluoride and for children ages 3 to 6 years in areas where the drinking water contains 0.3 through 0.6 ppm of fluoride.

Each 1.0 mL dose of Vitamins A, C, D and Fluoride Drops 0.5 mg supplies sodium fluoride (0.5 mg fluoride) plus significant amounts of vitamins A, C, and D in a single easy-to-use preparation.

WARNINGS

As in the case of all medications, keep out of the reach of children.

PRECAUTIONS

The suggested dose should not be exceeded since dental fluorosis may result from continued ingestion of large amounts of fluoride. When recommending vitamin fluoride products:

1. Determine the fluoride content of the drinking water.
2. Make sure the child is not receiving significant amounts of fluoride from other medications and swallowed toothpaste.
3. Periodically check to make sure that the child does not develop significant dental fluorosis.

ADVERSE REACTIONS

Allergic rash or other idiosyncrasies have been rarely reported. To report SUSPECTED ADVERSE REACTIONS, contact H2-Pharma at 1-866-592-6438 or FDA at 1-800-FDA-1088 or via the web at www.fda.gov/medwatch for voluntary reporting of adverse reactions.

DOSAGE AND ADMINISTRATION

As directed by a physician. See American Dental Association (ADA) recommended supplement schedule below. May be dropped directly into the mouth with the enclosed dropper or mixed with cereal, fruit juice or other food.

ADA Dietary Fluoride Supplement Schedule for Children at High Caries Risk

Age	Fluoride ion level in drinking water
------------	---

Age	(ppm)*		
	<0.3 ppm	0.3-0.6 ppm	>0.6 ppm
Birth - 6 months	None	None	None
6 months - 3 years	0.25 mg/day†	None	None
3-6 years	0.50 mg/day	0.25 mg/day	None
6-16 years	1.0 mg/day	0.50 mg/day	None

* 1.0 part per million (ppm) = 1 milligram/liter (mg/L)

† 2.2 mg sodium fluoride contains 1 mg fluoride ion.

SHAKE WELL BEFORE USE.

HOW SUPPLIED

Vitamins A, C, D and Fluoride Drops 0.5 mg are available in 50 mL bottles with accompanying calibrated dropper.

Dispense in original container.

RECOMMENDED STORAGE

Store at controlled room temperature, 20°-25°C (66°-77°F) [See USP Controlled Room Temperature]. After opening, store away from direct sunlight. Close tightly after each use. REFRIGERATION IS NOT REQUIRED.

TAMPER EVIDENT: Do not use if seal under cap is torn, broken or missing.

Distributed by: **H2-Pharma, LLC**
Montgomery, AL 36117

PRINCIPAL DISPLAY PANEL - 50 mL Bottle Carton

61269-167-50

Drops

Vitamins
A, C, D
and Fluoride

Supplement Drops

1²/₃ fl. oz. (50 mL)

0.5 mg

H² pharma

61269-167-50



Drops

Vitamins A, C, D and Fluoride

Supplement Drops

1²/₃ fl. oz. (50 mL)

0.5 mg



Supplement Facts

Serving Size 1.0 mL
Servings Per Container 50

	Amount Per Serving	% Daily Value
		<small>Children over 4 & Adults</small>
Vitamin A	450 mcg	50%
Vitamin C	35 mg	39%
Vitamin D	10 mcg (400 IU)	50%
Fluoride	0.5 mg	*

*Daily Value not established

* U.S. Recommended Daily Allowance not established.

Consult your physician for use by infants and young children 6 months to 3 years of age.

Active ingredient for caries prophylaxis:

Fluoride as sodium fluoride.

Other ingredients: Ascorbic acid, caramel color, cherry flavor, cholecalciferol, citric acid, ferrous sulfate, glycerin, methyl paraben, polysorbate 80, purified water, sodium benzoate, sodium fluoride, sodium hydroxide, sucralose, vitamin A palmitate.

Distributed by: H2-Pharma, LLC
Montgomery, AL 36117
www.h2-pharma.com






0.5 mg
 1/2 fl. oz. (50 mL)
**Vitamins A, C, D
 and Fluoride
 Supplement Drops**

CLINICAL PHARMACOLOGY

It is well established that fluoridation of the water supply (0.7 mg/L fluoride) during the period of tooth development leads to a significant decrease in the incidence of dental caries. Hydroxyapatite is the principal crystal for all calcified tissue in the human body. The fluoride ion reacts with the hydroxyapatite in the tooth as it is formed to produce the more caries-resistant crystal, fluorapatite. The reaction may be expressed by the equation:



Three stages of fluoride deposition in tooth enamel can be distinguished:

1. Small amounts (reflecting the low levels of fluoride in tissue fluids) are incorporated into the enamel crystals while they are being formed.
2. After enamel has been laid down, fluoride deposition continues in the surface enamel. Diffusion of fluoride from the surface inward is apparently restricted.
3. After eruption, the surface enamel acquires fluoride from water, food, supplementary fluoride and small amounts from saliva.

Vitamins A, C, D and Fluoride Drops 0.5 mg provide supplementation of the diet with three essential vitamins, and with fluoride for caries prophylaxis. The American Academy of Pediatrics recommends that children up to age 16, in areas where drinking water contains less than optimal levels of fluoride, receive daily fluoride supplementation. Vitamins A, C, D and Fluoride Drops 0.5 mg provide fluoride in drop form for infants and young children 6 months to 3 years of age in areas where the drinking water contains less than 0.3 ppm of fluoride and for children ages 3 to 6 years in areas where the drinking water contains 0.3 through 0.6 ppm of fluoride.

Each 1.0 mL dose of Vitamins A, C, D and Fluoride Drops 0.5 mg supplies sodium fluoride (0.5 mg fluoride) plus significant amounts of vitamins A, C, and D in a single easy-to-use preparation.

WARNINGS As in the case of all medications, keep out of the reach of children.

PRECAUTIONS The suggested dose should not be exceeded since dental fluorosis may result from continued ingestion of large amounts of fluoride. When recommending vitamin fluoride products:

1. Determine the fluoride content of the drinking water.
2. Make sure the child is not receiving significant amounts of fluoride from other medications and swallowed toothpaste.
3. Periodically check to make sure that the child does not develop significant dental fluorosis.

ADVERSE REACTIONS Allergic rash or other idiosyncrasies have been rarely reported. To report SUSPECTED ADVERSE REACTIONS, contact H2-Pharma at 1-866-592-6438 or FDA at 1-800-FDA-1088 or via the web at www.fda.gov/medwatch for voluntary reporting of adverse reactions.

DOSAGE AND ADMINISTRATION: As directed by a physician. See American Dental Association (ADA) recommended supplement schedule below. May be dropped directly into the mouth with the enclosed dropper or mixed with cereal, fruit juice or other food.

Age	Fluoride ion level in drinking water (ppm)*		
	<0.3 ppm	0.3-0.6 ppm	>0.6 ppm
Birth - 6 months	None	None	None
6 months - 3 years	0.25 mg/day**	None	None
3-6 years	0.50 mg/day	0.25 mg/day	None
6-16 years	1.0 mg/day	0.50 mg/day	None

* 1.0 part per million (ppm) = 1 milligram/liter (mg/L)
 ** 2.2 mg sodium fluoride contains 1 mg fluoride ion.

SHAKE WELL BEFORE USE.

HOW SUPPLIED Vitamins A, C, D and Fluoride Drops 0.5 mg are available in 50 mL bottles with accompanying calibrated dropper.

Dispense in original container.

RECOMMENDED STORAGE Store at controlled room temperature, 20°-25°C (66°-77°F) (See USP Controlled Room Temperature). After opening, store away from direct sunlight. Close tightly after each use. REFRIGERATION IS NOT REQUIRED.

TAMPER EVIDENT: Do not use if seal under cap is



1800063[02]

VITAMINS A, C, D AND FLUORIDE DROPS

vitamin a palmitate, ascorbic acid, cholecalciferol, and sodium fluoride solution/ drops

Product Information

Product Type	DIETARY SUPPLEMENT	Item Code (Source)	NHRIC:61269-167
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
VITAMIN A PALMITATE (UNII: 1D1K0N0VVC) (VITAMIN A - UNII:81G40H8B0T)	VITAMIN A	450 [iU] in 1 mL
ASCORBIC ACID (UNII: PQ6CK8PD0R) (ASCORBIC ACID - UNII:PQ6CK8PD0R)	ASCORBIC ACID	35 mg in 1 mL
CHOLECALCIFEROL (UNII: 1C6V77QF41) (CHOLECALCIFEROL - UNII:1C6V77QF41)	CHOLECALCIFEROL	10 [iU] in 1 mL
SODIUM FLUORIDE (UNII: 8ZYQ1474W7) (FLUORIDE ION - UNII:Q80VPU4080)	SODIUM FLUORIDE	0.5 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
CARAMEL (UNII: T9D99G2B1R)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FERROUS SULFATE (UNII: 39R4TAN1VT)	
CHERRY (UNII: BUC5I9595W)	
GLYCERIN (UNII: PDC6A3C0OX)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NHRIC:61269-167-50	1 in 1 CARTON		

1	50 mL in 1 BOTTLE		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
DIETARY SUPPLEMENT		10/22/2018	05/31/2026

Supplement Facts	
Serving Size :	Serving per Container :
Amount Per Serving	% Daily Value
color	
flavor	

Labeler - H2-Pharma, LLC (028473634)

Revised: 9/2025

H2-Pharma, LLC