

IBUPROFEN - ibuprofen capsule, liquid filled
Walgreen Company

Active ingredient(s)

(in each capsule)

Solubilized Ibuprofen equal to
200 mg ibuprofen (NSAID)*

(present as the free acid and potassium salt)

* nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Use(s)

- temporarily relieves minor aches and pain due to :
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alerts: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen,

or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because Ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask doctor if

- you experience any of following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that dose not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in painful area
- any new symptoms appear

Pregnancy/Breastfeeding

ask a health professional before use. It is especially important not to use Ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause a problems in the unborn child or complications during delivery.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison ControlCenter right away.(1-800-222-1222)

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever dose not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use.
- store at 20-25 °C (68-77 °F)
- avoid excessive heat above 40 °C (104 °F)

Inactive ingredients

Ammonium hydroxide, FD&C green no.3, Gelatin, Iron oxide black, Medium chain triglycerides, Polyethylene glycol, Propylene glycol, Potassium hydroxide, Purified water, Shellac, Sorbitol, Sorbitan monooleate

Principal Display Panel



Drug Facts (continued)

Some side effects may include: **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 65 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or over-the-counter NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed

The risk is higher if you: take a blood thinner are taking a diuretic you have kidney problems, such as heart failure you have high blood pressure, heart disease, liver disease, kidney disease or asthma

Ask a doctor or pharmacist before use if you are: under a doctor's care for any serious condition taking aspirin or heart attack or stroke, because ibuprofen may decrease the benefit of aspirin taking any other drug

NO COAT

NO COAT

Drug Facts

Active ingredient (in each capsule): Solifenilopantolol mesylate (NSAID)
200 mg (aspirin (NSAID))
(aspirin is the trade name for aspirin)
Nonsteroidal anti-inflammatory drug

Uses: temporarily relieves minor aches and pains due to: headache toothache backache menstrual cramps the common cold sinus pain sprain muscle aches minor pain of arthritis

Warnings: Many side effects may occur. Stop use and seek medical help right away if: an allergic reaction occurs, such as rash, hives, or swelling you experience any of the following signs of stomach bleeding: black or bloody stools vomit that looks like coffee grounds stomach pain that does not get better

Stop use and ask a doctor if: you experience any of the following signs of stomach bleeding: vomit that looks like coffee grounds stomach pain that does not get better you experience any of the following signs of stomach bleeding: vomit that looks like coffee grounds stomach pain that does not get better

When using this product: stay with food or milk if stomach upset occurs the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

NO COAT

NO COAT

Drug Facts (continued)

Other information:

- Each capsule contains potassium 20 mg
- Read all warnings and directions before use. Keep carton.
- Store between 20°-25°C (68°-77°F)
- Avoid excessive heat above 40°C (104°F)

Inactive ingredients: Acetaminophen hydrochloride, FD&C green no. 3, Gelatin, Iron oxide black, Medium chain triglycerides, Polyethylene glycol, Povidone, Potassium hydroxide, Purified water, Shellac, Sorbitol sorbitate, Sorbitol monohydrate

Questions or comments? 1-877-896-7679

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Walgreens

Compare to Advil® Liqui-gels® active ingredient*

NDC 0363-0003-20

IBUPROFEN

Ibuprofen Capsules 200 mg
Pain Reliever/Fever Reducer (NSAID)

SEE NEW WARNINGS INFORMATION

Liquid Filled

20 SOFTGELS**
(**Liquid filled capsules)

SOFTGELS**

DO NOT USE IF SEAL UNDER BOTTLE CAP INDICATED WITH REDDASH FROM YOUR PHARMACEUTICAL IS BROKEN OR MISSING

This product is not recommended for distribution by Walgreens Consumer Healthcare, owner of the registered trademark Advil®



ITEM 157164

3 1191714367 5

DISTRIBUTED BY: WALGREEN CO.
200 WILMOTT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
Walgreens.com ©2012 Walgreen Co.

Lot No.:
Exp. Date:

20ct_Carton



ITEM 157164

Active ingredient (in each capsule)
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*...Pain reliever/fever reducer (present as the free acid and potassium salt) *nonsteroidal anti-inflammatory drug

Purpose
Temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever

Uses
Temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever

Warnings
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.
Allergy alert
Liver warning
Stomach bleeding warning
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
■ do not take more than directed
■ the smallest effective dose should be used
■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
■ if pain or fever does not respond to 1 capsule, 2 capsules may be used
■ do not exceed 6 capsules in 24 hours, unless directed by a doctor
■ children under 12 years: ask a doctor

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2012 Walgreen Co.

Lot No.:

Exp Date:

20ct_Label

Questions or Comments? 1-877-586-7979

IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0003
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
AMMONIA (UNII: 5138Q19F1X)	
FD&C GREEN NO. 3 (UNII: 3P3ONR601S)	
GELATIN (UNII: 2G86QN327L)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0K00R)	
SHELLAC (UNII: 46N107B710)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	

Product Characteristics

Color	GREEN	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	133
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0003-20	1 in 1 CARTON		
1		20 in 1 BOTTLE		
2	NDC:0363-0003-40	1 in 1 CARTON		
2		40 in 1 BOTTLE		
3	NDC:0363-0003-80	1 in 1 CARTON		
3		80 in 1 BOTTLE		
4	NDC:0363-0003-16	1 in 1 CARTON		
4		160 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079205	05/09/2011	

Labeler - Walgreen Company (008965063)**Establishment**

Name	Address	ID/FEI	Business Operations
Aphena Pharma Solutions - Kentucky, LLC.		965013498	repack

Establishment

Name	Address	ID/FEI	Business Operations
Marksans Pharma Limited		925822975	manufacture

Revised: 5/2012

Walgreen Company