

ANTIPHLAMINE- l-menthol, l-camphor, methyl salicylate lotion
Lydia Co., Ltd.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

L-Menthol, L-Camphor, Methyl Salicylate

■ temporary relief aches and pains of muscles and joints associated with:

- arthritis • simple backache
- sprains • strains • bruises

Keep out of reach of children

■ Adults and children 2 years of age and older: apply to the affected area no more than 3 to 4 times daily

■ children under 2 years of age: consult a doctor

Warnings

For external use only.

Flammable

■ Keep away from fire, flame, sparks and heated surfaces. Tight cap firmly.

Stop use and ask a doctor if

■ Condition worsens or symptoms persist for more than 7days or clear up and occur again within a few days

When using this product

- Use only as direct
- Avoid contact with eyes
- Do not apply to wounds or damaged skin
- Do not bandage tightly

If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children

acetone, chloroxylenol, water

For external use only



| ANTIPHLAMINE LOTION | |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------|
| <i>Composition Active Ingredients (in 1g)</i> | <i>Purpose</i> |
| L-Menthol 8mg ----- | Topical analgesic |
| L-Camphor 59mg ----- | Topical analgesic |
| Methyl Salicylate 46mg ----- | Topical analgesic |
| Uses | |
| <ul style="list-style-type: none"> ■ for temporary relief aches and pains of muscles and joints associated with: <ul style="list-style-type: none"> • arthritis • simple backache • sprains • strains • bruises | |
| Warnings | |
| For external use only. | |
| Flammable | |
| <ul style="list-style-type: none"> ■ Keep away from fire, flame, sparks and heated surfaces. Tight cap firmly. | |
| Stop use and ask a doctor if | |
| <ul style="list-style-type: none"> ■ Condition worsens or symptoms persist for more than 7days or clear up and occur again within a few days | |
| When using this product | |
| <ul style="list-style-type: none"> ■ Use only as direct ■ Avoid contact with eyes ■ Do not apply to wounds or damaged skin ■ Do not bandage tightly | |
| If pregnant or breast-feeding, ask a health professional before use | |
| Keep out of reach of children. Store at room temperature | |
| Directions | |
| <ul style="list-style-type: none"> ■ Adults and children 2 years of age and older: apply to the affected area no more than 3 to 4 times daily ■ children under 2 years of age: consult a doctor | |
| Inactive Ingredient | |
| acetone, chloroxylenol, water | |
| Distributed By:Bio Mission Group,Inc/562-273-0692 | |
| Made in South Korea | |

ANTIPHLAMINE

I-menthol, I-camphor, methyl salicylate lotion

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:72988-0028 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--------------------------------------------------------------------------------|--------------------------|-----------------|
| CAMPHOR, (-)- (UNII: 213N3S8275) (CAMPHOR, (-)- - UNII:213N3S8275) | CAMPHOR, (-)- | 59 mg in 1 g |
| LEVOMENTHOL (UNII: BZ1R15MTK7) (LEVOMENTHOL - UNII:BZ1R15MTK7) | LEVOMENTHOL | 8 mg in 1 g |
| METHYL SALICYLATE (UNII: LAV5U5022Y) (SALICYLIC ACID - UNII:O414PZ4LPZ) | METHYL SALICYLATE | 46 mg in 1 g |

Inactive Ingredients

| Ingredient Name | Strength |
|-----------------------------------------|-----------------|
| ACETONE (UNII: 1364PS73AF) | |
| CHLOROXYLENOL (UNII: 0F32U78V2Q) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|----------|------------------|--------------------------------------------------|-----------------------------|---------------------------|
| 1 | NDC:72988-0028-1 | 30 g in 1 JAR; Type 0: Not a Combination Product | 01/02/2022 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------------|-------------------------------------------------|-----------------------------|---------------------------|
| unapproved drug other | | 01/02/2022 | |

Labeler - Lydia Co., Ltd. (695735569)

Registrant - Lydia Co., Ltd. (695735569)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-----------------|----------------|---------------|----------------------------|
| Lydia Co., Ltd. | | 695735569 | manufacture(72988-0028) |

Revised: 7/2023

Lydia Co., Ltd.