

PRO-SAN LV BASE- sodium chlorite liquid IBA

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Pro-San LV Base

Active Ingredient

Sodium Chlorite 0.7%

Mixing Directions

To make an RTU solution of Pro-San LV Teat Dip, add equal volumes of Pro-San LV Base and Pro-San LV Activator into a clean dip cup/container and mix until the color is uniform throughout. Do not dilute. Do not mix with any other products. Mix only the amount needed for a 24 hour period. Discard any unused teat dip. Unused teat dip may be diluted with water and safely flushed down the drain.

Recommended Procedures for Teat Sanitation

A. PRE-MILKING (PRE-DIPPING)

1. Remove visible soils from the teats
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip, spray or foam the cow's teats full length with READY-TO-USE PRO-SAN LV TEAT DIP.
4. Wait at least 15-30 seconds.
5. Remove all teat dip by thoroughly drying the teats, teat orifices and udder with a clean, single service paper towel(s) to avoid contamination of milk.
6. Attach milking unit.

B. POST-MILKING (POST-DIPPING)

Immediately after milking, dip, spray or foam all teats full length with READY-TO-USE PRO-SAN LV TEAT DIP. Ensure good coverage of all sides of the teats and teat orifices. Allow to dry. Do not wipe. Do not turn cows out in freezing weather until teat dip is completely dry.

Caution

Keep out of reach of children. Avoid contact with food. Do not take internally. Not for human use.

First Aid

EYE CONTACT: Rinse cautiously with fresh, tepid water for several minutes. Remove

contact lenses if present and easy to do. Continue rinsing for 15 minutes. Get immediate medical advice/attention. **SKIN CONTACT:** For RTU products wash any sensitive areas. **INGESTION:** Do not induce vomiting. Never give anything by mouth to an unconscious person. Drink large amounts of water. If vomiting occurs, keep airways open. Keep head lower than the hips to prevent aspiration into the lungs. Get immediate medical advice/attention.

Warning

Do not dilute this product.

Do not return teat dip left in the teat dipper to storage container.

Ensure that dipped teats are dry before exposing the animals to weather conditions which may cause damage to the teats.

Store at room temperature. Always store away from direct sunlight.

Protect this product from freezing. However, freezing will not affect Pro-San LV provided that the contents are thawed completely and agitated thoroughly before mixing base and activator together.

Pro-San LV Base

NDC 29983-1092-8

**NPE
FREE**



**NPE
FREE**

PRO-SAN LV

SANITIZING PRE-MILKING AND POST-MILKING TEAT DIP

A Highly Effective Germicidal Non-Iodine Formulation

ACTIVE INGREDIENT:
Sodium Chlorite.....0.70%

GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.

CAUTION:
KEEP OUT OF REACH OF CHILDREN.
AVOID CONTACT WITH FOOD.
DO NOT TAKE INTERNALLY.
NOT FOR HUMAN USE.

FIRST AID PROCEDURES

EYE CONTACT: RINSE CAUTIOUSLY WITH FRESH, TEPID WATER FOR SEVERAL MINUTES. REMOVE CONTACT LENSES IF PRESENT AND EASY TO DO. CONTINUE RINSING FOR 15 MINUTES. GET IMMEDIATE MEDICAL ADVICE/ATTENTION.


SKIN CONTACT: FOR READY-TO-USE PRODUCTS, WASH ANY SENSITIVE AREAS.

INGESTION: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS, KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS. GET IMMEDIATE MEDICAL ADVICE/ATTENTION.

**ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION
FOR MEDICAL EMERGENCIES CALL 1-800-424-9300**

BASE

**FOR USE ONLY WITH
PRO-SAN LV
ACTIVATOR**



An Aid In The Prevention
Of New Intramammary
Infections

MIXING DIRECTIONS: TO MAKE A READY-TO-USE SOLUTION OF PRO-SAN LV TEAT DIP, ADD EQUAL VOLUMES OF PRO-SAN LV BASE AND PRO-SAN LV ACTIVATOR INTO A CLEAN DIP CUP/CONTAINER AND MIX UNTIL THE COLOR IS UNIFORM THROUGHOUT. DO NOT DILUTE. DO NOT MIX WITH ANY OTHER PRODUCTS. MIX ONLY THE AMOUNT NEEDED FOR A 24 HOUR PERIOD. DISCARD ANY UNUSED TEAT DIP. UNUSED TEAT DIP MAY BE DILUTED WITH WATER AND SAFELY FLUSHED DOWN THE DRAIN.

RECOMMENDED PROCEDURES FOR TEAT SANITATION

A. PRE-MILKING (PRE-DIPPING)

1. Remove visible soils from the teats.
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip, spray or foam the cow's teats full length with **READY-TO-USE PRO-SAN LV TEAT DIP**.
4. Wait at least 15-30 seconds.
5. Remove all teat dip by thoroughly drying the teats, teat orifices and udder with a clean, single service paper towel(s) to avoid contamination of milk.
6. Attach milking unit.

B. POST-MILKING (POST-DIPPING)

Immediately after milking, dip, spray or foam all teats full length with **READY-TO-USE PRO-SAN LV TEAT DIP**. Ensure good coverage of all sides of the teats and teat orifices. Allow to dry. Do not wipe. Do not turn cows out in freezing weather until teat dip is completely dry.

WARNING

**DO NOT DILUTE THIS PRODUCT.
DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.
ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.
STORE AT ROOM TEMPERATURE. ALWAYS STORE AWAY FROM DIRECT SUNLIGHT.
PROTECT THIS PRODUCT FROM FREEZING. HOWEVER, FREEZING WILL NOT AFFECT PRO-SAN LV PROVIDED THAT THE CONTENTS ARE THAWED COMPLETELY AND AGITATED THOROUGHLY BEFORE MIXING BASE AND ACTIVATOR TOGETHER.**

A PART OF THE COMPLETE IBA SANITATION PROGRAM

EXP. DATE:
LOT NO.:
NET CONTENTS: 55 GALLONS
IBA STOCK # 262210
323W

MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 103 GILMORE DRIVE, SUTTON, MA 01590

PRO-SAN LV BASE

sodium chlorite liquid

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:29983-1092
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORITE (UNII: G538EBV4VF) (CHLORITE ION - UNII:Z63H374SB6)	SODIUM CHLORITE	0.7 kg in 100 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29983-1092-8	210.25 kg in 1 DRUM		
2	NDC:29983-1092-9	1051.25 kg in 1 CONTAINER		
3	NDC:29983-1092-4	19.11 kg in 1 PAIL		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		04/26/2023	

Labeler - IBA (019494160)

Registrant - IBA (019494160)

Establishment

Name	Address	ID/FEI	Business Operations
Morgan Gallacher Inc. DBA in Custom Chemical Formulators Inc.		028311595	manufacture, api manufacture, pack

Revised: 12/2025

IBA