

HYDROGEN PEROXIDE- hydrogen peroxide solution
Delon Laboratories (1990) Ltd

Hydrogen Peroxide 3%

Active ingredient

Hydrogen peroxide 3%

Purpose

first aid antiseptic / oral debriding agent

Uses

- first aid to help prevent the risk of infection in minor cuts, scrapes and burns
- aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings

For external use only

Do not use

- in the eyes or apply over large areas of the body
- longer than 1 week unless directed by a doctor or dentist

Ask a doctor before use if you have

- deep or puncture wounds, animal bites, or serious burns

Stop use and ask a doctor if

- the condition persists or gets worse
- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens
- swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

first aid antiseptic:

- clean the affected area
- apply a small amount of this product on the area 1 to 3 times daily
- may be covered with a sterile bandage
- if bandaged, let dry first

oral debriding agent (oral rinse):

- adults and children 2 years of age and older:
 - mix with an equal amount of warm water
 - swish around in the mouth over the affected area for at least 1 minute and then spit out
 - use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor
-
- children under 12 years of age should be supervised in the use of the product
 - children under 2 years of age: consult a dentist or doctor

Other information

- keep tightly closed in a cool dark place
- do not shake bottle
- hold away from face when opening

Inactive ingredient

purified water

Delon Hydrogen Peroxide 3%

<p>DELON </p> <p>NDC 61734-010-01</p> <p>Hydrogen Peroxide 3% USP</p> <p>FIRST AID ANTISEPTIC ORAL DEBRIDING AGENT</p> <p>KEEP OUT OF REACH OF CHILDREN</p> <p>FOR EXTERNAL USE ONLY</p> <p>16 FL OZ</p>	<p>Tamper evident: Do not use if safety seal under cap is broken or missing.</p> <table border="1"><tr><td>Drug Facts</td><td></td></tr><tr><td>Active ingredient</td><td>Purpose</td></tr><tr><td>Hydrogen peroxide 3%</td><td>First aid antiseptic/oral debriding agent</td></tr><tr><td colspan="2">Uses</td></tr><tr><td colspan="2">■ first aid to help prevent the risk of infection in minor cuts, scrapes and burns</td></tr><tr><td colspan="2">■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth</td></tr><tr><td colspan="2">Warnings</td></tr><tr><td colspan="2">For external use only</td></tr><tr><td colspan="2">Do not use</td></tr><tr><td colspan="2">■ in the eyes or apply over large areas of the body</td></tr><tr><td colspan="2">■ longer than 1 week unless directed by a doctor or dentist</td></tr><tr><td colspan="2">Ask a doctor before use if you have</td></tr><tr><td colspan="2">■ deep or puncture wounds, animal bites, or serious burns</td></tr><tr><td colspan="2">Stop use and ask a doctor if</td></tr><tr><td colspan="2">■ the condition persists or gets worse</td></tr><tr><td colspan="2">■ sore mouth symptoms do not improve in 7 days</td></tr><tr><td colspan="2">■ irritation, pain, or redness persists or worsens</td></tr><tr><td colspan="2">■ swelling, rash, or fever develops</td></tr><tr><td colspan="2">Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</td></tr><tr><td colspan="2">Directions</td></tr><tr><td colspan="2">first aid antiseptic:</td></tr><tr><td colspan="2">■ clean the affected area</td></tr><tr><td colspan="2">■ apply a small amount of this product on the area 1 to 3 times daily</td></tr><tr><td colspan="2">■ may be covered with a sterile bandage</td></tr><tr><td colspan="2">■ if bandaged, let dry first</td></tr><tr><td colspan="2">oral debriding agent (oral rinse):</td></tr><tr><td colspan="2">■ adults and children 2 years of age and older:</td></tr><tr><td colspan="2">■ mix with an equal amount of warm water</td></tr><tr><td colspan="2">■ swish around in the mouth over the affected area for at least 1 minute and then spit out</td></tr><tr><td colspan="2">■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor</td></tr><tr><td colspan="2">■ children under 12 years of age should be supervised in the use of the product</td></tr><tr><td colspan="2">■ children under 2 years of age: consult a dentist or doctor</td></tr><tr><td colspan="2">Other information</td></tr><tr><td colspan="2">■ keep tightly closed in a cool dark place</td></tr><tr><td colspan="2">■ do not shake bottle</td></tr><tr><td colspan="2">■ hold away from face when opening</td></tr><tr><td colspan="2">Inactive ingredient purified water</td></tr></table> <p>SPACE FOR LOT # AND EXPIRY DO NOT PRINT</p>	Drug Facts		Active ingredient	Purpose	Hydrogen peroxide 3%	First aid antiseptic/oral debriding agent	Uses		■ first aid to help prevent the risk of infection in minor cuts, scrapes and burns		■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth		Warnings		For external use only		Do not use		■ in the eyes or apply over large areas of the body		■ longer than 1 week unless directed by a doctor or dentist		Ask a doctor before use if you have		■ deep or puncture wounds, animal bites, or serious burns		Stop use and ask a doctor if		■ the condition persists or gets worse		■ sore mouth symptoms do not improve in 7 days		■ irritation, pain, or redness persists or worsens		■ swelling, rash, or fever develops		Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.		Directions		first aid antiseptic:		■ clean the affected area		■ apply a small amount of this product on the area 1 to 3 times daily		■ may be covered with a sterile bandage		■ if bandaged, let dry first		oral debriding agent (oral rinse):		■ adults and children 2 years of age and older:		■ mix with an equal amount of warm water		■ swish around in the mouth over the affected area for at least 1 minute and then spit out		■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor		■ children under 12 years of age should be supervised in the use of the product		■ children under 2 years of age: consult a dentist or doctor		Other information		■ keep tightly closed in a cool dark place		■ do not shake bottle		■ hold away from face when opening		Inactive ingredient purified water		<p>NOT TESTED ON ANIMALS</p>    <p>1630-3 Made in Canada by: DELON LABORATORIES (1990) INC. Pointe-Claire, Quebec, Canada, H9R 1E2 www.delon.com</p>  <p>0 59338 01003 8</p>
Drug Facts																																																																												
Active ingredient	Purpose																																																																											
Hydrogen peroxide 3%	First aid antiseptic/oral debriding agent																																																																											
Uses																																																																												
■ first aid to help prevent the risk of infection in minor cuts, scrapes and burns																																																																												
■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth																																																																												
Warnings																																																																												
For external use only																																																																												
Do not use																																																																												
■ in the eyes or apply over large areas of the body																																																																												
■ longer than 1 week unless directed by a doctor or dentist																																																																												
Ask a doctor before use if you have																																																																												
■ deep or puncture wounds, animal bites, or serious burns																																																																												
Stop use and ask a doctor if																																																																												
■ the condition persists or gets worse																																																																												
■ sore mouth symptoms do not improve in 7 days																																																																												
■ irritation, pain, or redness persists or worsens																																																																												
■ swelling, rash, or fever develops																																																																												
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.																																																																												
Directions																																																																												
first aid antiseptic:																																																																												
■ clean the affected area																																																																												
■ apply a small amount of this product on the area 1 to 3 times daily																																																																												
■ may be covered with a sterile bandage																																																																												
■ if bandaged, let dry first																																																																												
oral debriding agent (oral rinse):																																																																												
■ adults and children 2 years of age and older:																																																																												
■ mix with an equal amount of warm water																																																																												
■ swish around in the mouth over the affected area for at least 1 minute and then spit out																																																																												
■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor																																																																												
■ children under 12 years of age should be supervised in the use of the product																																																																												
■ children under 2 years of age: consult a dentist or doctor																																																																												
Other information																																																																												
■ keep tightly closed in a cool dark place																																																																												
■ do not shake bottle																																																																												
■ hold away from face when opening																																																																												
Inactive ingredient purified water																																																																												

HYDROGEN PEROXIDE

hydrogen peroxide solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:61734-010
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROGEN PEROXIDE (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)	HYDROGEN PEROXIDE	3 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	blue	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61734-010-01	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/08/2014	
2	NDC:61734-010-02	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/08/2014	12/07/2014
3	NDC:61734-010-03	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/08/2014	12/04/2014
4	NDC:61734-010-04	946 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/08/2014	12/04/2014
5	NDC:61734-010-05	1000 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/08/2014	12/04/2014
6	NDC:61734-010-06	100 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product	10/08/2014	12/04/2014
7	NDC:61734-010-07	175 mL in 1 BOTTLE, PUMP; Type 0: Not a Combination Product	10/08/2014	10/31/2019

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	10/08/2014	

Labeler - Delon Laboratories (1990) Ltd (248364184)

Establishment

Name	Address	ID/FEI	Business Operations
Laboratoires Delon		208896216	manufacture(61734-010) , pack(61734-010) , label(61734-010)

Revised: 10/2025

Delon Laboratories (1990) Ltd