

**TOPCARE PAIN RELIEF PATCH- lidocaine, menthol patch**  
**Topco Associates LLC**

-----  
**TopCare Pain Relief Patch Lidocaine plus Menthol**

**Active ingredients**

Lidocaine 4%

Menthol 1%

**Purposes**

Topical anesthetic

Topical analgesic

**Use**

for the temporary relief of pain

**Warnings**

**For external use only**

**Do not use**

- more than 1 patch on your body at a time
- On cut, irritated, or swollen skin
- on puncture wounds
- For more than one week without consulting a doctor
- If you are allergic to any active or inactive ingredients
- if pouch is damaged or opened

**When using this product**

- use only as directed
- Read and follow all directions and warnings on this carton
- Do not allow contact with the eyes
- Do not use at the same time as other topical analgesics
- Do not bandage tightly or apply local heat (such as heating pads) to the area of use
- Do not microwave
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

**Stop use and ask a doctor if**

- condition worsens
- Redness is present
- Irritation develops
- Symptoms persist for more than 7 days or clear up and occur again within a few days
- you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children and pets.**

If swallowed, get medical help or contact a Poison Control Center 800-222-1222 right away.

**Directions****adults and children 12 years of age and over:**

- clean and dry affected area
- carefully remove backing from patch starting at a corner.
- Apply sticky side of patch to affected area.
- Use one patch for up to 12 hours
- Dispatch patch after a single use.
- **children under 12 years of age:** consult a physician

**Inactive ingredients**

aluminum glycinat, glycerin, kaolin, methylparaben, polyacrylic acid, polysorbate 80, propylene glycol, propylparaben, PVP, sodium polyacrylate, tartaric acid, titanium dioxide, water.

**Package/Label Principal Display Panel**



**Drug Facts**

**Active Ingredients:** Lidocaine 4.0% (Local anesthetic), Menthol 1.0% (Topical analgesic)

**Use:** Temporarily relieves minor pain.

**Warnings:** For external use only. Do not use in more than one patch on your body at a time or on cut, irritated or swollen skin. Do not use on puncture wounds or for more than one week without consulting a doctor. Do not use on any active or inactive ingredients if it pricks or stings or causes irritation.

**When using this product:** Use only on directed site and follow all directions and warnings on this section. Do not use other ointments with the area. Do not use of the area that is other topical medication. Do not breathe deeply or apply heat (such as heating pad) to the area of use. Do not reinsert or reuse of used patch. It is normal that always keep product away from children and pets. Used patches will contain the drug product that can produce serious adverse effects if a child or pet chews or ingests the patch.

**Stop use and consult a doctor if:** Condition worsens or becomes present in Irritation develops. If symptoms persist for more than 7 days or clear up and clear again within a few days as you experience signs of skin injury, such as pain, swelling or blistering where the product was applied.

**If pregnant or breastfeeding, seek health professional before use.**

**Keep out of reach of children and pets. If swallowed, get medical help or contact a Poison Control Center 800-235-1223 right away.**

**Directions:** Adults and children 12 years of age and over: Clean and dry affected area. Carefully remove backing from patch starting at a corner. Apply sticky side of patch to affected area. Use one patch for up to 12 hours. Discard patch after single use. Children under 12 years of age: consult a physician.

**Other Information:** Store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.

**Inactive Ingredients:** aluminum glycidate, glycerin, isobutyl alcohol, polyacrylic acid, polyacrylate G1, propylene glycol, polyisobutylene, PVP, sodium polyacrylate, butylacrylate, butyl methacrylate, urethane.

DISTRIBUTED BY TOPCARE ASSOCIATES LLC  
 1155 BRIDGE WILLOW, EL MONTE, CA 91734  
 CONTACT NUMBER: 1-800-425-8138  
 topcare.com www.topcare.com  
 MADE IN CHINA

Scan here for more information  
 or call 1-800-425-8138

0 36800 49585 2

TopCare health

MAXIMUM STRENGTH LIDOCAINE  
**Pain Relief Patch**  
 4% Lidocaine • 1 % Menthol

NDC 76162-007-05

TopCare health

COMPARE TO KY-HOT® LIDOCAINE PATCH PLUS MENTHOL ACTIVE INGREDIENTS\*


MAXIMUM STRENGTH LIDOCAINE  
**Pain Relief Patch**  
 4% Lidocaine • 1 % Menthol

- Desensitizes Aggravated Nerves & Relieves Pain
- No-Mess, Single-Use Application
- Stay-Put Flexible Patch



BACK, NECK,  
 LEG & SHOULDER

**5 PATCHES**  
 3.53" X 5.67" (10 cm X 14 cm) EACH



MAXIMUM STRENGTH LIDOCAINE  
**Pain Relief Patch**  
 4% Lidocaine • 1 % Menthol

\*KY-HOT® is a registered trademark of Chullian, Inc. Chullian, Inc. is not affiliated with TopCare or this product.



This TopCare® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.

Child-Resistant Packaging  
 Model: 23-912  
 92-52018 01/22 VCS10046

LOT  
 EXP

## TOPCARE PAIN RELIEF PATCH

lidocaine, menthol patch

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:76162-007
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LIDOCAINE</b> (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	4 g in 100 g
<b>MENTHOL, UNSPECIFIED FORM</b> (UNII: L7T10EIP3A) (MENTHOL, UNSPECIFIED FORM - UNII:L7T10EIP3A)	MENTHOL, UNSPECIFIED FORM	1 g in 100 g

### Inactive Ingredients

Ingredient Name	Strength
<b>DIHYDROXYALUMINUM AMINOACETATE ANHYDROUS</b> (UNII: 1K713C615K)	
<b>SODIUM POLYACRYLATE (2500000 MW)</b> (UNII: 05115JN12J)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	

<b>GLYCERIN</b> (UNII: PDC6A3C00X)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>KAOLIN</b> (UNII: 24H4NWX5CO)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>POLYACRYLIC ACID (250000 MW)</b> (UNII: 9G2MAD7J6W)	
<b>TARTARIC ACID</b> (UNII: W4888I119H)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>WATER</b> (UNII: 059QF0K00R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76162-007-05	5 in 1 CARTON	01/01/2022	
1		1 g in 1 POUCH; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	01/01/2022	

**Labeler** - Topco Associates LLC (006935977)

### Establishment

Name	Address	ID/FEI	Business Operations
Foshan Aqua Gel Biotech Co., Ltd		529128763	manufacture(76162-007)

Revised: 12/2024

Topco Associates LLC