

BRIGHTER DAY BROAD SPECTRUM SPF 30 SUNSCREEN- zinc oxide lotion
SQN Lab

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

SQN - BRIGHTER DAY BROAD SPECTRUM SPF30 (82495-101)

ACTIVE INGREDIENT

ZINC OXIDE

PURPOSE

SUNSCREEN

USE

HELPS PREVENT SUNBURN

WARNINGS

FOR EXTERNAL USE ONLY.

DO NOT USE ON DAMAGED OR BROKEN SKIN.

WHEN USING THIS PRODUCT KEEP OUT OF EYES. RINSE WITH WATER TO REMOVE.

STOP USE AND ASK A DOCTOR IF RASH OCCURS.

KEEP OUT OF REACH OF CHILDREN. IF PRODUCT IS SWALLOWED, GET MEDICAL HELP OR CONTACT A POISON CONTROL CENTER RIGHT AWAY.

DIRECTIONS

- APPLY LIBERALLY 15 MINUTES BEFORE SUN EXPOSURE.
- CHILDREN UNDER 6 MONTHS OF AGE: ASK A DOCTOR.
- USE A WATER-RESISTANT SUNSCREEN IF SWIMMING OR SWEATING.
- REAPPLY AT LEAST EVERY 2 HOURS.

- SUN PROTECTION MEASURES.
SPENDING TIME IN THE SUN INCREASES YOUR RISK OF SKIN CANCER AND EARLY AGING. TO DECREASE THE RISK, REGULARLY USE A SUNSCREEN WITH A BROAD SPECTRUM SPF VALUE OF 15 OR HIGHER AND OTHER SUN PROTECTION MEASURES INCLUDING:
- LIMIT TIME IN THE SUN, ESPECIALLY FROM 10:00 A.M. - 2 P.M.
- WEAR LONG-SLEEVED SHIRTS, PANTS, HATS, AND SUNGLASSES.

INACTIVE INGREDIENTS

WATER, BUTYLENE GLYCOL, CAPRYLIC/CAPRIC TRIGLYCERIDE, DIETHYLHEXYL 2,6-NAPHTHALATE, HYDROXYETHYL ACRYLATE/SODIUM, ACRYLOYLDIMETHYL TAURATE COPOLYMER, SODIUM OLIVATE, XYLITYLGLUCOSIDE, ANHYDROXYLITOL, TRIETHOXYCAPRYLYLSILANE, XYLITOL, PENTYLENE GLYCOL, CITRIC ACID, TOCOPHEROL, CAMELLIA SINENSIS LEAF OIL, DIMETHICONE, PHENOXYETHANOL, ETHYLHEXYLGLYCERIN, IRON OXIDES.

OTHER INFORMATION

- PROTECT THE PRODUCT IN THIS CONTAINER FROM EXCESSIVE HEAT AND DIRECT SUN.

an everyday essential to protect, replenish and deliver calming actives and hydration when you need it most

What matters

- + 10% zinc oxide
- + triglycerides
- + antioxidants

**BRIGHTER DAY
BROAD SPECTRUM
SPF 30
SUNSCREEN**



HYDRATING FORMULA

Drug Facts

Active ingredient	Purpose
Zinc Oxide 10%	Sunscreen

Use
Helps Prevent Sunburn

Warnings
For External Use Only.
Do not use on damaged or broken skin.
When using this product keep out of eyes. Rinse with water to remove.
Stop use and ask a doctor if rash occurs.
Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Apply liberally 15 minutes before sun exposure.
- Reapply under 80 minutes if you are a doctor.
- Use a water-resistant sunscreen if swimming or sweating.
- Reapply at least every 2 hours.
- Use Protective Measures.

Spending time in the sun increases your risk of skin cancer and early aging. To decrease the risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:

- 1. Limit time in the sun, especially from 10:00 a.m. - 2 p.m.
- 2. Wear protective clothing, hats, and sunglasses.

Inactive ingredients
Water, Butylene Glycol, Caprylic/Capric Triglyceride, Dihydroxyethyl Hexanediol, Hydroxyethyl Acrylate, Acryloyl Dimethyl Taurate Copolymer, Sodium Olivolate, Xylitylglucoside, Anhydroxylytol, Triethoxycaprylylsilane, Xylitol, Pentylene Glycol, Citric Acid, Isophenyl Camellia Sinensis Leaf Oil, Dimethicone, Phenoxyethanol, Ethylhexylglycerin, Iron Oxides.

Other information

- Protect the product in this container from excessive heat and direct sun.





See how we made it

2 fl oz / 60 ml



SON dist. by son lab | sonlab.com
Chattsworth CA 91311, USA

Made from responsibly sourced ingredients.

BRIGHTER DAY BROAD SPECTRUM SPF 30 SUNSCREEN

zinc oxide lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82495-101
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	10 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
DIETHYLHEXYL 2,6-NAPHTHALATE (UNII: I0DQJ7YGXM)	
TOCOPHEROL (UNII: R0ZB2556P8)	
ETHYLHEXYLGLYCERIN (UNII: 147D247K3P)	
WATER (UNII: 059QF0KO0R)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROXYETHYL ACRYLATE/SODIUM ACRYLOYLDIMETHYL TAURATE COPOLYMER (45000 MPA.S AT 1%) (UNII: 86FQE96TZ4)	
SODIUM OLEATE (UNII: 399SL044HN)	
XYLITYLGLUCOSIDE (UNII: 00IEZ166FB)	
ANHYDROXYLITOL (UNII: 8XWR7NN42F)	
XYLITOL (UNII: VCQ006KQ1E)	
PENTYLENE GLYCOL (UNII: 50C1307PZG)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
TEA LEAF OIL (UNII: VC855RRT77)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82495-101-11	1 in 1 BOX	12/20/2021	
1		50 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	12/20/2021	

Labeler - SQN Lab (118483828)