OMEPRAZOLE- omeprazole tablet, delayed release Little Pharma, Inc.

Curist Heartburn Relief (Omeprazole Tablets)

Drug Facts

Active ingredient (in each tablet)

Omeprazole USP 20 mg

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs <u>2 or more</u> days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert

Do not use if you are allergic to omeprazole.

Do not use if you have

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

your heartburn continues or worsens

- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20 to 25° C (68 to 77° F) and protect from moisture

Inactive ingredients

anhydrous lactose, hypromellose, hypromellose acetate succinate, iron oxide red, iron oxide yellow, lactose monohydrate, methyl cellulose, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, triethyl citrate and titanium dioxide.

The imprinting ink contains ammonium hydroxide, black iron oxide, n-butyl alcohol, propylene glycol and shellac.

Questions or Comments?

Call toll free 1-800-818-4555 weekdays.

Distributed by:

Little Pharma, Inc. New York, NY 10023

PRINCIPAL DISPLAY PANEL - 20 mg Tablet Bottle Carton

curist


Compare To Prilosec OTC ®*
Heartburn Relief

Omeprazole Delayed-release Tablets 20 mg Acid Reducer

24 HR; Treats Frequent Heartburn!

42 TABLETS


Three 14-day courses of treatment May take 1 to 4 days for full effect



ISS, 10/2021

curist





Compare To Prilosec OTC®*



curistrelief.com



Heartburn Relief

Omeprazole Delayed-release Tablets 20 mg **Acid Reducer**

24 HR; Treats Frequent Heartburn!

42 TABLETS

Three 14-day courses of treatment May take 1 to 4 days for full effect

20

Unvarnish Area: 44x33 mm

*All trademarks are the property of their respective owner. This product is not affiliated with the maker/owner of Prilosec OTC®.

Distributed by: Little Pharma, Inc. New York, NY 10023

Made In India

DNH/DRUGS/NH/138

5224457

Drug Facts

Active ingredient (in each tablet)

Purpose

- treats frequent heartburn (occurs 2 or more days a
- week)

 not intended for immediate relief of heartburn; this drug
 may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

- Stop use and ask a doctor if:

 your hearthurn continues or worsens

 you need to take this product for more than 14 days

 you need to take more than 1 course of treatment every
 4 months

 you get diarrhea

 you det diarrhea

 you det weekop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.


Drug Facts (continued)

do not take for more than 14 days or more often
 than every 4 months unless directed by a doctor
 children under 18 years of age: ask a doctor. Heartburn
 in children may sometimes be caused by a serious
 condition.

- Other information
 read the directions and warnings before use
 keep the carbon. It contains important information.
 store at 20 to 25° C (68 to 77° F) and protect from

Inactive ingredients anhydrous lactose

Tips for Managing Heartburn
■ Do not lie flat or bend over after eating ■ Do not wear tight-fitting clothing around the stomach ■ Do not eat before bedtime ■ Raise the head of your bed ■ Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocotate, caffeine, alsohol and certain fruits and vegetables ■ Eat slowly and avoid big meals ■ If overweight, lose weight ■ Quit smoking



OMEPRAZOLE

omeprazole tablet, delayed release

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72559-014	
Route of Administration	ORAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
OMEPRAZOLE (UNII: KG60484QX9) (OMEPRAZOLE - UNII:KG60484QX9)	OMEPRAZ OLE	20 mg	

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
HYPROMELLOSE ACETATE SUCCINATE 12070923 (3 MM2/S) (UNII: 36BGF0E889)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MONOETHANOLAMINE (UNII: 5KV86114PT)	
METHYLCELLULOSE (1500 MPA.S) (UNII: PONTE48364)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
SODIUM STEARATE (UNII: QU7E2XA9TG)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	

TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
AMMONIA (UNII: 5138Q19F1X)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
BUTYL ALCOHOL (UNII: 8PJ61P6TS3)	
SHELLAC (UNII: 46N107B710)	

Product Characteristics			
Color	brown (brownish pink)	Score	no score
Shape	OVAL (bioconvex)	Size	12mm
Flavor		Imprint Code	20
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72559-014- 06	3 in 1 CARTON	07/15/2020	
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:72559-014- 07	1 in 1 CARTON	07/15/2020	
2		14 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:72559-014- 36	4 in 1 PACKAGE, COMBINATION	06/18/2024	
3		3 in 1 CARTON		
3		14 in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA207891	07/15/2020	

Labeler - Little Pharma, Inc. (074328189)

Establishment				
Na me	Address	ID/FEI	Business Operations	
Ohm Laboratories		184769029	manufacture(72559-014)	

Revised: 6/2024 Little Pharma, Inc.