


**DEXAMETHASONE SODIUM PHOSPHATE - dexamethasone sodium phosphate solution**  
**Dispensing Solutions, Inc.**

-----  
**Dexamethasone Sodium Phosphate Ophthalmic Solution, USP**  
**Equivalent to dexamethasone phosphate 0.1%**  
**STERILE**

**DESCRIPTION:**

Dexamethasone sodium phosphate ophthalmic solution is a clear, colorless to pale yellow topical steroid solution for ophthalmic or otic administration. Glucocorticoids are adrenocortical steroids, both naturally occurring and synthetic. Dexamethasone is a synthetic analog of naturally occurring glucocorticoids (hydrocortisone and cortisone). Dexamethasone sodium phosphate is a water soluble, inorganic ester of dexamethasone. It is approximately three thousand times more soluble in water at 25°C than hydrocortisone.

Dexamethasone sodium phosphate is 9-fluoro-11 $\beta$ , 17-dihydroxy-16 $\alpha$ -methyl-21-(phosphonoxy)pregna-1,4-diene-3, 20-dione disodium salt and has the following structural formula:



**Each mL contains: Active:** dexamethasone sodium phosphate equivalent to 1mg (0.1%) dexamethasone phosphate. **Preservative:** benzalkonium chloride 0.01%. **Inactives:** monobasic sodium phosphate, sodium chloride, dibasic sodium phosphate, edetate disodium, monobasic sodium phosphate and/or dibasic sodium phosphate to adjust pH (6.6 to 7.8) and water for injection.

**CLINICAL PHARMACOLOGY:**

Dexamethasone sodium phosphate suppresses the inflammatory response to a variety of agents and it probably delays or slows healing. No generally accepted explanation of these steroid properties has been advanced.

**INDICATIONS AND USAGE:**

For the treatment of the following conditions:

**Ophthalmic:** Steroid responsive inflammatory conditions of the palpebral and bulbar conjunctiva, cornea, and anterior segment of the globe, such as allergic conjunctivitis, acne rosacea, superficial punctuate keratitis, herpes zoster keratitis, iritis, cyclitis, selected infective conjunctivitis when the inherent hazard of steroid use is accepted to obtain an advisable diminution in edema and inflammation;

corneal injury from chemical or thermal burns, or penetration of foreign bodies.

**Otic:** Steroid responsive inflammatory conditions of the external auditory meatus, such as allergic otitis externa, selected purulent and nonpurulent infective otitis externa when the hazard of steroid use is accepted to obtain an advisable diminution in edema and inflammation.

### **CONTRAINDICATIONS:**

Epithelial herpes simplex keratitis (dendritic keratitis). Acute infectious stages of vaccinia, varicella, and many other viral diseases of the cornea and conjunctiva.

Mycobacterial infection of the eye.

Fungal diseases of ocular or auricular structures.

Hypersensitivity to any component of this product.

Perforation of a drum membrane.

### **WARNINGS:**

Prolonged use may result in ocular hypertension and/or glaucoma, with damage to the optic nerve, defects in visual acuity and fields of vision, and posterior subcapsular cataract formation. Prolonged use may suppress the host response and thus increase the hazard of secondary ocular infections. In those diseases causing thinning of the cornea or sclera, perforations have been known to occur with the use of topical corticosteroids. In acute purulent conditions of the eye or ear, corticosteroids may mask infection or enhance existing infection. If these products are used for 10 days or longer, intraocular pressure should be routinely monitored even though it may be difficult in children and uncooperative patients.

Employment of corticosteroid medication in the treatment of herpes simplex other than epithelial herpes simplex keratitis, in which it is contraindicated, requires great caution; periodic slit-lamp microscopy is essential.

### **PRECAUTIONS:**

#### **General:**

The possibility of persistent fungal infections of the cornea should be considered after prolonged corticosteroid dosing. There have been reports of bacterial keratitis associated with the use of multiple dose containers of topical ophthalmic products. These containers had been inadvertently contaminated by patients who, in most cases, had a concurrent corneal disease or a disruption of the ocular epithelial surface. (See PRECAUTIONS, Information for Patients.)

#### **Information for Patients :**

Patients should be instructed to avoid allowing the tip of the dispensing container to contact the eye or surrounding structures. Patients should also be instructed that ocular solutions, if handled improperly, can become contaminated by common bacteria known to cause ocular infections. Serious damage to the eye and subsequent loss of vision may result from using contaminated solutions. (See PRECAUTIONS, General.)

Patients should also be advised that if they develop an intercurrent ocular condition (e.g., trauma, ocular surgery or infection), they should immediately seek their physician's advice concerning the continued use of the present multidose container. One of the preservatives in dexamethasone sodium phosphate ophthalmic solution, benzalkonium chloride, may be absorbed by soft contact lenses. Patients wearing soft contact lenses should be instructed to wait at least 15 minutes after instilling dexamethasone sodium phosphate ophthalmic solution before they insert their lenses.

**Carcinogenesis, Mutagenesis, Impairment of Fertility:**

Long-term animal studies have not been performed to evaluate the carcinogenic potential or the effect on fertility of dexamethasone sodium phosphate ophthalmic solution.

**Pregnancy:**

Pregnancy Category C: Dexamethasone has been shown to be teratogenic in mice and rabbits following topical ophthalmic application in multiples of the therapeutic dose.

In the mouse, corticosteroids produce fetal resorptions and a specific abnormality, cleft palate. In the rabbit, corticosteroids have produced fetal resorptions and multiple abnormalities involving the head, ears, limbs, palate, etc.

There are no adequate or well-controlled studies in pregnant women. Dexamethasone sodium phosphate ophthalmic solution should be used during pregnancy only if the potential benefit to the mother justifies the potential risk to the embryo or fetus. Infants born of mothers who have received substantial doses of corticosteroids during pregnancy should be observed carefully for signs of hypoadrenalism.

**Nursing Mothers :**

Topically applied steroids are absorbed systemically. Therefore, because of the potential for serious adverse reactions in nursing infants from dexamethasone sodium phosphate, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

**Pediatric Use:**

Safety and effectiveness in pediatric patients have not been established.

**Geriatric Use:**

No overall differences in safety and effectiveness have been observed between elderly and younger patients.

**ADVERSE REACTIONS:**

Glaucoma with optic nerve damage, visual acuity and field defects, posterior subcapsular cataract formation, secondary ocular infection from pathogens including herpes simplex, perforation of the globe.

Rarely, filtering blebs have been reported when topical steroids have been used following cataract surgery.

Rarely, stinging or burning may occur.

**DOSAGE AND ADMINISTRATION:**

The duration of treatment will vary with the type of lesion and may extend from a few days to several weeks, according to therapeutic response. Relapses, more common in chronic active lesions than in self-limited conditions, usually respond to treatment.

**Eye:** Instill one or two drops of solution into the conjunctival sac every hour during the day and every two hours during the night as initial therapy. When a favorable response is observed, reduce dosage to one drop every four hours. Later, further reduction in dosage to one drop three or four times daily may suffice to control symptoms.

**Ear:** Clean the aural canal thoroughly and sponge dry. Instill the solution directly into the aural canal. A suggested initial dosage is three or four drops two or three times a day. When a favorable response is obtained, reduce dosage gradually and eventually discontinue. If preferred, the aural canal may be packed with a gauze wick saturated with solution. Keep the wick moist with the preparation and remove from the ear after 12 to 24 hours. Treatment may be repeated as often as necessary at the discretion of the physician.

**HOW SUPPLIED:**

Dexamethasone sodium phosphate ophthalmic solution USP, 0.1% is available in a 5 mL DROP-TAINER®\* Dispenser Bottle.

NDC 61314-294-05

Store at controlled room temperature 15° - 30°C (59° - 86°F).

**Rx Only**

Revised: March 2006

\*DROP-TAINER is a registered trademark of Alcon Research, Ltd.

**9004571-0409**

Dist. By:

**FALCON Pharmaceuticals, Ltd.**

Fort Worth, Texas 76134 USA

Mfd. By:

**Alcon Laboratories, Inc.**

6201 South Freeway

Fort Worth, Texas 76134 USA

1-800-757-9195

MedInfo@AlconLabs.com

**PRINCIPAL DISPLAY PANEL**

**BULK SOURCE DATA**

DIST. BY: FALCON PHARMACEUTICALS, LTD.  
FORT WORTH, TEXAS 76134

**PRODUCT ID:**

WHITE / BLACK / RED / PINK  
BOX PRINTED NDC 61314-294-05

**BULK SOURCE NDC:** 61314-0294-05

MFR. LOT: XXXXXX

PEDIGREE #: 17515859

DISPENSE IN THIS  
TIGHT/LIGHT RESISTANT CONTAINER



INSTILL \_\_\_ DROP(S) INTO  
AFFECTED EYE(S) EVERY  
\_\_\_ HOURS \_\_\_ TIMES A DAY  
FOR \_\_\_ DAY(S).



**DEXAMETHASONE\* 0.1%**  
**5 mL OPTHALMIC SOLUTION**  
**NDC 55045-1755-05**  
**PRODUCT # 9D0477**

Rev. Date: 12/09

EACH mL CONTAINS:  
\*DEXAMETHASONE SODIUM PHOSPHATE USP  
EQUIVALENT TO 1 mg (0.1%)  
DEXAMETHASONE PHOSPHATE

LOT# SAMPLE EXP: 00-00 Rx # 21150930

**RX ONLY**

WARNING: KEEP OUT OF  
CHILDREN'S REACH  
STORE AT 59° - 86° F

9D0477 NDC 55045-1755-05  
DEXAMETHASONE\* 0.1%  
5 mL OPTHALMIC SOLUTION  
LOT # SAMPLE EXP: 00-00  
MN 61314-0294-05 RX#21150930

9D0477 NDC 55045-1755-05  
DEXAMETHASONE\* 0.1%  
5 mL OPTHALMIC SOLUTION  
LOT # SAMPLE EXP: 00-00  
MN 61314-0294-05 RX#21150930

9D0477 NDC 55045-1755-05  
DEXAMETHASONE\* 0.1%  
5 mL OPTHALMIC SOLUTION  
LOT # SAMPLE EXP: 00-00  
MN 61314-0294-05 RX#21150930



Packaged Exclusively By:  
**DISPENSING SOLUTIONS** inc  
Santa Ana, CA 92704

NDC 55045-1755-05 Rx Only  
Dexamethasone Sodium Phosphate  
Ophthalmic Solution, USP  
equivalent to dexamethasone phosphate  
0.1%  
5 mL STERILE

**DEXAMETHASONE SODIUM PHOSPHATE**

dexamethasone sodium phosphate solution

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:55045-1755(NDC:61314-294)
<b>Route of Administration</b>	OPHTHALMIC		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
DEXAMETHASONE SODIUM PHOSPHATE (UNII: A19376Y64P) (DEXAMETHASONE - UNII:7S5I7G3JQL)	DEXAMETHASONE	1 mg in 1 mL

**Inactive Ingredients**

Ingredient Name	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
SODIUM PHOSPHATE, MONOBASIC (UNII: 3980JIH2SW)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM PHOSPHATE, DIBASIC (UNII: GR686LBA74)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
WATER (UNII: 059QF0KO0R)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55045-1755-5	5 mL in 1 BOTTLE, PLASTIC		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA088771	07/01/2004	

**Labeler** - Dispensing Solutions, Inc. (066070785)**Establishment**

Name	Address	ID/FEI	Business Operations
Dispensing Solutions, Inc.		066070785	relabel, repack

Revised: 10/2011

Dispensing Solutions, Inc.