

BRIVARACETAM - brivaracetam tablet, film coated
Zydus Pharmaceuticals USA Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use BRIVARACETAM TABLETS safely and effectively. See full prescribing information for BRIVARACETAM TABLETS.

BRIVARACETAM tablets, for oral use, CV

Initial U.S. Approval: 2016

INDICATIONS AND USAGE

Brivaracetam tablets are indicated for the treatment of partial-onset seizures in patients 4 years of age and older. (1)

DOSAGE AND ADMINISTRATION

- **Adults (16 Years and Older):** The recommended starting dosage for monotherapy or adjunctive therapy is 50 mg twice daily (100 mg per day). Based on individual patient tolerability and therapeutic response, the dosage may be adjusted down to 25 mg twice daily (50 mg per day) or up to 100 mg twice daily (200 mg per day). (2.1)
- **Pediatric Patients (4 Years to less than 16 Years):** The recommended dosage is based on body weight and is administered orally twice daily (2.1)
- **Hepatic Impairment:** Dose adjustment is recommended for all stages of hepatic impairment. (2.5)

DOSAGE FORMS AND STRENGTHS

Tablets: 10 mg, 25 mg, 50 mg, 75 mg and 100 mg (3)

CONTRAINDICATIONS

Hypersensitivity to brivaracetam or any of the inactive ingredients in brivaracetam tablets. (4)

WARNINGS AND PRECAUTIONS

- **Suicidal Behavior and Ideation:** Monitor patients for suicidal behavior and ideation. (5.1)
- **Neurological Adverse Reactions:** Monitor for somnolence and fatigue and advise patients not to drive or operate machinery until they have gained sufficient experience on brivaracetam. (5.2)
- **Psychiatric Adverse Reactions:** Behavioral reactions including psychotic symptoms, irritability, depression, aggressive behavior and anxiety; monitor patients for symptoms. (5.3)
- **Hypersensitivity: Bronchospasm and Angioedema:** Advise patients to seek immediate medical care. Discontinue and do not restart brivaracetam if hypersensitivity occurs. (5.4)
- **Withdrawal of Antiepileptic Drugs:** Brivaracetam should be gradually withdrawn. (5.5)

ADVERSE REACTIONS

Adults: Most common adverse reactions (at least 5% for brivaracetam and at least 2% more frequently than placebo) are somnolence/sedation, dizziness, fatigue and nausea/vomiting. (6.1)

Pediatric Patients: Most common adverse reactions are similar to those seen in adult patients. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Zydus Pharmaceuticals (USA) Inc. at 1-877-993-8779 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- **Rifampin:** Because of decreased concentrations, increasing brivaracetam dosage in patients on concomitant rifampin is recommended. (2.6, 7.1)
- **Carbamazepine:** Because of increased exposure to carbamazepine metabolite, if tolerability issues arise, consider reducing carbamazepine dosage in patients on concomitant brivaracetam. (7.2)
- **Phenytoin:** Because phenytoin concentrations can increase, phenytoin levels should be monitored in patients on concomitant brivaracetam. (7.3)
- **Levetiracetam:** Brivaracetam had no added therapeutic benefit when co-administered with levetiracetam. (7.4)

USE IN SPECIFIC POPULATIONS

Pregnancy: Based on animal data, may cause fetal harm. (8.1)

Additional pediatric use information is approved for UCB, Inc's Briviact (brivaracetam) tablets. However, due to UCB, Inc's marketing exclusivity rights, this drug product is not labeled with that pediatric information.

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosage Information
- 2.2 Administration Instructions for Brivaracetam Tablets
- 2.4 Discontinuation of Brivaracetam
- 2.5 Patients with Hepatic Impairment
- 2.6 Co-administration with Rifampin

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Suicidal Behavior and Ideation
- 5.2 Neurological Adverse Reactions
- 5.3 Psychiatric Adverse Reactions
- 5.4 Hypersensitivity: Bronchospasm and Angioedema
- 5.5 Withdrawal of Antiepileptic Drugs
- 5.6 Risk of Allergic Reactions Due to Tartrazine

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

7 DRUG INTERACTIONS

- 7.1 Rifampin
- 7.2 Carbamazepine
- 7.3 Phenytoin
- 7.4 Levetiracetam

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

9 DRUG ABUSE AND DEPENDENCE

- 9.1 Controlled Substance
- 9.2 Abuse
- 9.3 Dependence

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

16.2 Storage and Handling

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Brivaracetam tablets are indicated for the treatment of partial-onset seizures in patients 4 years of age and older.

Additional pediatric use information is approved for UCB, Inc's Briviact (brivaracetam) tablets. However, due to UCB, Inc's marketing exclusivity rights, this drug product is not labeled with that pediatric information.

2 DOSAGE AND ADMINISTRATION

2.1 Dosage Information

Monotherapy or Adjunctive Therapy

The recommended dosage for patients 4 years of age and older is included in Table 1. In pediatric patients weighing less than 50 kg, the recommended dosing regimen is dependent upon body weight. When initiating treatment, gradual dose escalation is not required. Dosage should be adjusted based on clinical response and tolerability.

Table 1 Recommended Dosage for Patients 4 Years of Age and Older

Age and Body Weight	Initial Dosage	Minimum and Maximum Maintenance Dosage
Adults (16 years and older)	50 mg twice daily (100 mg per day)	25 mg to 100 mg twice daily (50 mg to 200 mg per day)
Pediatric patients weighing 50 kg or more	25 mg to 50 mg twice daily (50 mg to 100 mg per day)	25 mg to 100 mg twice daily (50 mg to 200 mg per day)
Pediatric patients weighing 20 kg to less than 50 kg	0.5 mg/kg to 1 mg/kg twice daily (1 mg/kg to 2 mg/kg per day)	0.5 mg/kg to 2 mg/kg twice daily (1 mg/kg to 4 mg/kg per day)
Pediatric patients weighing 11 kg to less than 20 kg	0.5 mg/kg to 1.25 mg/kg twice daily (1 mg/kg to 2.5 mg/kg per day)	0.5 mg/kg to 2.5 mg/kg twice daily (1 mg/kg to 5 mg/kg per day)

1000 mg/day

day)

1000 mg/day

Additional pediatric use information is approved for UCB, Inc's Briviact (brivaracetam) tablets. However, due to UCB, Inc's marketing exclusivity rights, this drug product is not labeled with that pediatric information.

2.2 Administration Instructions for Brivaracetam Tablets

Brivaracetam can be initiated with oral administration.

Brivaracetam tablets may be taken with or without food.

Brivaracetam tablets should be swallowed whole with liquid. Brivaracetam tablets should not be chewed or crushed.

2.4 Discontinuation of Brivaracetam

Avoid abrupt withdrawal from brivaracetam in order to minimize the risk of increased seizure frequency and status epilepticus [see *Warnings and Precautions (5.5)* and *Clinical Studies (14)*].

2.5 Patients with Hepatic Impairment

The recommended dosage for patients with hepatic impairment is included in Table 2 [see *Use in Specific Populations (8.7)* and *Clinical Pharmacology (12.3)*].

Table 2 Recommended Dosage for Patients with Hepatic Impairment

Age and Body Weight	Initial Dosage	Maximum Maintenance Dosage
Adults (16 years and older)		
Pediatric patients weighing 50 kg or more	25 mg twice daily (50 mg per day)	75 mg twice daily (150 mg per day)
Pediatric patients weighing 20 kg to less than 50 kg	0.5 mg/kg twice daily (1 mg/kg per day)	1.5 mg/kg twice daily (3 mg/kg per day)
Pediatric patients weighing 11 kg to less than 20 kg	0.5 mg/kg twice daily (1 mg/kg per day)	2 mg/kg twice daily (4 mg/kg per day)

Additional pediatric use information is approved for UCB, Inc's Briviact (brivaracetam) tablets. However, due to UCB, Inc's marketing exclusivity rights, this drug product is not labeled with that pediatric information.

2.6 Co-administration with Rifampin

Increase the brivaracetam dosage in patients on concomitant rifampin by up to 100% (i.e., double the dosage) [see *Drug Interactions (7.1)* and *Clinical Pharmacology (12.3)*].

3 DOSAGE FORMS AND STRENGTHS

- Brivaracetam tablets, 10 mg are white to off-white, round, biconvex, film-coated tablets debossed with "U" on one side and plain on the other.
- Brivaracetam tablets, 25 mg are light blue, oval, biconvex, film-coated tablets debossed with "UI" on one side and plain on the other.
- Brivaracetam tablets, 50 mg are white to off-white, oval, biconvex, film-coated tablets debossed with "Ull" on one side and plain on the other.
- Brivaracetam tablets, 75 mg are purple, oval, biconvex, film-coated tablets debossed with "1587" on one side and plain on the other.
- Brivaracetam tablets, 100 mg are light green, oval, biconvex, film-coated tablets debossed with "1588" on one side and plain on the other.

4 CONTRAINDICATIONS

Hypersensitivity to brivaracetam or any of the inactive ingredients in brivaracetam tablets (bronchospasm and angioedema have occurred) [see *Warnings and Precautions* (5.4)].

5 WARNINGS AND PRECAUTIONS

5.1 Suicidal Behavior and Ideation

Antiepileptic drugs (AEDs), including brivaracetam, increase the risk of suicidal thoughts or behavior in patients taking these drugs for any indication. Patients treated with any AED for any indication should be monitored for the emergence or worsening of depression, suicidal thoughts or behavior and/or any unusual changes in mood or behavior.

Pooled analyses of 199 placebo-controlled clinical trials (mono- and adjunctive therapy) of 11 different AEDs showed that patients randomized to one of the AEDs had approximately twice the risk (adjusted Relative Risk 1.8, 95% CI:1.2, 2.7) of suicidal thinking or behavior compared to patients randomized to placebo. In these trials, which had a median treatment duration of 12 weeks, the estimated incidence rate of suicidal behavior or ideation among 27,863 AED-treated patients was 0.43%, compared to 0.24% among 16,029 placebo-treated patients, representing an increase of approximately one case of suicidal thinking or behavior for every 530 patients treated. There were four suicides in drug-treated patients in the trials and none in placebo-treated patients, but the number is too small to allow any conclusion about drug effect on suicide.

The increased risk of suicidal thoughts or behavior with AEDs was observed as early as one week after starting drug treatment with AEDs and persisted for the duration of treatment assessed. Because most trials included in the analysis did not extend beyond 24 weeks, the risk of suicidal thoughts or behavior beyond 24 weeks could not be assessed.

The risk of suicidal thoughts or behavior was generally consistent among drugs in the data analyzed. The finding of increased risk with AEDs of varying mechanisms of action and across a range of indications suggests that the risk applies to all AEDs used for any

indication. The risk did not vary substantially by age (5 years to 100 years) in the clinical trials analyzed. Table 3 shows absolute and relative risk by indication for all evaluated AEDs.

Table 3 Risk of Suicidal Thoughts or Behaviors by Indication for Antiepileptic Drugs in the Pooled Analysis

Indication	Placebo Patients with Events Per 1,000 Patients	Drug Patients with Events Per 1,000 Patients	Relative Risk: Incidence of Events in Drug Patients/Incidence in Placebo Patients	Risk Difference: Additional Drug Patients with Events Per 1,000 Patients
Epilepsy	1	3.4	3.5	2.4
Psychiatric	5.7	8.5	1.5	2.9
Other	1	1.8	1.9	0.9
Total	2.4	4.3	1.8	1.9

The relative risk for suicidal thoughts or behavior was higher in clinical trials in patients with epilepsy than in clinical trials in patients with psychiatric or other conditions, but the absolute risk differences were similar for the epilepsy and psychiatric indications.

Anyone considering prescribing brivaracetam or any other AED must balance the risk of suicidal thoughts or behaviors with the risk of untreated illness. Epilepsy and many other illnesses for which AEDs are prescribed are themselves associated with morbidity and mortality and an increased risk of suicidal thoughts and behavior. Should suicidal thoughts and behavior emerge during treatment, consider whether the emergence of these symptoms in any given patient may be related to the illness being treated.

5.2 Neurological Adverse Reactions

Brivaracetam causes somnolence, fatigue, dizziness and disturbance in coordination. Patients should be monitored for these signs and symptoms and advised not to drive or operate machinery until they have gained sufficient experience on brivaracetam to gauge whether it adversely affects their ability to drive or operate machinery.

Somnolence and Fatigue

Brivaracetam causes dose-dependent increases in somnolence and fatigue-related adverse reactions (fatigue, asthenia, malaise, hypersomnia, sedation and lethargy) [see *Adverse Reactions (6.1)*]. In the Phase 3 controlled adjunctive epilepsy trials, these events were reported in 25% of patients randomized to receive brivaracetam at least 50 mg/day (20% at 50 mg/day, 26% at 100 mg/day and 27% at 200 mg/day) compared to 14% of patients who received placebo. The risk is greatest early in treatment but can occur at any time.

Dizziness and Disturbance in Gait and Coordination

Brivaracetam causes adverse reactions related to dizziness and disturbance in gait and coordination (dizziness, vertigo, balance disorder, ataxia, nystagmus, gait disturbance

and abnormal coordination) [see *Adverse Reactions (6.1)*]. In the Phase 3 controlled adjunctive epilepsy trials, these events were reported in 16% of patients randomized to receive brivaracetam at least 50 mg/day compared to 10% of patients who received placebo. The risk is greatest early in treatment but can occur at any time.

5.3 Psychiatric Adverse Reactions

Brivaracetam causes psychiatric adverse reactions. In the Phase 3 controlled adjunctive epilepsy trials, psychiatric adverse reactions were reported in approximately 13% of patients who received brivaracetam (at least 50 mg/day) compared to 8% of patients who received placebo. Psychiatric events included both non-psychotic symptoms (irritability, anxiety, nervousness, aggression, belligerence, anger, agitation, restlessness, depression, depressed mood, tearfulness, apathy, altered mood, mood swings, affect lability, psychomotor hyperactivity, abnormal behavior and adjustment disorder) and psychotic symptoms (psychotic disorder along with hallucination, paranoia, acute psychosis and psychotic behavior). A total of 1.7% of adult patients treated with brivaracetam discontinued treatment because of psychiatric reactions compared to 1.3% of patients who received placebo.

Psychiatric adverse reactions were also observed in open-label pediatric trials and were generally similar to those observed in adults [see *Adverse Reactions (6.1)* and *Use in Specific Populations (8.4)*].

5.4 Hypersensitivity: Bronchospasm and Angioedema

Brivaracetam can cause hypersensitivity reactions. Bronchospasm and angioedema have been reported in patients taking brivaracetam. If a patient develops hypersensitivity reactions after treatment with brivaracetam, the drug should be discontinued. Brivaracetam tablets are contraindicated in patients with a prior hypersensitivity reaction to brivaracetam or any of the inactive ingredients [see *Contraindications (4)*].

5.5 Withdrawal of Antiepileptic Drugs

As with most antiepileptic drugs, brivaracetam should generally be withdrawn gradually because of the risk of increased seizure frequency and status epilepticus [see *Dosage and Administration (2.4)* and *Clinical Studies (14)*]. But if withdrawal is needed because of a serious adverse event, rapid discontinuation can be considered.

5.6 Risk of Allergic Reactions Due to Tartrazine

Brivaracetam tablets, 100 mg contain FD&C Yellow No. 5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. Although the overall incidence of FD&C Yellow No. 5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity.

6 ADVERSE REACTIONS

The following serious adverse reactions are described elsewhere in labeling:

- Suicidal Behavior and Ideation [see *Warnings and Precautions (5.1)*]
- Neurological Adverse Reactions [see *Warnings and Precautions (5.2)*]
- Psychiatric Adverse Reactions [see *Warnings and Precautions (5.3)*]
- Hypersensitivity: Bronchospasm and Angioedema [see *Warnings and Precautions*

(5.4)]

- Withdrawal of Antiepileptic Drugs [see *Warnings and Precautions* (5.5)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In all controlled and uncontrolled trials performed in adult epilepsy patients, brivaracetam was administered as adjunctive therapy to 2,437 patients. Of these patients, 1,929 were treated for at least 6 months, 1,500 for at least 12 months, 1,056 for at least 24 months and 758 for at least 36 months. A total of 1,558 patients (1,099 patients treated with brivaracetam and 459 patients treated with placebo) constituted the safety population in the pooled analysis of Phase 3 placebo-controlled studies in patients with partial-onset seizures (Studies 1, 2 and 3) [see *Clinical Studies* (14)]. The adverse reactions presented in Table 4 are based on this safety population; the median length of treatment in these studies was 12 weeks. Of the patients in those studies, approximately 51% were male, 74% were Caucasian and the mean age was 38 years.

In the Phase 3 controlled epilepsy studies, adverse events occurred in 68% of patients treated with brivaracetam and 62% treated with placebo. The most common adverse reactions occurring at a frequency of at least 5% in patients treated with brivaracetam doses of at least 50 mg/day and greater than placebo were somnolence and sedation (16%), dizziness (12%), fatigue (9%) and nausea and vomiting symptoms (5%).

The discontinuation rates due to adverse events were 5%, 8% and 7% for patients randomized to receive brivaracetam at the recommended doses of 50 mg, 100 mg and 200 mg/day, respectively, compared to 4% in patients randomized to receive placebo.

Table 4 lists adverse reactions for brivaracetam that occurred at least 2% more frequently for brivaracetam doses of at least 50 mg/day than placebo.

Table 4 Adverse Reactions in Pooled Placebo-Controlled Adjunctive Therapy Studies in Adult Patients with Partial-Onset Seizures (Brivaracetam 50 mg/day, 100 mg/day and 200 mg/day)

Adverse Reactions	Brivaracetam (N=803) %	Placebo (N=459) %
Gastrointestinal disorders		
Nausea/vomiting symptoms	5	3
Constipation	2	0
Nervous system disorders		
Somnolence and sedation	16	8
Dizziness	12	7
Fatigue	9	4
Cerebellar coordination and balance disturbances*	3	1
Psychiatric disorders		

*Cerebellar coordination and balance disturbances includes ataxia, balance disorder, coordination abnormal and nystagmus.

There was no apparent dose-dependent increase in adverse reactions listed in Table 4 with the exception of somnolence and sedation.

Pediatric Patients

Safety of brivaracetam was evaluated in two open-label, safety and pharmacokinetic trials in pediatric patients 4 years to less than 16 years of age. Across studies of pediatric patients with partial onset seizures, 149 patients received brivaracetam oral solution or tablet, of whom 107 received brivaracetam for at least 12 months. Adverse reactions reported in clinical studies of pediatric patients were generally similar to those seen in adult patients.

Hematologic Abnormalities

Brivaracetam can cause hematologic abnormalities. In the Phase 3 controlled adjunctive epilepsy studies, a total of 1.8% of brivaracetam-treated patients and 1.1% of placebo-treated patients had at least one clinically significant decreased white blood cell count ($< 3 \times 10^9/L$) and 0.3% of brivaracetam-treated patients and 0% of placebo-treated patients had at least one clinically significant decreased neutrophil count ($< 1 \times 10^9/L$).

Comparison by Sex

There were no significant differences by sex in the incidence of adverse reactions.

Additional pediatric use information is approved for UCB, Inc's Briviact (brivaracetam) tablets. However, due to UCB, Inc's marketing exclusivity rights, this drug product is not labeled with that pediatric information.

7 DRUG INTERACTIONS

7.1 Rifampin

Co-administration with rifampin decreases brivaracetam plasma concentrations likely because of CYP2C19 induction [see *Clinical Pharmacology (12.3)*]. Prescribers should increase the brivaracetam dose by up to 100% (i.e., double the dosage) in patients while receiving concomitant treatment with rifampin [see *Dosage and Administration (2.6)*].

7.2 Carbamazepine

Co-administration with carbamazepine may increase exposure to carbamazepine-epoxide, the active metabolite of carbamazepine. Though available data did not reveal any safety concerns, if tolerability issues arise when co-administered, carbamazepine dose reduction should be considered [see *Clinical Pharmacology (12.3)*].

7.3 Phenytoin

Because brivaracetam can increase plasma concentrations of phenytoin, phenytoin levels should be monitored in patients when concomitant brivaracetam is added to or discontinued from ongoing phenytoin therapy [see *Clinical Pharmacology (12.3)*].

7.4 Levetiracetam

Brivaracetam provided no added therapeutic benefit to levetiracetam when the two drugs were co-administered [see *Clinical Studies* (14)].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to antiepileptic drugs (AEDs), such as brivaracetam, during pregnancy. Encourage patients who are taking brivaracetam during pregnancy to enroll in the North American Antiepileptic Drug (NAAED) Pregnancy Registry by calling the toll free number 1-888-233-2334 or visiting <http://www.aedpregnancyregistry.org/>.

Risk Summary

There are no adequate data on the developmental risks associated with use of brivaracetam in pregnant women. In animal studies, brivaracetam produced evidence of developmental toxicity (increased embryofetal mortality and decreased fetal body weights in rabbits; decreased growth, delayed sexual maturation and long-term neurobehavioral changes in rat offspring) at maternal plasma exposures greater than clinical exposures [see *Data*].

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively. The background risk of major birth defects and miscarriage for the indicated population is unknown.

Data

Animal Data

Oral administration of brivaracetam (0 mg/kg/day, 150 mg/kg/day, 300 mg/kg/day or 600 mg/kg/day) to pregnant rats during the period of organogenesis did not produce any significant maternal or embryofetal toxicity. The highest dose tested was associated with maternal plasma exposures (AUC) approximately 30 times exposures in humans at the maximum recommended dose (MRD) of 200 mg/day.

Oral administration of brivaracetam (0 mg/kg/day, 30 mg/kg/day, 60 mg/kg/day, 120 mg/kg/day or 240 mg/kg/day) to pregnant rabbits during the period of organogenesis resulted in embryofetal mortality and decreased fetal body weights at the highest dose tested, which was also maternally toxic. The highest no-effect dose (120 mg/kg/day) was associated with maternal plasma exposures approximately 4 times human exposures at the MRD.

When brivaracetam (0 mg/kg/day, 150 mg/kg/day, 300 mg/kg/day or 600 mg/kg/day) was orally administered to rats throughout pregnancy and lactation, decreased growth, delayed sexual maturation (female) and long-term neurobehavioral changes were observed in the offspring at the highest dose. The highest no-effect dose (300 mg/kg/day) was associated with maternal plasma exposures approximately 7 times human exposures at the MRD.

Brivaracetam was shown to readily cross the placenta in pregnant rats after a single oral (5 mg/kg) dose of ¹⁴C-brivaracetam. From 1 hour post dose, radioactivity levels in fetuses, amniotic fluid and placenta were similar to those measured in maternal blood.

8.2 Lactation

Risk Summary

No data are available regarding the presence of brivaracetam in human milk, the effects on the breastfed infant or the effects of the drug on milk production. Studies in lactating rats have shown excretion of brivaracetam or metabolites in milk [see *Data*].

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for brivaracetam and any potential adverse effects on the breastfed infant from brivaracetam or from the underlying maternal condition.

Data

Animal Data

Following a single oral (5 mg/kg) dose of ¹⁴C-brivaracetam to lactating rats, radioactivity was secreted in milk and rapidly reached levels similar to those in plasma.

8.4 Pediatric Use

Safety and effectiveness of brivaracetam have been established in pediatric patients 4 years to less than 16 years of age. Use of brivaracetam in these age groups is supported by evidence from adequate and well-controlled studies of brivaracetam in adults with partial-onset seizures, pharmacokinetic data from adult and pediatric patients and safety data in pediatric patients 4 years to less than 16 years of age [see *Dosage and Administration* (2.1), *Warnings and Precautions* (5.3), *Adverse Reactions* (6.1), *Clinical Pharmacology* (12.3) and *Clinical Studies* (14)].

Safety and effectiveness in pediatric patients below the age of 1 month have not been established.

Juvenile Animal Toxicity Data

The potential adverse effects of brivaracetam on postnatal growth and development were investigated in juvenile rats and dogs. Oral administration (0 mg/kg/day, 150 mg/kg/day, 300 mg/kg/day or 600 mg/kg/day) to rats during the neonatal and juvenile periods of development (approximately equivalent to neonatal through adolescent development in humans) resulted in increased mortality, decreased body weight gain, delayed male sexual maturation and adverse neurobehavioral effects at the highest dose tested and decreased brain size and weight at all doses. Therefore, a no-effect dose was not established; the lowest dose tested in juvenile rats was associated with plasma exposures (AUC) approximately 2 times those in children and adolescents at the recommended maintenance dose. In dogs, oral administration (0 mg/kg/day, 15 mg/kg/day, 30 mg/kg/day or 100 mg/kg/day) throughout the neonatal and juvenile periods of development induced liver changes similar to those observed in adult animals at the highest dose but produced no adverse effects on growth, bone density or strength, neurological testing or neuropathology evaluation. The overall no-effect dose (30 mg/kg/day) and the no-effect dose for adverse effects on developmental parameters (100 mg/kg/day) were associated with plasma exposures approximately equal to and 4 times, respectively, those in children and adolescents at the

recommended maintenance dose.

Additional pediatric use information is approved for UCB, Inc's Briviact (brivaracetam) tablets. However, due to UCB, Inc's marketing exclusivity rights, this drug product is not labeled with that pediatric information.

8.5 Geriatric Use

There were insufficient numbers of patients 65 years of age and older in the double-blind, placebo-controlled epilepsy trials (n=38) to allow adequate assessment of the effectiveness of brivaracetam in this population. In general, dose selection for an elderly patient should be judicious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function and of concomitant disease or other drug therapy [see *Clinical Pharmacology* (12.3)].

8.6 Renal Impairment

Dose adjustments are not required for patients with impaired renal function. There are no data in patients with end-stage renal disease undergoing dialysis and use of brivaracetam is not recommended in this patient population [see *Clinical Pharmacology* (12.3)].

8.7 Hepatic Impairment

Because of increases in brivaracetam exposure, dosage adjustment is recommended for all stages of hepatic impairment [see *Dosage and Administration* (2.5) and *Clinical Pharmacology* (12.3)].

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

Brivaracetam tablet contains brivaracetam and is listed as a Schedule V controlled substance.

9.2 Abuse

In a human abuse potential study, single doses of brivaracetam at therapeutic and supratherapeutic doses were compared to alprazolam (C-IV) (1.5 mg and 3 mg). Brivaracetam at the recommended single dose (50 mg) caused fewer sedative and euphoric effects than alprazolam; however, brivaracetam at supratherapeutic single doses (200 mg and 1,000 mg) was similar to alprazolam on other measures of abuse.

9.3 Dependence

There was no evidence of physical dependence potential or a withdrawal syndrome with brivaracetam in a pooled review of placebo-controlled adjunctive therapy studies [see *Warnings and Precautions* (5.5)].

10 OVERDOSAGE


There is limited clinical experience with brivaracetam overdose in humans. Somnolence

and dizziness were reported in a patient taking a single dose of 1,400 mg (14 times the highest recommended single dose) of brivaracetam. The following adverse reactions were reported with brivaracetam overdose: vertigo, balance disorder, fatigue, nausea, diplopia, anxiety and bradycardia. In general, the adverse reactions associated with brivaracetam overdose were consistent with the known adverse reactions.

There is no specific antidote for overdose with brivaracetam. In the event of overdose, standard medical practice for the management of any overdose should be used. An adequate airway, oxygenation and ventilation should be ensured; monitoring of cardiac rate and rhythm and vital signs is recommended. A certified poison control center should be contacted for updated information on the management of overdose with brivaracetam. There are no data on the removal of brivaracetam using hemodialysis, but because less than 10% of brivaracetam is excreted in urine, hemodialysis is not expected to enhance brivaracetam clearance.

11 DESCRIPTION

The chemical name of brivaracetam is (2S)-2-[(4R)-2-oxo-4-propyl-tetrahydro-1*H*-pyrrol-1-yl] butanamide. Its molecular formula is C₁₁H₂₀N₂O₂ and its molecular weight is 212.29. The chemical structure is:



Brivaracetam is a almost white to creamish yellow powder. It is very soluble in ethanol and glacial acetic acid. It is freely soluble in buffer (pH 1.2, 4.5, and 7.4), acetonitrile, toluene, acetone and isopropyl acetate and soluble in water and methanol. It is slightly soluble in n-hexane and diisopropyl ether.

Brivaracetam tablets are for oral administration and contain the following inactive ingredients: anhydrous lactose, croscarmellose sodium, glyceryl monocaprylocaprate type 1, lactose monohydrate, magnesium stearate, polyvinyl alcohol partially hydrolyzed, sodium lauryl sulfate, talc and titanium dioxide.

Additionally, each 25 mg tablet contains FD&C blue #2 Aluminum Lake and FD&C red #40 Aluminum Lake; each 75 mg tablet contains FD&C blue #1 Aluminum Lake and D&C red #27 Aluminum Lake and each 100 mg tablet contains FD&C yellow #5 Aluminum Lake, FD&C blue #1 Aluminum Lake and iron oxide yellow.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The precise mechanism by which brivaracetam exerts its anticonvulsant activity is not known. Brivaracetam displays a high and selective affinity for synaptic vesicle protein 2A(SV2A) in the brain, which may contribute to the anticonvulsant effect.

12.2 Pharmacodynamics

Interactions with Alcohol

In a pharmacokinetic and pharmacodynamic interaction study in healthy subjects, co-administration of brivaracetam (single dose 200 mg [2 times greater than the highest recommended single dose]) and ethanol (continuous intravenous infusion to achieve a blood alcohol concentration of 60 mg/100 mL during 5 hours) increased the effects of alcohol on psychomotor function, attention and memory. Co-administration of brivaracetam and ethanol caused a larger decrease from baseline in saccadic peak velocity, smooth pursuit, adaptive tracking performance and Visual Analog Scale (VAS) alertness and a larger increase from baseline in body sway and in saccadic reaction time compared with brivaracetam alone or ethanol alone. The immediate word recall scores were generally lower for brivaracetam when co-administered with ethanol.

Cardiac Electrophysiology

At a dose 4 times the maximum recommended dose, brivaracetam did not prolong the QT interval to a clinically relevant extent.

12.3 Pharmacokinetics

Brivaracetam tablets and oral solution can be used interchangeably. Brivaracetam exhibits linear and time-independent pharmacokinetics at the approved doses.

The pharmacokinetics of brivaracetam are similar when used as monotherapy or as adjunctive therapy for the treatment of partial-onset seizures.

Absorption

Brivaracetam is highly permeable and is rapidly and almost completely absorbed after oral administration. Pharmacokinetics is dose-proportional from 10 mg to 600 mg (a range that extends beyond the minimum and maximum single-administration dose levels described in Dosage and Administration [see *Dosage and Administration (2.1)*]). The median T_{max} for tablets taken without food is 1 hour (range 0.25 hour to 3 hours). Co-administration with a high-fat meal slowed absorption, but the extent of absorption remained unchanged. Specifically, when a 50 mg tablet was administered with a high-fat meal, C_{max} (maximum brivaracetam plasma concentration during a dose interval, an exposure metric) was decreased by 37% and T_{max} was delayed by 3 hours, but AUC (area under the brivaracetam plasma concentration versus time curve, an exposure metric) was essentially unchanged (decreased by 5%).

Distribution

Brivaracetam is weakly bound to plasma proteins ($\leq 20\%$). The volume of distribution is 0.5 L/kg, a value close to that of the total body water. Brivaracetam is rapidly and evenly distributed in most tissues.

Elimination

Metabolism

Brivaracetam is primarily metabolized by hydrolysis of the amide moiety to form the corresponding carboxylic acid metabolite and secondarily by hydroxylation on the propyl side chain to form the hydroxy metabolite. The hydrolysis reaction is mediated by hepatic and extra-hepatic amidase. The hydroxylation pathway is mediated primarily by CYP2C19. In human subjects possessing genetic variations in CYP2C19, production of the hydroxy metabolite is decreased 2-fold or 10-fold, while the blood level of brivaracetam itself is increased by 22% or 42%, respectively, in individuals with one or both mutated alleles. CYP2C19 poor metabolizers and patients using inhibitors of CYP2C19 may require dose reduction. An additional hydroxy acid metabolite is created by hydrolysis of the amide moiety on the hydroxy metabolite or hydroxylation of the propyl side chain on the carboxylic acid metabolite (mainly by CYP2C9). None of the 3 metabolites are pharmacologically active.

Excretion

Brivaracetam is eliminated primarily by metabolism and by excretion in the urine. More than 95% of the dose, including metabolites, is excreted in the urine within 72 hours after intake. Fecal excretion accounts for less than 1% of the dose. Less than 10% of the dose is excreted unchanged in the urine. Thirty-four percent of the dose is excreted as the carboxylic acid metabolite in urine. The terminal plasma half-life ($t_{1/2}$) is approximately 9 hours.

Specific Populations

Age

Pediatric Patients (4 years to less than 16 years): An open-label, single-arm, multicenter, pharmacokinetic study with a 3-week evaluation period and fixed 3-step up-titration using brivaracetam oral solution was conducted in 59 pediatric patients 4 years to less than 16 years of age. In those patients, plasma concentrations were shown to be dose-proportional. The pediatric pharmacokinetic profile for brivaracetam was determined in a population pharmacokinetic analysis using sparse plasma concentration data obtained in two open-label studies in 191 adult and pediatric patients with epilepsy 4 years to 17 years of age that received oral solution or oral tablet formulations.

A weight-based dosing regimen is necessary to achieve brivaracetam exposures in pediatric patients 4 years to less than 16 years of age that are similar to those observed in adults treated at effective doses of brivaracetam [see *Dosage and Administration (2.2)*]. The estimated plasma clearance was 1.61 L/h; 2.18 L/h and 3.19 L/h for pediatric patients weighing 20 kg, 30 kg and 50 kg, respectively. In comparison, plasma clearance was estimated at 3.58 L/h in adult patients (70 kg body weight).

Geriatric Population: In a study in elderly subjects (65 years to 79 years old; creatinine clearance 53 mL/min/1.73 m² to 98 mL/min/1.73 m²) receiving brivaracetam 200 mg twice daily (2 times the highest recommended dosage), the plasma half-life of brivaracetam was 7.9 hours and 9.3 hours in the 65 years to 75 and > 75 years groups, respectively. The steady-state plasma clearance of brivaracetam was slightly lower (0.76 mL/min/kg) than in young healthy controls (0.83 mL/min/kg).

Sex

There were no differences observed in the pharmacokinetics of brivaracetam between male and female subjects.

Race/Ethnicity

A population pharmacokinetic analysis comparing Caucasian and non-Caucasian patients showed no significant pharmacokinetic difference.

Renal Impairment

A study in adult subjects with severe renal impairment (creatinine clearance < 30 mL/min/1.73m² and not requiring dialysis) revealed that the plasma AUC of brivaracetam was moderately increased (21%) relative to healthy controls, while the AUCs of the acid, hydroxy and hydroxyacid metabolites were increased 3-fold, 4-fold and 21-fold, respectively. The renal clearance of these inactive metabolites was decreased 10-fold. Brivaracetam has not been studied in patients undergoing hemodialysis [see *Use in Specific Populations* (8.6)].

Hepatic Impairment

A pharmacokinetic study in adult subjects with hepatic cirrhosis, Child-Pugh grades A, B and C, showed 50%, 57% and 59% increases in brivaracetam exposure, respectively, compared to matched healthy controls. The effect of hepatic impairment on brivaracetam pharmacokinetics in pediatric patients is expected to be comparable to the effect observed in adults [see *Dosage and Administration* (2.5) and *Use in Specific Populations* (8.7)].

Drug Interaction Studies

In Vitro Assessment of Drug Interactions

Drug-Metabolizing Enzyme Inhibition

Brivaracetam did not inhibit CYP1A2, 2A6, 2B6, 2C8, 2C9, 2D6 or 3A4. Brivaracetam weakly inhibited CYP2C19 and would not be expected to cause significant inhibition of CYP2C19 in humans. Brivaracetam was an inhibitor of epoxide hydrolase, (IC₅₀ = 8.2 µM), suggesting that brivaracetam can inhibit the enzyme *in vivo*.

Drug-Metabolizing Enzyme Induction

Brivaracetam at concentrations up to 10 µM caused little or no change of mRNA expression of CYP1A2, 2B6, 2C9, 2C19, 3A4 and epoxide hydrolase. It is unlikely that brivaracetam will induce these enzymes *in vivo*.

Transporters

Brivaracetam was not a substrate of P-gp, MRP1 or MRP2. Brivaracetam did not inhibit or weakly inhibit BCRP, BSEP, MATE1, MATE2/K, MRP2, OAT1, OAT3, OCT1, OCT2, OATP1B1, OATP1B3 or P-gp, suggesting that brivaracetam is unlikely to inhibit these transporters *in vivo*.

In Vivo Assessment of Drug Interactions

Drug Interaction Studies with Antiepileptic Drugs (AEDs)

Potential interactions between brivaracetam (25 mg twice daily to 100 mg twice daily) and other AEDs were investigated in a pooled analysis of plasma drug concentrations from all Phase 2 and 3 studies and in a population exposure-response analysis of placebo-controlled, Phase 3 studies in adjunctive therapy in the treatment of partial-onset seizures. None of the interactions require changes in the dose of brivaracetam. Interactions with carbamazepine and phenytoin can be clinically important [see *Drug Interactions* (7.2) and (7.3)]. The interactions are summarized in Table 5.

Table 5 Drug Interactions Between Brivaracetam and Concomitant Antiepileptic Drugs

Concomitant AED	Influence of AED on Brivaracetam	Influence of Brivaracetam on AED
Carbamazepine	26% decrease in plasma concentration	None for carbamazepine Increase of carbamazepine-epoxide metabolite* [see <i>Drug Interactions (7.2)</i>]
Lacosamide	No data	None
Lamotrigine	None	None
Levetiracetam	None	None
Oxcarbazepine	None	None on the active monohydroxy metabolite derivative (MHD)
Phenobarbital	19% decrease in plasma concentration	None
Phenytoin	21% decrease in plasma concentration	Up to 20% increase in plasma concentration [see <i>Drug Interactions (7.3)</i>]**
Pregabalin	No data	None
Topiramate	None	None
Valproic acid	None	None
Zonisamide	No data	None

* Brivaracetam is a reversible inhibitor of epoxide hydrolase resulting in an increased concentration of carbamazepine epoxide, an active metabolite of carbamazepine. The carbamazepine epoxide plasma concentration increased up to 198% at a brivaracetam dose of 100 mg twice daily.

** At a supratherapeutic dose of 400 mg/day brivaracetam, there was a 20% increase in phenytoin plasma concentration.

Drug Interaction Studies with Other Drugs

Effect of Other Drugs on Brivaracetam

Co-administration with CYP inhibitors or transporter inhibitors is unlikely to significantly affect brivaracetam exposure.

Co-administration with rifampin decreases brivaracetam plasma concentrations by 45%, an effect that is probably the result of CYP2C19 induction [see *Dosage and Administration (2.6)* and *Drug Interactions (7.1)*].

Oral Contraceptives

Co-administration of brivaracetam 200 mg twice daily (twice the recommended maximum daily dosage) with an oral contraceptive containing ethinylestradiol (0.03 mg) and levonorgestrel (0.15 mg) reduced estrogen and progestin AUCs by 27% and 23%, respectively, without impact on suppression of ovulation. However, co-administration of brivaracetam 50 mg twice daily with an oral contraceptive containing ethinylestradiol (0.03 mg) and levonorgestrel (0.15 mg) did not significantly influence the

pharmacokinetics of either substance. The interaction is not expected to be of clinical significance.

Additional pediatric use information is approved for UCB, Inc's Briviact (brivaracetam) tablets. However, due to UCB, Inc's marketing exclusivity rights, this drug product is not labeled with that pediatric information.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

In a carcinogenicity study in mice, oral administration of brivaracetam (0 mg/kg/day, 400 mg/kg/day, 550 mg/kg/day or 700 mg/kg/day) for 104 weeks increased the incidence of liver tumors (hepatocellular adenoma and carcinoma) in male mice at the two highest doses tested. At the dose (400 mg/kg) not associated with an increase in liver tumors, plasma exposures (AUC) were approximately equal to those in humans at the maximum recommended dose (MRD) of 200 mg/day. Oral administration (0 mg/kg/day, 150 mg/kg/day, 230 mg/kg/day, 450 mg/kg/day or 700 mg/kg/day) to rats for 104 weeks resulted in an increased incidence of thymus tumors (benign thymoma) in female rats at the highest dose tested. At the highest dose not associated with an increase in thymus tumors, plasma exposures were approximately 9 times those in humans at the MRD.

Mutagenesis

Brivaracetam was negative for genotoxicity in *in vitro* (Ames, mouse lymphoma and CHO chromosomal aberration) and *in vivo* (rat bone marrow micronucleus) assays.

Impairment of Fertility

Oral administration of brivaracetam (0 mg/kg/day, 100 mg/kg/day, 200 mg/kg/day or 400 mg/kg/day) to male and female rats prior to and throughout mating and early gestation produced no adverse effects on fertility. The highest dose tested was associated with plasma exposures approximately 6 (males) and 13 (females) times those in humans at the MRD.

14 CLINICAL STUDIES

The effectiveness of brivaracetam in partial-onset seizures with or without secondary generalization was established in 3 fixed-dose, randomized, double-blind, placebo-controlled, multicenter studies (Studies 1, 2 and 3), which included 1,550 patients. Patients enrolled had partial-onset seizures that were not adequately controlled with 1 to 2 concomitant antiepileptic drugs (AEDs). In each of these studies, 72% to 86% of patients were taking 2 or more concomitant AEDs with or without vagal nerve stimulation. The median baseline seizure frequency across the 3 studies was 9 seizures per 28 days. Patients had a mean duration of epilepsy of approximately 23 years.

All trials had an 8-week baseline period, during which patients were required to have at least 8 partial-onset seizures. The baseline period was followed by a 12-week treatment period. There was no titration period in these studies. Study 1 compared doses of brivaracetam 50 mg/day and 100 mg/day with placebo. Study 2 compared a dose of brivaracetam 50 mg/day with placebo. Study 3 compared doses of brivaracetam 100

mg/day and 200 mg/day with placebo. Brivaracetam was administered in equally divided twice daily doses. Upon termination of brivaracetam treatment, patients were down-titrated over a 1-week, 2-week and 4-week duration for patients receiving 25 mg, 50 mg and 100 mg twice daily brivaracetam, respectively.

The primary efficacy outcome in Study 1 and Study 2 was the percent reduction in 7-day partial-onset seizure frequency over placebo, while the primary outcome for Study 3 was the percent reduction in 28-day partial-onset seizure frequency over placebo. The criteria for statistical significance for all 3 studies was $p<0.05$. Table 6 presents the primary efficacy outcome of the percent change in seizure frequency over placebo, based upon each study's protocol-defined 7-day and 28-day seizure frequency efficacy outcome.

Table 6 Percent Reduction in Partial-Onset Seizure Frequency over Placebo (Studies 1, 2 and 3)

	Percent Reduction Over Placebo (%)
STUDY 1^a	
Placebo (n=100)	-
50 mg/day (n=99)	9.5
100 mg/day (n=100)	17
STUDY 2^a	
Placebo (n=96)	-
50 mg/day (n=101)	16.9*
STUDY 3^b	
Placebo (n=259)	-
100 mg/day (n=252)	25.2*
200 mg/day (n= 249)	25.7*

*Statistically significant based on testing procedure with alpha = 0.05


^aBased upon 7-day seizure frequency

^bBased upon 28-day seizure frequency

Figure 1 presents the percentage of patients by category of reduction from baseline in partial-onset seizure frequency per 28 days for all pooled patients in the 3 pivotal studies. Patients in whom the seizure frequency increased are shown at left as "worse." Patients with an improvement in percent reduction from baseline partial-onset seizure frequency are shown in the 4 right-most categories.

Figure 1

Proportion of Patients by Category of Seizure Response for Brivaracetam and Placebo Across all Three Double-Blind Trials



Treatment with Levetiracetam

In Studies 1 and 2, which evaluated brivaracetam dosages of 50 mg and 100 mg daily, approximately 20% of the patients were on concomitant levetiracetam. Although the numbers of patients were limited, brivaracetam provided no added benefit when it was added to levetiracetam.

Although patients on concomitant levetiracetam were excluded from Study 3, which evaluated 100 mg and 200 mg daily, approximately 54% of patients in this study had prior exposure to levetiracetam.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Brivaracetam tablets, 10 mg are white to off-white, round, biconvex, film-coated tablets debossed with "U" on one side and plain on the other and are supplied as follows:

NDC 70710-1584-6 in bottles of 60 tablets with child-resistant closure.

NDC 70710-1584-8 in bottles of 180 tablets with child-resistant closure.

NDC 70710-1584-4 in unit-dose blister cartons of 100 tablets (100 x 1 unit-dose)

Brivaracetam tablets, 25 mg are light blue, oval, biconvex, film-coated tablets debossed with "U1" on one side and plain on the other and are supplied as follows:

NDC 70710-1585-6 in bottles of 60 tablets with child-resistant closure.

NDC 70710-1585-8 in bottles of 180 tablets with child-resistant closure.

NDC 70710-1585-4 in unit-dose blister cartons of 100 tablets (100 x 1 unit-dose)

Brivaracetam tablets, 50 mg are white to off-white, oval, biconvex, film-coated tablets debossed with "U1" on one side and plain on the other and are supplied as follows:

NDC 70710-1586-6 in bottles of 60 tablets with child-resistant closure.

NDC 70710-1586-8 in bottles of 180 tablets with child-resistant closure.

NDC 70710-1586-4 in unit-dose blister cartons of 100 tablets (100 x 1 unit-dose)

Brivaracetam tablets, 75 mg are purple, oval, biconvex, film-coated tablets debossed with "1587" on one side and plain on the other and are supplied as follows:

NDC 70710-1587-6 in bottles of 60 tablets with child-resistant closure.

NDC 70710-1587-8 in bottles of 180 tablets with child-resistant closure.

NDC 70710-1587-4 in unit-dose blister cartons of 100 tablets (100 x 1 unit-dose)

Brivaracetam tablets, 100 mg are light green, oval, biconvex, film-coated tablets debossed with "1588" on one side and plain on the other and are supplied as follows:

NDC 70710-1588-6 in bottles of 60 tablets with child-resistant closure.

NDC 70710-1588-8 in bottles of 180 tablets with child-resistant closure.

NDC 70710-1588-4 in unit-dose blister cartons of 100 tablets (100 x 1 unit-dose)

16.2 Storage and Handling

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C to 30°C (59°F to 86°F) [See USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

Suicidal Behavior and Ideation

Counsel patients, their caregivers and/or families that antiepileptic drugs, including brivaracetam, may increase the risk of suicidal thoughts and behavior and advise patients to be alert for the emergence or worsening of symptoms of depression; unusual changes in mood or behavior; or suicidal thoughts, behavior or thoughts about self-harm. Advise patients, their caregivers and/or families to report behaviors of concern immediately to a healthcare provider [see *Warnings and Precautions (5.1)*].

Neurological Adverse Reactions

Counsel patients that brivaracetam causes somnolence, fatigue, dizziness and gait disturbance. These adverse reactions, if observed, are more likely to occur early in treatment but can occur at any time. Advise patients not to drive or operate machinery

until they have gained sufficient experience on brivaracetam to gauge whether it adversely affects their ability to drive or operate machinery [see *Warnings and Precautions* (5.2)].

Psychiatric Adverse Reactions

Advise patients that brivaracetam causes changes in behavior (e.g., aggression, agitation, anger, anxiety and irritability) and psychotic symptoms. Instruct patients to report these symptoms immediately to their healthcare provider [see *Warnings and Precautions* (5.3)].

Hypersensitivity: Bronchospasm and Angioedema

Advise patients that symptoms of hypersensitivity including bronchospasm and angioedema can occur with brivaracetam. Instruct them to seek immediate medical care should they experience signs and symptoms of hypersensitivity [see *Warnings and Precautions* (5.4)].

Withdrawal of Antiepileptic Drugs

Advise patients not to discontinue use of brivaracetam without consulting with their healthcare provider. Brivaracetam should normally be gradually withdrawn to reduce the potential for increased seizure frequency and status epilepticus [see *Warnings and Precautions* (5.5)].

Pregnancy

Advise patients to notify their healthcare provider if they become pregnant or intend to become pregnant during brivaracetam therapy. Encourage patients to enroll in the North American Antiepileptic Drug Pregnancy Registry if they become pregnant. This registry is collecting information about the safety of antiepileptic drugs during pregnancy [see *Use in Specific Populations* (8.1)].

Dosing Instructions

Counsel patients that brivaracetam may be taken with or without food. Instruct patients that brivaracetam tablets should be swallowed whole with liquid and not chewed or crushed [see *Dosage and Administration* (2.2)].

Medication Guide available at www.zydususa.com/medguides or call 1-877-993-8779.

Manufactured by:

Zydus Lifesciences Ltd.

Ahmedabad, India.

Distributed by:

Zydus Pharmaceuticals (USA) Inc.

Pennington, NJ 08534

Rev.: 09/22

MEDICATION GUIDE

Brivaracetam (briv" a ra' se tam) tablets, CV

What is the most important information I should know about

brivaracetam tablets?

Brivaracetam tablets are a federally controlled substance (CV) because it can be abused or lead to dependence. Keep brivaracetam tablets in a safe place to prevent misuse and abuse. Selling or giving away brivaracetam tablets may harm others and is against the law.

Like other antiepileptic drugs, brivaracetam tablets may cause suicidal thoughts or actions in a very small number of people, about 1 in 500 people taking it.

Call a healthcare provider right away if you have any of these symptoms, especially if they are new, worse or worry you:

- thoughts about suicide or dying
- new or worse depression
- feeling agitated or restless
- trouble sleeping (insomnia)
- acting aggressive, feeling angry or being violent
- an extreme increase in activity and talking (mania)
- attempts to commit suicide
- new or worse anxiety
- panic attacks
- new or worse irritability
- acting on dangerous impulses
- other unusual changes in behavior or mood

Suicidal thoughts or actions can be caused by things other than medicines. If you have suicidal thoughts or actions, your healthcare provider may check for other causes.

How can I watch for early symptoms of suicidal thoughts and actions?

- Pay attention to any changes, especially sudden changes, in mood, behaviors, thoughts or feelings.
- Keep all follow-up visits with your healthcare provider as scheduled.

Call your healthcare provider between visits as needed, especially if you are worried about symptoms.

Do not stop brivaracetam tablets without first talking to a healthcare provider.

- Stopping brivaracetam tablets suddenly can cause serious problems.
- Stopping a seizure medicine suddenly can cause seizures that will not stop (status epilepticus).

What are brivaracetam tablets?

Brivaracetam tablets are a prescription medicine used to treat partial-onset seizures in people 4 years of age and older.

It is not known if brivaracetam tablets is safe and effective in children younger than 1 month of age.

Who should not take brivaracetam tablets?

Do not take brivaracetam tablets if you are allergic to brivaracetam or any of the ingredients in brivaracetam tablets. See the end of this Medication Guide for a complete list of ingredients in brivaracetam tablets.

What should I tell my healthcare provider before starting brivaracetam tablets?

Before taking brivaracetam tablets, tell your healthcare provider about all of your medical conditions, including if you:

- have or had depression, mood problems or suicidal thoughts or behavior.
- have liver problems.
- have abused or been dependent on prescription medicines, street drugs or alcohol.
- are pregnant or plan to become pregnant. It is not known if brivaracetam tablets will harm your unborn baby. Tell your healthcare provider right away if you become pregnant while taking brivaracetam tablets. You and your healthcare provider will have to decide if you should take brivaracetam tablets while you are pregnant. If you become pregnant while taking brivaracetam tablets, talk to your healthcare provider about registering with the North American Antiepileptic Drug Pregnancy Registry. You can enroll in this registry by calling 1-888-233-2334. The purpose of this registry is to collect information about the safety of brivaracetam tablets and other antiepileptic medicines during pregnancy.
- are breastfeeding or plan to breastfeed. It is not known if brivaracetam passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you take brivaracetam tablets.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements. Brivaracetam tablets may affect the way other medicines work and other medicines may affect how brivaracetam tablet works. Do not start a new medicine without first talking with your healthcare provider. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist each time you get a new medicine.

How should I take brivaracetam tablets?

- Take brivaracetam tablets exactly as your healthcare provider tells you.
- Your healthcare provider will tell you how much brivaracetam tablets to take and when to take it.
- Your healthcare provider may change your dose if needed. Do not change your dose without talking to your healthcare provider.
- Take brivaracetam tablets with or without food.
- Swallow brivaracetam tablets whole with a liquid. Do not chew or crush brivaracetam tablets before swallowing.
- If you take too much brivaracetam tablets, call your Poison Control Center or go to the nearest emergency room right away.

What should I avoid while taking brivaracetam tablets?

Do not drive or operate machinery until you know how brivaracetam

tablets affect you. Brivaracetam tablets may cause drowsiness, tiredness, dizziness and problems with your balance and coordination.

What are the possible side effects of brivaracetam tablets?

Brivaracetam tablets may cause serious side effects, including:

- See "**What is the most important information I should know about brivaracetam tablets?**"
- **Nervous system problems.** Drowsiness, tiredness and dizziness are common with brivaracetam tablets, but can be severe. See "**What should I avoid while taking brivaracetam tablets?**" brivaracetam tablets can also cause problems with balance and coordination.
- **Mental (psychiatric) symptoms.** Brivaracetam tablets can cause mood and behavior changes such as aggression, agitation, anger, anxiety, apathy, mood swings, depression, hostility and irritability. Irritability and anxiety are common with brivaracetam tablets and can be severe. People who take brivaracetam tablets can also get psychotic symptoms such as hallucinations (seeing or hearing things that are really not there), delusions (false or strange thoughts or beliefs) and unusual behavior.

The most common side effects of brivaracetam tablets in adults include:

- sleepiness
- dizziness
- feeling tired
- nausea and vomiting

Side effects of brivaracetam tablets in children 4 years to less than 16 years of age are similar to those seen in adults.

These are not all the possible side effects of brivaracetam tablets. For more information, ask your healthcare provider or pharmacist. Tell your healthcare provider about any side effect that bothers you or that does not go away. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store brivaracetam tablets?

- Store brivaracetam tablets at room temperature between 59°F to 86°F (15°C to 30°C).
- Drug brivaracetam tablets come in child-resistant bottles of 60's and 180's.

Keep brivaracetam tablets and all medicines out of the reach of children.

General information about the safe and effective use of brivaracetam tablets.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use brivaracetam tablets for a condition for which it was not prescribed. Do not give brivaracetam tablets to other people, even if they have the same symptoms that you have. It may harm them.

This Medication Guide summarizes the most important information about

brivaracetam tablets. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about brivaracetam tablets that is written for health professionals.

What are the ingredients in brivaracetam tablets?

Active ingredient: brivaracetam

Inactive ingredients: anhydrous lactose, croscarmellose sodium, glyceryl monocaprylocaprate type 1, lactose monohydrate, magnesium stearate, polyvinyl alcohol partially hydrolyzed, sodium lauryl sulfate, talc and titanium dioxide.

Additionally, each 25 mg tablet contains FD&C blue #2 Aluminum Lake and FD&C red #40 Aluminum Lake, 75 mg tablet contains FD&C blue #1 Aluminum Lake and D&C red #27 Aluminum Lake and 100 mg tablet contains FD&C yellow #5 Aluminum Lake, FD&C blue #1 Aluminum Lake and iron oxide yellow.

For more information, go to MedicalAffairs@zydususa.com or Tel.: 1-877-993-8779.

Medication Guide available at www.zydususa.com/medguides or call 1-877-993-8779.

Additional pediatric use information is approved for UCB, Inc's Briviact (brivaracetam) tablets. However, due to UCB, Inc's marketing exclusivity rights, this drug product is not labeled with that pediatric information.

Manufactured by:
Zydus Lifesciences Ltd.

Ahmedabad, India.

Distributed by:
Zydus Pharmaceuticals (USA) Inc.
Pennington, NJ 08534

Rev: 05/22

This Medication Guide has been approved by the U.S. Food and Drug Administration

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 70710-1584-6

Brivaracetam tablets, 10 mg

60 Tablets

Rx only

Zydus



GTIN 00000000000000
SN 00000000000000
EXP DDMMYYYY
LOT XXXXXX

Over Coding Template
No Varnished Area (Do Not Print)
(18 x 41 mm)

NDC 70710-1584-6

Brivaracetam Tablets

10 mg

PHARMACIST: Dispense the Medication
Guide provided separately to each patient.



60 Tablets
Rx only

Each film-coated tablet contains
10 mg brivaracetam.

Usual Dosage: See accompanying
prescribing information.

This package is child-resistant.


Store at 20°C to 25°C (68°F to 77°F);
excursions permitted between 15°C
to 30°C (59°F to 86°F) [See USP
Controlled Room Temperature].

Keep this and all drugs out of
the reach of children.

Medication Guide available at
www.zydususa.com/medguides or
call 1-877-993-8779.

Mfg. by: Zydus Lifesciences Ltd.
Ahmedabad, India

Dist. by: Zydus Pharmaceuticals (USA) Inc.
Pennington, NJ 08534



Rev.: 05/22

NDC 70710-1585-6

Brivaracetam tablets, 25 mg

60 Tablets

Rx only

Zydus



GTIN 00000000000000
SN 00000000000000
EXP DDMMYYYY
LOT XXXXXX

Over Coding Template
No Varnished Area (Do Not Print)
(18 x 41 mm)

NDC 70710-1585-6

Brivaracetam Tablets

25 mg

PHARMACIST: Dispense the Medication
Guide provided separately to each patient.



Rev.: 05/22



60 Tablets
Rx only

Each film-coated tablet contains
25 mg brivaracetam.

Usual Dosage: See accompanying
prescribing information.

This package is child-resistant.

Store at 20°C to 25°C (68°F to 77°F);
excursions permitted between 15°C
to 30°C (59°F to 86°F) [See USP
Controlled Room Temperature].

Keep this and all drugs out of
the reach of children.

Medication Guide available at
www.zydususa.com/medguides or
call 1-877-993-8779.

Mfg. by: Zydus Lifesciences Ltd.
Ahmedabad, India

Dist. by: Zydus Pharmaceuticals (USA) Inc.
Pennington, NJ 08534

NDC 70710-1586-6

Brivaracetam tablets, 50 mg

60 Tablets

Rx only

Zydus



GTIN 00000000000000
SN 00000000000000
EXP DDMMYYYY
LOT XXXXXXXX

Over Coding Template
No Varnished Area (Do Not Print)
(18 x 41 mm)

NDC 70710-1586-6

Brivaracetam Tablets

50 mg

PHARMACIST: Dispense the Medication
Guide provided separately to each patient.



60 Tablets
Rx only

Each film-coated tablet contains
50 mg brivaracetam.

Usual Dosage: See accompanying
prescribing information.

This package is child-resistant.

Store at 20°C to 25°C (68°F to 77°F);
excursions permitted between 15°C
to 30°C (59°F to 86°F) [See USP
Controlled Room Temperature].

Keep this and all drugs out of
the reach of children.

Medication Guide available at
www.zydususa.com/medguides or
call 1-877-993-8779.

Mfg. by: Zydus Lifesciences Ltd.
Ahmedabad, India

Dist. by: Zydus Pharmaceuticals (USA) Inc.
Pennington, NJ 08534



Rev.: 05/22


NDC 70710-1587-6

Brivaracetam tablets, 75 mg

60 Tablets

Rx only

Zydus



Over Coding Template
No Varnished Area (Do Not Print)
(18 x 41 mm)

NDC 70710-1587-6

Brivaracetam Tablets

75 mg

PHARMACIST: Dispense the Medication
Guide provided separately to each patient.



60 Tablets
Rx only

Each film-coated tablet contains
75 mg brivaracetam.

Usual Dosage: See accompanying
prescribing information.

This package is child-resistant.
Store at 20°C to 25°C (68°F to 77°F);
excursions permitted between 15°C
to 30°C (59°F to 86°F) [See USP
Controlled Room Temperature].

**Keep this and all drugs out of
the reach of children.**

Medication Guide available at
www.zydususa.com/medguides or
call 1-877-993-8779.

Mfg. by: Zydus Lifesciences Ltd.
Ahmedabad, India
Dist. by: Zydus Pharmaceuticals (USA) Inc.
Pennington, NJ 08534



Rev.: 05/22

NDC 70710-1588-6

Brivaracetam tablets, 100 mg

60 Tablets

Rx only

Zydus



GTIN 00000000000000
SN 00000000000000
EXP DDMMYYYY
LOT XXXXXX

Over Coding Template
No Varnished Area (Do Not Print)
(18 x 41 mm)



Rev.: 05/22

NDC 70710-1588-6

Brivaracetam Tablets

100 mg

PHARMACIST: Dispense the Medication
Guide provided separately to each patient.



60 Tablets
Rx only

Each film-coated tablet contains
100 mg brivaracetam.

Contains FD&C Yellow No. 5 (tartrazine)
as a color additive.

Usual Dosage: See accompanying
prescribing information.

This package is child-resistant.
Store at 20°C to 25°C (68°F to 77°F);
excursions permitted between 15°C
to 30°C (59°F to 86°F) [See USP
Controlled Room Temperature].

Keep this and all drugs out of
the reach of children.

Medication Guide available at
www.zydususa.com/medguides or
call 1-877-993-8779.

Mfg. by: Zydus Lifesciences Ltd.
Ahmedabad, India

Dist. by: Zydus Pharmaceuticals (USA) Inc.
Pennington, NJ 08534

BRIVARACETAM

brivaracetam tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70710-1584
Route of Administration	ORAL	DEA Schedule	CV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BRIVARACETAM (UNII: U863JGG21A) (BRIVARACETAM - UNII:U863JGG21A)	BRIVARACETAM	10 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
GLYCERYL MONO AND DICAPRYLOCAPRATE (UNII: U72Q2I8C85)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	

TALC (UNII: 7SEV7J4R1U)**TITANIUM DIOXIDE** (UNII: 15FIX9V2JP)**Product Characteristics**

Color	WHITE (off-white)	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	U
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70710-1584-6	60 in 1 BOTTLE; Type 0: Not a Combination Product	01/21/2026	
2	NDC:70710-1584-8	180 in 1 BOTTLE; Type 0: Not a Combination Product	01/21/2026	
3	NDC:70710-1584-4	100 in 1 CARTON	01/21/2026	
3	NDC:70710-1584-7	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214501	01/21/2026	

BRIVARACETAM

brivaracetam tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDc:70710-1585
Route of Administration	ORAL	DEA Schedule	CV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BRIVARACETAM (UNII: U863JGG2IA) (BRIVARACETAM - UNII:U863JGG2IA)	BRIVARACETAM	25 mg

Inactive Ingredients

Ingredient Name	Strength
ALUMINUM OXIDE (UNII: LMI26O6933)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	

FD&C BLUE NO. 2 (UNII: L06K8R7DQK)
FD&C RED NO. 40 (UNII: WZB9127XOA)
GLYCERYL MONO AND DICAPRYLOCAPRATE (UNII: U72Q2I8C85)
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)
MAGNESIUM STEARATE (UNII: 70097M6I30)
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)
SODIUM LAURYL SULFATE (UNII: 368GB5141J)
TALC (UNII: 7SEV7J4R1U)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

Color	BLUE (light blue)	Score	no score
Shape	OVAL	Size	9mm
Flavor		Imprint Code	U1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70710-1585-6	60 in 1 BOTTLE; Type 0: Not a Combination Product	01/21/2026	
2	NDC:70710-1585-8	180 in 1 BOTTLE; Type 0: Not a Combination Product	01/21/2026	
3	NDC:70710-1585-4	100 in 1 CARTON	01/21/2026	
3	NDC:70710-1585-7	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214501	01/21/2026	

BRIVARACETAM

brivaracetam tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70710-1586
Route of Administration	ORAL	DEA Schedule	CV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BRIVARACETAM (UNII: U863JGG2IA) (BRIVARACETAM - UNII:U863JGG2IA)	BRIVARACETAM	50 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
GLYCERYL MONO AND DICAPRYLOCAPRATE (UNII: U72Q2I8C85)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE (off-white)	Score	no score
Shape	OVAL	Size	12mm
Flavor		Imprint Code	U11
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70710-1586-6	60 in 1 BOTTLE; Type 0: Not a Combination Product	01/21/2026	
2	NDC:70710-1586-8	180 in 1 BOTTLE; Type 0: Not a Combination Product	01/21/2026	
3	NDC:70710-1586-4	100 in 1 CARTON	01/21/2026	
3	NDC:70710-1586-7	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214501	01/21/2026	

BRIVARACETAM

brivaracetam tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70710-1587
Route of Administration	ORAL	DEA Schedule	CV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BRIVARACETAM (UNII: U863JGG2IA) (BRIVARACETAM - UNII:U863JGG2IA)	BRIVARACETAM	75 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
FD&C BLUE NO. 1--ALUMINUM LAKE (UNII: J9EQA3S2JM)	
GLYCERYL MONO AND DICAPRYLOCAPRATE (UNII: U72Q2I8C85)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	PURPLE	Score	no score
Shape	OVAL	Size	13mm
Flavor		Imprint Code	1587
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70710-1587-6	60 in 1 BOTTLE; Type 0: Not a Combination Product	01/21/2026	
2	NDC:70710-1587-8	180 in 1 BOTTLE; Type 0: Not a Combination Product	01/21/2026	
3	NDC:70710-1587-4	100 in 1 CARTON	01/21/2026	
3	NDC:70710-1587-7	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214501	01/21/2026	

BRIVARACETAM

brivaracetam tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70710-1588
Route of Administration	ORAL	DEA Schedule	CV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BRIVARACETAM (UNII: U863JGG2IA) (BRIVARACETAM - UNII:U863JGG2IA)	BRIVARACETAM	100 mg

Inactive Ingredients

Ingredient Name	Strength
ALUMINUM OXIDE (UNII: LMI2606933)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 1--ALUMINUM LAKE (UNII: J9EQA3S2JM)	
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
GLYCERYL MONO AND DICAPRYLOCAPRATE (UNII: U72Q2I8C85)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	GREEN (light-green)	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	1588
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70710-1588-6	60 in 1 BOTTLE; Type 0: Not a Combination Product	01/21/2026	
2	NDC:70710-1588-8	180 in 1 BOTTLE; Type 0: Not a Combination Product	01/21/2026	
3	NDC:70710-1588-4	100 in 1 CARTON	01/21/2026	
3	NDC:70710-1588-7	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214501	01/21/2026	

Labeler - Zydus Pharmaceuticals USA Inc. (156861945)

Registrant - Zydus Pharmaceuticals USA Inc. (156861945)

Establishment

Name	Address	ID/FEI	Business Operations
Zydus Lifesciences Limited		863362789	ANALYSIS(70710-1584, 70710-1585, 70710-1586, 70710-1587, 70710-1588) , MANUFACTURE(70710-1584, 70710-1585, 70710-1586, 70710-1587, 70710-1588)

Revised: 9/2022

Zydus Pharmaceuticals USA Inc.