

**SINUS RELIEF CONGESTION AND COUGH MAXIMUM STRENGTH-  
acetaminophen, diphenhydramine hci, phenylephrine hcl liquid  
CVS Pharmacy**

-----  
**Drug Facts**

**Active ingredients (in each 20 mL)**

Acetaminophen 650 mg

Diphenhydramine HCl 25 mg

Phenylephrine HCl 10 mg

**Purposes**

Pain Reliever

Antihistamine / cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves
  - nasal congestion
  - headache
  - minor aches and pains
  - cough
  - sinus congestion and pressure
  - runny nose and sneezing
  - temporarily promotes nasal and/or sinus drainage
  - controls cough to help you get to sleep

**Warnings**

**Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

### **Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin
- for children under 12 years of age

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a Doctor or Pharmacist before use if

- you are taking the blood thinning drug warfarin
- you are taking sedatives or tranquilizers

### **Ask a Doctor or pharmacist before use if**

- you are taking the blood thinning drug warfarin
- you are taking sedatives or tranquilizers

### **When using this product,**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

- cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warnings:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- dose as follows or as directed by a doctor
- adults and children 12 years and older:
  - 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

### **Other information**

- **each 20 mL contains:** sodium 12 mg
- store between 20-25°C (68°-77°F). Do not refrigerate.

### **Inactive ingredients**

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, sodium citrate, xanthan gum

### **Questions or comments?**

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

### **Principal Display Panel**

Compare to the active ingredients in Maximum Strength Mucinex® SINUS-MAX Night Time Congestion & Cough\*

Nighttime

### **Severe Cough & Congestion Relief**

ACETAMINOPHEN - Pain reliever

DIPHENHYDRAMINE HCl / Antihistamine / Cough suppressant

PHENYLEPHRINE HCl - Nasal decongestant

## **MAXIMUM STRENGTH**

### **Multi-Symptom**

- Clears sinus congestion
- Relieves headache
- relieves runny nose & sneezing

For Ages 12 & Over

FL OZ (mL)

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® SINUS-MAX® Night Time Congestion & Cough.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

**Distributed by: CVS Pharmacy, Inc.**

One CVS Drive, Woonsocket, RI 02895

CVS.com

**Package Label**

**CVS Health.** Compare to the active ingredients in Maximum Strength Mucinex® SINUS-MAX® Night Time Congestion & Cough\* NDC 69842-461-06

# Nighttime Severe Congestion Relief

**ACETAMINOPHEN**  
Pain reliever

**DIPHENHYDRAMINE HCl**  
Antihistamine; Cough suppressant

**PHENYLEPHRINE HCl**  
Nasal decongestant

**MAXIMUM STRENGTH**  
Multi-Symptom

- Clears sinus congestion
- Relieves headache
- Relieves runny nose & sneezing

For Ages 12 & Over

**6 FL OZ (177 mL)**

P.L.D-A378C LB007739

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® SINUS-MAX® Night Time Congestion & Cough.

#547736



PLD-A378C LB007740

**Distributed by: CVS Pharmacy, Inc.**  
One CVS Drive, Woonsocket, RI 02895  
© 2020 CVS/pharmacy  
CVS.com 1-800-SHOP CVS  
V-13114



**PEEL CORNER FOR DRUG FACTS**

### Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg	Pain reliever
Diphenhydramine HCl 25 mg	Antihistamine/cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant

### Uses

- temporarily relieves
  - nasal congestion
  - minor aches and pains
  - sinus congestion and pressure
  - runny nose and sneezing
- temporarily promotes nasal and/or sinus drainage
- controls cough to help you get to sleep

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

### Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

#### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- diabetes
- thyroid disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

### Drug Facts (continued)

#### Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin
- you are taking sedatives or tranquilizers

#### When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or

### Drug Facts (continued)

contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- dose as follows or as directed by a doctor
- adults and children 12 years and older: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

#### Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

#### Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

#### Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

**PEEL CORNER FOR MORE DRUG FACTS**

## CVS HEALTH Nighttime Severe Congestion Relief

# SINUS RELIEF CONGESTION AND COUGH MAXIMUM STRENGTH

acetaminophen, diphenhydramine hci, phenylephrine hcl liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69842-461
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg in 20 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYL GALLATE</b> (UNII: 8D4SNN7V92)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:69842-461-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/30/2018	

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M012	04/30/2018	

**Labeler** - CVS Pharmacy (062312574)

Revised: 3/2024

CVS Pharmacy