

GALANTAMINE HYDROBROMIDE- galantamine hydrobromide solution

Hikma Pharmaceuticals USA Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use GALANTAMINE ORAL SOLUTION safely and effectively. See full prescribing information for GALANTAMINE ORAL SOLUTION.

GALANTAMINE oral solution
Initial U.S. Approval: 2001

INDICATIONS AND USAGE

Galantamine is a cholinesterase inhibitor indicated for the treatment of mild to moderate dementia of the Alzheimer's type. (1)

DOSAGE AND ADMINISTRATION

- Galantamine Oral Solution: Recommended starting dosage is 4 mg (1 mL) twice daily; increase dose to initial maintenance dosage of 8 mg (2 mL) twice daily after a minimum of 4 weeks. Based on clinical benefit and tolerability, dosage may be increased to 12 mg (3 mL) twice daily after a minimum of 4 weeks at 8 mg (2 mL) twice daily. (2.2)
- Take with meals; ensure adequate fluid intake during treatment. (2.2)
- Hepatic Impairment: should not exceed 16 mg (4 mL)/day for moderate hepatic impairment; do not use in patients with severe hepatic impairment. (2.3)
- Renal Impairment: should not exceed 16 mg (4 mL)/day for creatinine clearance 9 mL to 59 mL/min; do not use in patients with creatinine clearance less than 9 mL/min. (2.4)

DOSAGE FORMS AND STRENGTHS

Oral Solution – 4 mg/mL (3)

CONTRAINDICATIONS

Known hypersensitivity to galantamine hydrobromide or any excipients. (4)

WARNINGS AND PRECAUTIONS

- Serious Skin Reactions: discontinue at first appearance of skin rash. (5.1)
- All patients should be considered at risk for adverse effects on cardiac conduction, including bradycardia and AV block, due to vagotonic effects on sinoatrial and atrioventricular nodes. (5.3)
- Active or Occult Gastrointestinal Bleeding: monitor, especially those with an increased risk for developing ulcers. (5.4)
- Cholinomimetics may cause bladder outflow obstruction. (5.5)
- Monitor for respiratory adverse events in patients with a history of severe asthma or obstructive pulmonary disease. (5.7)

ADVERSE REACTIONS

The most common adverse reactions ($\geq 5\%$) were nausea, vomiting, diarrhea, dizziness, headache, and decreased appetite. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Hikma Pharmaceuticals USA Inc. at 1-800-962-8364 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Potential to interfere with the activity of anticholinergic medications. (7.1)
- Synergistic effect expected when given concurrently with succinylcholine, other cholinesterase inhibitors, similar neuromuscular blocking agents, or cholinergic agonists. (7.2)

USE IN SPECIFIC POPULATIONS

Pregnancy: Based on animal data may cause fetal harm. (8.1)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 10/2023

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.2 Galantamine Oral Solution
- 2.3 Dosage in Patients with Hepatic Impairment
- 2.4 Dosage in Patients with Renal Impairment

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Serious Skin Reactions
- 5.2 Anesthesia
- 5.3 Cardiovascular Conditions
- 5.4 Gastrointestinal Conditions
- 5.5 Genitourinary Conditions
- 5.6 Neurological Conditions
- 5.7 Pulmonary Conditions
- 5.8 Deaths in Subjects with Mild Cognitive Impairment (MCI)

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Use with Anticholinergics
- 7.2 Use with Cholinomimetics and Other Cholinesterase Inhibitors

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment
- 8.7 Renal Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

- 14.1 Study Outcome Measures
- 14.2 Immediate-Release Tablets

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Storage and Handling

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Galantamine is indicated for the treatment of mild to moderate dementia of the Alzheimer's type.

2 DOSAGE AND ADMINISTRATION

2.2 Galantamine Oral Solution

The dosage of galantamine tablets shown to be effective in controlled clinical trials is 16 mg to 32 mg/day given as twice daily dosing. As the dosage of 32 mg/day is less well tolerated than lower dosages and does not provide increased effectiveness, the recommended dosage range is 16 mg to 24 mg/day given twice daily. The dosage of 24 mg/day did not provide a statistically significant greater clinical benefit than 16 mg/day. It is possible, however, that a daily dosage of 24 mg of galantamine might provide additional benefit for some patients.

The recommended starting dosage of galantamine oral solution is 4 mg (1 mL) twice a day (8 mg (2 mL)/day). The dosage should be increased to the initial maintenance dosage of 8 mg (2 mL) twice a day (16 mg (4 mL)/day) after a minimum of 4 weeks. A further increase to 12 mg (3 mL) twice a day (24 mg (6 mL)/day) should be attempted after a minimum of 4 weeks at 8 mg (2 mL) twice a day (16 mg (4 mL)/day).

Dosage increases should be based upon assessment of clinical benefit and tolerability of the previous dose.

Galantamine oral solution should be administered twice a day, preferably with morning and evening meals.

Patients and caregivers should be advised to ensure adequate fluid intake during treatment. If therapy has been interrupted for more than three days, the patient should be restarted at the lowest dosage and the dosage escalated to the current dose.

The abrupt withdrawal of galantamine in those patients who had been receiving dosages in the effective range was not associated with an increased frequency of adverse events in comparison with those continuing to receive the same dosages of that drug. The beneficial effects of galantamine are lost, however, when the drug is discontinued.

2.3 Dosage in Patients with Hepatic Impairment

In patients with moderate hepatic impairment (Child-Pugh score of 7 to 9), the dosage should generally not exceed 16 mg (4 mL)/day. The use of galantamine in patients with severe hepatic impairment (Child-Pugh score of 10 to 15) is not recommended [see *Clinical Pharmacology* (12.3)].

2.4 Dosage in Patients with Renal Impairment

In patients with creatinine clearance of 9 mL to 59 mL/min, the dosage should generally not exceed 16 mg (4 mL)/day. In patients with creatinine clearance less than 9 mL/min, the use of galantamine is not recommended [see *Clinical Pharmacology* (12.3)].

3 DOSAGE FORMS AND STRENGTHS

Galantamine Oral Solution USP, 4 mg/mL is a clear, colorless to pale yellow solution supplied in 100 mL bottles with a calibrated (in milliliters) syringe. The minimum calibrated volume is 0.5 mL, while the maximum calibrated volume is 5 mL.

4 CONTRAINDICATIONS

Galantamine is contraindicated in patients with known hypersensitivity to galantamine hydrobromide or to any excipients used in the formulation.

5 WARNINGS AND PRECAUTIONS

5.1 Serious Skin Reactions

Serious skin reactions (Stevens-Johnson syndrome and acute generalized exanthematous pustulosis) have been reported in patients receiving galantamine. Inform patients and caregivers that the use of galantamine should be discontinued at the first appearance of a skin rash, unless the rash is clearly not drug-related. If signs or symptoms suggest a serious skin reaction, use of this drug should not be resumed and alternative therapy should be considered.

5.2 Anesthesia

Galantamine, as a cholinesterase inhibitor, is likely to exaggerate the neuromuscular blocking effects of succinylcholine-type and similar neuromuscular blocking agents during anesthesia.

5.3 Cardiovascular Conditions

Because of their pharmacological action, cholinesterase inhibitors have vagotonic effects on the sinoatrial and atrioventricular nodes, leading to bradycardia and AV block. Bradycardia and all types of heart block have been reported in patients both with and without known underlying cardiac conduction abnormalities [see *Adverse Reactions*(6.1, 6.2)]. Therefore, all patients should be considered at risk for adverse effects on cardiac conduction.

Patients treated with galantamine up to 24 mg/day using the recommended dosing schedule showed a dose-related increase in risk of syncope (placebo 0.7% [2/286]; 4 mg twice daily 0.4% [3/692]; 8 mg twice daily 1.3% [7/552]; 12 mg twice daily 2.2% [6/273]).

5.4 Gastrointestinal Conditions

Through their primary action, cholinomimetics may be expected to increase gastric acid secretion due to increased cholinergic activity. Therefore, patients should be monitored closely for symptoms of active or occult gastrointestinal bleeding, especially those with an increased risk for developing ulcers, e.g., those with a history of ulcer disease or patients using concurrent nonsteroidal anti-inflammatory drugs (NSAIDs). Clinical studies of galantamine have shown no increase, relative to placebo, in the incidence of

either peptic ulcer disease or gastrointestinal bleeding.

Galantamine, as a predictable consequence of its pharmacological properties, has been shown to produce nausea, vomiting, diarrhea, anorexia, and weight loss. During therapy, the patient's weight should be monitored.

5.5 Genitourinary Conditions

Although this was not observed in clinical trials with galantamine, cholinomimetics may cause bladder outflow obstruction.

5.6 Neurological Conditions

Seizures:

Cholinesterase inhibitors are believed to have some potential to cause generalized convulsions [see *Adverse Reactions* (6.2)]. Seizure activity may also be a manifestation of Alzheimer's disease. Patients with Alzheimer's disease should be monitored closely for seizures while taking galantamine.

5.7 Pulmonary Conditions

Because of its cholinomimetic action, galantamine should be prescribed with care to patients with a history of severe asthma or obstructive pulmonary disease. Respiratory function should be monitored closely for the occurrence of respiratory adverse effects.

5.8 Deaths in Subjects with Mild Cognitive Impairment (MCI)

In two randomized placebo controlled trials of 2 years duration in patients with mild cognitive impairment (MCI), a total of 13 patients on galantamine (n=1026) and 1 patient on placebo (n=1022) died. The deaths were due to various causes which could be expected in an elderly population; about half of the galantamine deaths appeared to result from various vascular causes (myocardial infarction, stroke, and sudden death).

Although the difference in mortality between galantamine- and placebo-treated groups in these two studies was significant, the results are highly discrepant with other studies of galantamine. Specifically, in these two MCI studies, the mortality rate in the placebo-treated patients was markedly lower than the rate in placebo-treated patients in trials of galantamine in Alzheimer's disease or other dementias (0.7 per 1000 person years compared to 22 to 61 per 1000 person years, respectively). Although the mortality rate in the galantamine-treated MCI patients was also lower than that observed in galantamine-treated patients in Alzheimer's disease and other dementia trials (10.2 per 1000 person years compared to 23 to 31 per 1000 person years, respectively), the relative difference was much less. When the Alzheimer's disease and other dementia studies were pooled (n=6000), the mortality rate in the placebo group numerically exceeded that in the galantamine group. Furthermore, in the MCI studies, no patients in the placebo group died after 6 months, a highly unexpected finding in this population.

Individuals with mild cognitive impairment demonstrate isolated memory impairment greater than expected for their age and education, but do not meet current diagnostic criteria for Alzheimer's disease.

6 ADVERSE REACTIONS

Serious adverse reactions are discussed in more detail in the following sections of the labeling:

- Serious Skin Reactions [see Warnings and Precautions (5.1)]
- Cardiovascular Conditions [see Warnings and Precautions (5.3)]
- Gastrointestinal Conditions [see Warnings and Precautions (5.4)]
- Genitourinary Conditions [see Warnings and Precautions (5.5)]
- Neurological Conditions [see Warnings and Precautions (5.6)]
- Pulmonary Conditions [see Warnings and Precautions (5.7)]
- Deaths in Subjects with Mild cognitive impairment (MCI) [see Warnings and Precautions (5.8)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The most common adverse reactions in galantamine-treated patients from double-blind clinical trials ($\geq 5\%$) were nausea, vomiting, diarrhea, dizziness, headache, and decreased appetite.

The most common adverse reactions associated with discontinuation ($\geq 1\%$) in galantamine-treated patients from double-blind clinical trials were nausea (6.2%), vomiting (3.3%), decreased appetite (1.5%), and dizziness (1.3%).

The safety of the extended-release capsule and immediate-release tablet formulations of galantamine was evaluated in 3956 galantamine-treated patients who participated in 8 placebo-controlled clinical studies and 1454 subjects in 5 open-label clinical studies with mild to moderate dementia of the Alzheimer's type. In clinical studies, the safety profile of once-daily treatment with extended-release galantamine was similar in frequency and nature to that seen with tablets. The information presented in this section was derived from pooled double-blind studies and from pooled open-label data.

Commonly-Observed Adverse Reactions in Double-Blind, Placebo-Controlled Clinical Trials:

Table 1 lists the adverse reactions reported in $\geq 1\%$ of galantamine-treated patients in 8 placebo-controlled, double-blind clinical trials.

Table 1: Adverse Reactions Reported by $\geq 1\%$ of Galantamine-Treated Patients in Pooled Placebo-Controlled, Double-Blind Clinical Trials

System/Organ Class Adverse Reaction	Galantamine (n=3956) %	Placebo (n=2546) %
Metabolism and Nutrition Disorders		
Decreased Appetite	7.4	2.1
Psychiatric Disorders		
Depression	3.6	2.3
Nervous System Disorders		
Headache	7.1	5.5

Dizziness	7.5	3.4
Tremor	1.6	0.7
Somnolence	1.5	0.8
Syncope	1.4	0.6
Lethargy	1.3	0.4
Cardiac Disorders		
Bradycardia	1.0	0.3
Gastrointestinal Disorders		
Nausea	20.7	5.5
Vomiting	10.5	2.3
Diarrhea	7.4	4.9
Abdominal Discomfort	2.1	0.7
Abdominal Pain	3.8	2.0
Dyspepsia	1.5	1.0
Musculoskeletal and Connective Tissue Disorders		
Muscle Spasms	1.2	0.5
General Disorders and Administration Site Conditions		
Fatigue	3.5	1.8
Asthenia	2.0	1.5
Malaise	1.1	0.5
Investigations		
Decreased Weight	4.7	1.5
Injury, Poisoning and Procedural Complications		
Fall	3.9	3.0
Laceration	1.1	0.5

The majority of these adverse reactions occurred during the dose-escalation period. In those patients who experienced the most frequent adverse reaction, nausea, the median duration of the nausea was 5 to 7 days.

Other Adverse Reactions Observed in Clinical Trials of Galantamine:

The following adverse reactions occurred in <1% of all galantamine-treated patients (n=3956) in the above double-blind, placebo-controlled clinical trial data sets. In addition, the following also includes all adverse reactions reported at any frequency rate in patients (n=1454) who participated in open-label studies. Adverse reactions listed in Table 1 above were not included below:

Metabolism and Nutrition Disorders: Dehydration

Nervous System Disorders: Dysgeusia, Hypersomnia, Paresthesia

Eye Disorders: Blurred vision

Cardiac Disorders: First degree atrioventricular block, Palpitations, Sinus bradycardia, Supraventricular extrasystoles

Vascular Disorders: Flushing, Hypotension

Gastrointestinal Disorders: Retching

Skin and Subcutaneous Tissue Disorders: Hyperhidrosis

Musculoskeletal and Connective Tissue Disorders: Muscular weakness

Discontinuations Due to Adverse Reactions:

In the 8 placebo-controlled studies of adults, 418 (10.6%) galantamine-treated patients (n=3956) and 56 (2.2%) placebo patients (n=2546) discontinued due to an adverse reaction. Those events with an incidence of $\geq 0.5\%$ in the galantamine-treated patients included nausea (245, 6.2%), vomiting (129, 3.3%), decreased appetite (60, 1.5%), dizziness (50, 1.3%), diarrhea (31, 0.8%), headache (29, 0.7%), and decreased weight (26, 0.7%). The only event with an incidence of $\geq 0.5\%$ in placebo patients was nausea (17, 0.7%).

In the 5 open-label studies, 103 (7.1%) patients (n=1454) discontinued due to an adverse reaction. Those events with an incidence of $\geq 0.5\%$ included nausea (43, 3.0%), vomiting (23, 1.6%), decreased appetite (13, 0.9%), headache (12, 0.8%), decreased weight (9, 0.6%), dizziness (8, 0.6%), and diarrhea (7, 0.5%).

6.2 Postmarketing Experience

The following additional adverse reactions have been identified during post-approval use of galantamine. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency:

Immune System Disorders: Hypersensitivity

Psychiatric Disorders: Hallucinations

Nervous System Disorders: Seizures

Ear and Labyrinth Disorders: Tinnitus

Cardiac Disorders: Complete atrioventricular block

Vascular Disorders: Hypertension

Hepatobiliary Disorders: Hepatitis, Increased hepatic enzyme

Skin and Subcutaneous Tissue Disorders: Stevens-Johnson syndrome, Acute generalized exanthematous pustulosis, Erythema multiforme

7 DRUG INTERACTIONS

7.1 Use with Anticholinergics

Galantamine has the potential to interfere with the activity of anticholinergic medications [see *Clinical Pharmacology* (12.3)].

7.2 Use with Cholinomimetics and Other Cholinesterase Inhibitors

A synergistic effect is expected when cholinesterase inhibitors are given concurrently with succinylcholine, other cholinesterase inhibitors, similar neuromuscular blocking agents or cholinergic agonists such as bethanechol [see *Clinical Pharmacology* (12.3)].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

There are no adequate and well-controlled studies in pregnant women. In studies conducted in animals, administration of galantamine during pregnancy resulted in developmental toxicity (increased incidence of morphological abnormalities and decreased growth in offspring) at doses similar to or greater than those used clinically. Galantamine should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

In rats, administration of galantamine (oral doses of 2 mg, 8 mg, or 16 mg/kg/day), from day 14 (females) or day 60 (males) prior to mating and continuing in females through the period of organogenesis, resulted in an increased incidence of fetal skeletal variations at the two highest doses. The no-effect dose for embryo-fetal developmental toxicity in rats (2 mg/kg/day) is approximately equal to the maximum recommended human dose (MRHD of 24 mg/day) on a body surface area (mg/m^2) basis. When galantamine (oral doses of 4 mg, 12 mg, 28 mg, or 48 mg/kg/day) was administered to pregnant rabbits throughout the period of organogenesis, small increases in fetal visceral malformations and skeletal variations were observed at the highest dose. The no-effect dose for embryo-fetal developmental toxicity in rabbits (28 mg/kg/day) is approximately 20 times the MRHD on a mg/m^2 basis. In a study in which pregnant rats were orally dosed with galantamine (2 mg, 8 mg, or 16 mg/kg/day) from the beginning of organogenesis through day 21 post-partum, pup weights were decreased at birth and during the lactation period at the two highest doses. The no-effect dose for pre- and postnatal developmental toxicity in rats (2 mg/kg/day) is approximately equal to the MRHD on a mg/m^2 basis.

8.3 Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when galantamine is administered to a nursing woman.

8.4 Pediatric Use

The safety and effectiveness in pediatric patients have not been established.

8.5 Geriatric Use

Eight double-blind, placebo-controlled clinical trials and 5 open-label trials in a total of 6519 patients have investigated galantamine in the treatment of mild to moderate dementia of the Alzheimer's type [see *Adverse Reactions* (6.1) and *Clinical Studies* (14)]. The mean age of patients enrolled in these clinical studies was 75 years; 78% of these patients were between 65 and 84 years of age, and 10% of patients were 85 years of age or older.

8.6 Hepatic Impairment

In patients with moderate hepatic impairment, a dosage adjustment is recommended. The use of galantamine in patients with severe hepatic impairment is not recommended [see *Dosage and Administration* (2.3) and *Clinical Pharmacology* (12.3)].

8.7 Renal Impairment

In patients with a creatinine clearance of 9 mL to 59 mL/min, a dosage adjustment is recommended. The use of galantamine in patients with creatinine clearance less than 9 mL/min is not recommended [see *Dosage and Administration* (2.4) and *Clinical Pharmacology* (12.3)].

10 OVERDOSAGE

Because strategies for the management of overdose are continually evolving, it is advisable to contact a poison control center to determine the latest recommendations for the management of an overdose of any drug.

As in any case of overdose, general supportive measures should be utilized. Signs and symptoms of significant overdosing of galantamine are predicted to be similar to those of overdosing of other cholinomimetics. These effects generally involve the central nervous system, the parasympathetic nervous system, and the neuromuscular junction. In addition to muscle weakness or fasciculations, some or all of the following signs of cholinergic crisis may develop: severe nausea, vomiting, gastrointestinal cramping, salivation, lacrimation, urination, defecation, sweating, bradycardia, hypotension, respiratory depression, collapse and convulsions. Increasing muscle weakness is a possibility and may result in death if respiratory muscles are involved.


Tertiary anticholinergics such as atropine may be used as an antidote for galantamine overdosage. Intravenous atropine sulfate titrated to effect is recommended at an initial dose of 0.5 mg to 1.0 mg i.v. with subsequent doses based upon clinical response. Atypical responses in blood pressure and heart rate have been reported with other cholinomimetics when co-administered with quaternary anticholinergics. It is not known whether galantamine and/or its metabolites can be removed by dialysis (hemodialysis, peritoneal dialysis, or hemofiltration). Dose-related signs of toxicity in animals included hypoactivity, tremors, clonic convulsions, salivation, lacrimation, chromodacryorrhea, mucoid feces, and dyspnea.

In one postmarketing report, one patient who had been taking 4 mg of galantamine daily for a week inadvertently ingested eight 4 mg tablets (32 mg total) on a single day. Subsequently, she developed bradycardia, QT prolongation, ventricular tachycardia and torsades de pointes accompanied by a brief loss of consciousness for which she required hospital treatment. Two additional cases of accidental ingestion of 32 mg (nausea, vomiting, and dry mouth; nausea, vomiting, and substernal chest pain) and one of 40 mg (vomiting), resulted in brief hospitalizations for observation with full recovery. One patient, who was prescribed 24 mg/day and had a history of hallucinations over the previous two years, mistakenly received 24 mg twice daily for 34 days and developed hallucinations requiring hospitalization. Another patient, who was prescribed 16 mg/day of oral solution, inadvertently ingested 160 mg (40 mL) and experienced sweating, vomiting, bradycardia, and near-syncope one hour later, which necessitated hospital treatment. His symptoms resolved within 24 hours.

11 DESCRIPTION

Galantamine Oral Solution, USP contains galantamine hydrobromide, USP, a reversible, competitive acetylcholinesterase inhibitor, as the hydrobromide salt. Galantamine

hydrobromide is known chemically as (4a*S*,6*R*,8a*S*)-4a,5,9,10,11,12-hexahydro-3-methoxy-11-methyl-6*H*-benzofuro[3a,3,2-*ef*][2]benzazepin-6-ol hydrobromide. It has an empirical formula of C₁₇H₂₁NO₃ • HBr and a molecular weight of 368.27. Galantamine hydrobromide, USP is a white to off-white powder and is sparingly soluble in water. The structural formula is:



Galantamine Oral Solution, USP contains 4 mg (as 5.13 mg galantamine hydrobromide, USP) per mL. The inactive ingredients are methylparaben, propylparaben, purified water and saccharin sodium. Galantamine may contain hydrochloric acid and/or sodium hydroxide to adjust pH.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Although the etiology of cognitive impairment in Alzheimer's disease (AD) is not fully understood, it has been reported that acetylcholine-producing neurons degenerate in the brains of patients with Alzheimer's disease. The degree of this cholinergic loss has been correlated with degree of cognitive impairment and density of amyloid plaques (a neuropathological hallmark of Alzheimer's disease).

Galantamine, a tertiary alkaloid, is a competitive and reversible inhibitor of acetylcholinesterase. While the precise mechanism of galantamine's action is unknown, it is postulated to exert its therapeutic effect by enhancing cholinergic function. This is accomplished by increasing the concentration of acetylcholine through reversible inhibition of its hydrolysis by cholinesterase. If this mechanism is correct, galantamine's effect may lessen as the disease process advances and fewer cholinergic neurons remain functionally intact. There is no evidence that galantamine alters the course of the underlying dementing process.

12.3 Pharmacokinetics

The pharmacokinetics of galantamine are linear over a dose range of 8 mg to 32 mg/day.

Absorption and Distribution: Galantamine is absorbed with time to peak concentration of about 1 hour. The absolute bioavailability of galantamine is about 90%. The bioavailability of the tablet formulation was the same as the bioavailability of the oral solution formulation. Food did not affect the AUC of galantamine, but C_{max} was decreased by 25% and T_{max} was delayed by 1.5 hours, when galantamine was administered with food. The mean volume of distribution of galantamine is 175 L.

The plasma protein binding of galantamine is 18% at therapeutically relevant concentrations. In whole blood, galantamine is mainly distributed to blood cells (52.7%). The blood to plasma concentration ratio of galantamine is 1.2.

Metabolism and Elimination: Galantamine is metabolized by hepatic cytochrome P450 enzymes, glucuronidated, and excreted unchanged in the urine. *In vitro* studies indicate that cytochrome CYP2D6 and CYP3A4 were the major cytochrome P450 isoenzymes involved in the metabolism of galantamine, and inhibitors of both pathways increase oral bioavailability of galantamine modestly. O-demethylation, mediated by CYP2D6 was greater in extensive metabolizers of CYP2D6 than in poor metabolizers. In plasma from both poor and extensive metabolizers, however, unchanged galantamine and its glucuronide accounted for most of the sample radioactivity.

In studies of oral ^3H -galantamine, unchanged galantamine and its glucuronide, accounted for most plasma radioactivity in poor and extensive CYP2D6 metabolizers. Up to 8 hours post-dose, unchanged galantamine accounted for 39% to 77% of the total radioactivity in the plasma, and galantamine glucuronide for 14% to 24%. By 7 days, 93% to 99% of the radioactivity had been recovered, with about 95% in urine and about 5% in the feces. Total urinary recovery of unchanged galantamine accounted for, on average, 32% of the dose and that of galantamine glucuronide for another 12% on average.

After i.v. or oral administration, about 20% of the dose was excreted as unchanged galantamine in the urine in 24 hours, representing a renal clearance of about 65 mL/min, about 20% to 25% of the total plasma clearance of about 300 mL/min. Galantamine has a terminal half-life of about 7 hours.

Specific Populations: Elderly: Data from clinical trials in patients with Alzheimer's disease indicate that galantamine concentrations are 30% to 40% higher in those patients than in healthy young subjects.

Gender and Race: A population pharmacokinetic analysis (on 539 men and 550 women) indicates that galantamine clearance is about 20% lower in women than in men (which is explained by a lower body weight in women) and that race (n=1029 White, 24 Black, 13 Asian and 23 other) did not affect the clearance of galantamine.

Hepatic Impairment: Following a single 4 mg dose of galantamine tablets, the pharmacokinetics of galantamine in subjects with mild hepatic impairment (n=8; Child-Pugh score of 5 to 6) were similar to the pharmacokinetics of galantamine in healthy subjects. In patients with moderate hepatic impairment (n=8; Child Pugh score of 7 to 9), galantamine clearance was decreased by about 25% compared to galantamine clearance in normal volunteers. Exposure to galantamine would be expected to increase further with increasing degree of hepatic impairment [see *Dosage and Administration* (2.3) and *Use in Specific Populations* (8.6)].

Renal Impairment: Following a single 8 mg dose of galantamine tablets, AUC increased by 37% and 67% in patients with moderate and severe renal impairment, respectively, compared with normal volunteers [see *Dosage and Administration* (2.4) and *Use in Specific Populations* (8.7)].

CYP2D6 Poor Metabolizers: Approximately 7% of the normal population has a genetic variation that leads to reduced levels of activity of CYP2D6 isozyme. Such individuals have been referred to as poor metabolizers. After a single oral dose of 4 mg or 8 mg

galantamine, CYP2D6 poor metabolizers demonstrated a similar C_{max} and about 35% AUC_{∞} increase of unchanged galantamine compared to extensive metabolizers.

A total of 356 patients with Alzheimer's disease enrolled in two Phase 3 studies were genotyped with respect to CYP2D6 (n=210 hetero-extensive metabolizers, 126 homo-extensive metabolizers, and 20 poor metabolizers). Population pharmacokinetic analysis indicated that there was a 25% decrease in median clearance in poor metabolizers compared to extensive metabolizers. Dosage adjustment is not necessary in patients identified as poor metabolizers as the dose of drug is individually titrated to tolerability.

Drug-Drug Interactions: Multiple metabolic pathways and renal excretion are involved in the elimination of galantamine so no single pathway appears predominant. Based on *in vitro* studies, CYP2D6 and CYP3A4 were the major enzymes involved in the metabolism of galantamine. CYP2D6 was involved in the formation of O-desmethyl-galantamine, whereas CYP3A4 mediated the formation of galantamine-N-oxide. Galantamine is also glucuronidated and excreted unchanged in urine.

Effect of Other Drugs on Galantamine:

- CYP3A4 Inhibitors:
Ketoconazole: Ketoconazole, a strong inhibitor of CYP3A4 and an inhibitor of CYP2D6, when administered at a dose of 200 mg two times a day for 4 days, increased the AUC of galantamine by 30%.
Erythromycin: Erythromycin, a moderate inhibitor of CYP3A4, when administered at a dose of 500 mg four times a day for 4 days, affected the AUC of galantamine minimally (10% increase).
- CYP2D6 Inhibitors: A population pharmacokinetics analysis on a database of 852 patients with Alzheimer's disease showed that the clearance of galantamine was reduced about 25% to 33% by the concurrent administration of amitriptyline (n=17), fluoxetine (n=48), fluvoxamine (n=14), and quinidine (n=7), all of which are known inhibitors of CYP2D6.
Paroxetine: Paroxetine, a strong inhibitor of CYP2D6, when administered at a dose of 20 mg/day for 16 days, increased the oral bioavailability of galantamine by about 40%.
- H2 Antagonists: Galantamine was administered as a single dose of 4 mg on Day 2 of a 3-day treatment with either cimetidine (800 mg daily) or ranitidine (300 mg daily). Cimetidine increased the bioavailability of galantamine by approximately 16%. Ranitidine had no effect on the pharmacokinetics of galantamine.
- Memantine: Memantine, an N-methyl-D-aspartate receptor antagonist, when administered at a dose of 10 mg two times a day, had no effect on the pharmacokinetics of galantamine (16 mg/day) at steady state.

Effect of Galantamine on Other Drugs:

- In Vitro Studies: *In vitro* studies show that galantamine did not inhibit the metabolic pathways catalyzed by CYP1A2, CYP2A6, CYP3A4, CYP4A, CYP2C, CYP2D6 or CYP2E1. This indicates that the inhibitory potential of galantamine towards the major forms of cytochrome P450 is very low.
- In Vivo Studies: *Warfarin:* Multiple doses of galantamine at 24 mg/day had no effect on the pharmacokinetics of R- and S-warfarin (administered in a single dose of 25 mg) or on the increased prothrombin time induced by warfarin. The protein binding of warfarin was unaffected by galantamine.

Digoxin: Multiple doses of galantamine at 24 mg/day had no effect on the steady-state pharmacokinetics of digoxin (at a dose of 0.375 mg once daily) when those two drugs were co-administered. In that study, however, one healthy subject was hospitalized on account of 2nd and 3rd degree heart block and bradycardia.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis: In a 24-month oral carcinogenicity study in rats, an increase in endometrial adenocarcinomas was observed at 10 mg/kg/day (4 times the MRHD of 24 mg/day on a mg/m² basis or 6 times on a plasma exposure [AUC] basis) and 30 mg/kg/day (12 times MRHD on a mg/m² basis or 19 times on an AUC basis). No increase in neoplastic changes was observed in females at 2.5 mg/kg/day (equivalent to the MRHD on a mg/m² basis or 2 times on an AUC basis) or in males up to the highest dose tested of 30 mg/kg/day (12 times the MRHD on a mg/m² and AUC basis).

Galantamine was not carcinogenic in a 6-month carcinogenicity study in transgenic (P 53-deficient) mice at oral doses up to 20 mg/kg/day, or in a 24-month carcinogenicity study in mice at oral doses up to 10 mg/kg/day (equivalent to the MRHD on a plasma AUC basis).

Mutagenesis: Galantamine was negative in a battery of *in vitro* (bacterial reverse mutation, mouse lymphoma *tk*, and chromosomal aberration in mammalian cells) and *in vivo* (mouse micronucleus) genotoxicity assays.

Impairment of Fertility: No impairment of fertility was seen in rats given up to 16 mg/kg/day (7 times the MRHD on a mg/m² basis) for 14 days prior to mating in females and for 60 days prior to mating in males.

14 CLINICAL STUDIES

The effectiveness of galantamine as a treatment for Alzheimer's disease is demonstrated by the results of 5 randomized, double-blind, placebo-controlled clinical investigations in patients with probable Alzheimer's disease, 4 with the immediate-release tablet and 1 with the extended-release capsule [diagnosed by NINCDS-ADRDA criteria, with Mini-Mental State Examination scores that were ≥ 10 and ≤ 24]. Doses studied with the tablet formulation were 8 mg to 32 mg/day given as twice daily doses. In 3 of the 4 studies with the tablet, patients were started on a low dose of 8 mg, then titrated weekly by 8 mg/day to 24 mg or 32 mg as assigned. In the fourth study (USA 4-week Dose Escalation Fixed-Dose Study) dose escalation of 8 mg/day occurred over 4-week intervals. The mean age of patients participating in these 4 galantamine trials was 75 years with a range of 41 to 100. Approximately 62% of patients were women and 38% were men. The racial distribution was White 94%, Black 3% and other races 3%. Two other studies examined a three times daily dosing regimen; these also showed or suggested benefit but did not suggest an advantage over twice daily dosing.

14.1 Study Outcome Measures

In each study, the primary effectiveness of galantamine was evaluated using a dual outcome assessment strategy as measured by the Alzheimer's Disease Assessment

Scale (ADAS-cog) and the Clinician's Interview Based Impression of Change that required the use of caregiver information (CIBIC-plus).

The ability of galantamine to improve cognitive performance was assessed with the cognitive sub-scale of the Alzheimer's Disease Assessment Scale (ADAS-cog), a multi-item instrument that has been extensively validated in longitudinal cohorts of Alzheimer's disease patients. The ADAS-cog examines selected aspects of cognitive performance including elements of memory, orientation, attention, reasoning, language and praxis. The ADAS-cog scoring range is from 0 to 70, with higher scores indicating greater cognitive impairment. Elderly normal adults may score as low as 0 or 1, but it is not unusual for non-demented adults to score slightly higher.

The patients recruited as participants in each study using the tablet formulation had mean scores on ADAS-cog of approximately 27 units, with a range from 5 to 69. Experience gained in longitudinal studies of ambulatory patients with mild to moderate Alzheimer's disease suggests that they gain 6 to 12 units a year on the ADAS-cog. Lesser degrees of change, however, are seen in patients with very mild or very advanced disease because the ADAS-cog is not uniformly sensitive to change over the course of the disease. The annualized rate of decline in the placebo patients participating in galantamine trials was approximately 4.5 units per year.

The ability of galantamine to produce an overall clinical effect was assessed using a Clinician's Interview Based Impression of Change that required the use of caregiver information, the CIBIC-plus. The CIBIC-plus is not a single instrument and is not a standardized instrument like the ADAS-cog. Clinical trials for investigational drugs have used a variety of CIBIC formats, each different in terms of depth and structure. As such, results from a CIBIC-plus reflect clinical experience from the trial or trials in which it was used and cannot be compared directly with the results of CIBIC-plus evaluations from other clinical trials. The CIBIC-plus used in the trials was a semi-structured instrument based on a comprehensive evaluation at baseline and subsequent time-points of 4 major areas of patient function: general, cognitive, behavioral and activities of daily living. It represents the assessment of a skilled clinician based on his/her observation at an interview with the patient, in combination with information supplied by a caregiver familiar with the behavior of the patient over the interval rated. The CIBIC-plus is scored as a seven point categorical rating, ranging from a score of 1, indicating "markedly improved," to a score of 4, indicating "no change" to a score of 7, indicating "marked worsening." The CIBIC-plus has not been systematically compared directly to assessments not using information from caregivers (CIBIC) or other global methods.

14.2 Immediate-Release Tablets

U.S. Twenty-One Week Fixed-Dose Study: In a study of 21 weeks duration, 978 patients were randomized to doses of 8 mg, 16 mg, or 24 mg of galantamine per day, or to placebo, each given in 2 divided doses. Treatment was initiated at 8 mg/day for all patients randomized to galantamine, and increased by 8 mg/day every 4 weeks. Therefore, the maximum titration phase was 8 weeks and the minimum maintenance phase was 13 weeks (in patients randomized to 24 mg/day of galantamine).

Effects on the ADAS-cog: Figure 1 illustrates the time course for the change from baseline in ADAS-cog scores for all four dose groups over the 21 weeks of the study. At 21 weeks of treatment, the mean differences in the ADAS-cog change scores for the galantamine-treated patients compared to the patients on placebo were 1.7, 3.3, and 3.6

units for the 8 mg, 16 mg and 24 mg/day treatments, respectively. The 16 mg/day and 24 mg/day treatments were statistically significantly superior to placebo and to the 8 mg/day treatment. There was no statistically significant difference between the 16 mg/day and 24 mg/day dose groups.

Figure 1: Time-Course of the Change From Baseline in ADAS-cog Score for Patients Completing 21 Weeks (5 Months) of Treatment




Figure 1: Time-Course of the Change From Baseline in ADAS-cog Score for Patients Completing 21 Weeks (5 Months) of Treatment

Figure 2 illustrates the cumulative percentages of patients from each of the four treatment groups who had attained at least the measure of improvement in ADAS-cog score shown on the X-axis. Three change scores (10-point, 7-point and 4-point reductions) and no change in score from baseline have been identified for illustrative purposes, and the percent of patients in each group achieving that result is shown in the inset table.

The curves demonstrate that both patients assigned to galantamine and placebo have a wide range of responses, but that the galantamine groups are more likely to show the greater improvements.

Figure 2: Cumulative Percentage of Patients Completing 21 Weeks of Double-Blind Treatment With Specified Changes From Baseline in ADAS-cog Scores. The Percentages of Randomized Patients Who Completed the Study Were: Placebo 84%, 8 mg/day 77%, 16 mg/day 78% and 24 mg/day 78%.




Figure 2: Cumulative Percentage of Patients Completing 21 Weeks of Double-Blind Treatment With Specified Changes From Baseline in ADAS-cog Scores. The Percentages of Randomized Patients Who Completed the Study Were: Placebo 84%, 8 mg/day 77%, 16 mg/day 78% and 24 mg/day 78%.

Treatment	Change in ADAS-cog			
	-10	-7	-4	-0
Placebo	3.6%	7.6%	19.6%	41.8%
8 mg/day	5.9%	13.9%	25.7%	46.5%
16 mg/day	7.2%	15.9%	35.6%	65.4%
24 mg/day	10.4%	22.3%	37.0%	64.9%

Effects on the CIBIC-plus: Figure 3 is a histogram of the percentage distribution of CIBIC-plus scores attained by patients assigned to each of the four treatment groups who completed 21 weeks of treatment. The galantamine-placebo differences for these groups of patients in mean rating were 0.15, 0.41 and 0.44 units for the 8 mg, 16 mg and 24 mg/day treatments, respectively. The 16 mg/day and 24 mg/day treatments were statistically significantly superior to placebo. The differences vs. the 8 mg/day treatment for the 16 mg and 24 mg/day treatments were 0.26 and 0.29, respectively. There were no statistically significant differences between the 16 mg/day and 24 mg/day dose groups.

Figure 3: Distribution of CIBIC-plus Ratings at Week 21




Figure 3: Distribution of CIBIC-plus Ratings at Week 21

U.S. Twenty-Six Week Fixed-Dose Study: In a study of 26 weeks duration, 636 patients were randomized to either a dose of 24 mg or 32 mg of galantamine per day, or to placebo, each given in two divided doses. The 26-week study was divided into a 3-week dose titration phase and a 23-week maintenance phase.

Effects on the ADAS-cog: Figure 4 illustrates the time course for the change from baseline in ADAS-cog scores for all three dose groups over the 26 weeks of the study. At 26 weeks of treatment, the mean differences in the ADAS-cog change scores for the galantamine-treated patients compared to the patients on placebo were 3.9 and 3.8 units for the 24 mg/day and 32 mg/day treatments, respectively. Both treatments were statistically significantly superior to placebo, but were not significantly different from each other.

Figure 4: Time-Course of the Change From Baseline in ADAS-cog Score for Patients Completing 26 Weeks of Treatment




Figure 4: Time-Course of the Change From Baseline in ADAS-cog Score for Patients Completing 26 Weeks of Treatment

Figure 5 illustrates the cumulative percentages of patients from each of the three treatment groups who had attained at least the measure of improvement in ADAS-cog score shown on the X-axis. Three change scores (10-point, 7-point and 4-point reductions) and no change in score from baseline have been identified for illustrative purposes, and the percent of patients in each group achieving that result is shown in the inset table.

The curves demonstrate that both patients assigned to galantamine and placebo have a wide range of responses, but that the galantamine groups are more likely to show the greater improvements. A curve for an effective treatment would be shifted to the left of the curve for placebo, while an ineffective or deleterious treatment would be superimposed upon, or shifted to the right of the curve for placebo, respectively.

Treatment	Change in ADAS-cog			
	-10	-7	-4	-0
Placebo	2.1%	5.7%	16.6%	43.9%
24 mg/day	7.6%	18.3%	33.6%	64.1%
32 mg/day	11.1%	19.7%	33.3%	58.1%

Figure 5: Cumulative Percentage of Patients Completing 26 Weeks of Double-Blind Treatment With Specified Changes From Baseline in ADAS-cog Scores. The Percentages of Randomized Patients Who Completed the Study Were: Placebo 81%, 24 mg/day 68%, and 32 mg/day 58%.




Figure 5: Cumulative Percentage of Patients Completing 26 Weeks of Double-Blind Treatment With Specified Changes From Baseline in ADAS-cog Scores. The Percentages of Randomized Patients Who Completed the Study Were: Placebo 81%, 24 mg/day 68%, and 32 mg/day 58%.

Effects on the CIBIC-plus: Figure 6 is a histogram of the percentage distribution of CIBIC-plus scores attained by patients assigned to each of the three treatment groups who completed 26 weeks of treatment. The mean galantamine-placebo differences for these groups of patients in the mean rating were 0.28 and 0.29 units for 24 mg and 32 mg/day of galantamine, respectively. The mean ratings for both groups were statistically significantly superior to placebo, but were not significantly different from each other.

Figure 6: Distribution of CIBIC-plus Ratings at Week 26




Figure 6: Distribution of CIBIC-plus Ratings at Week 26

International Twenty-Six Week Fixed-Dose Study: In a study of 26 weeks duration identical in design to the USA 26-Week Fixed-Dose Study, 653 patients were randomized to either a dose of 24 mg or 32 mg of galantamine per day, or to placebo, each given in two divided doses. The 26-week study was divided into a 3-week dose titration phase and a 23-week maintenance phase.

Effects on the ADAS-cog: Figure 7 illustrates the time course for the change from baseline in ADAS-cog scores for all three dose groups over the 26 weeks of the study. At 26 weeks of treatment, the mean differences in the ADAS-cog change scores for the galantamine-treated patients compared to the patients on placebo were 3.1 and 4.1 units for the 24 mg/day and 32 mg/day treatments, respectively. Both treatments were statistically significantly superior to placebo, but were not significantly different from each other.

Figure 7: Time-Course of the Change From Baseline in ADAS-cog Score for Patients Completing 26 Weeks of Treatment




Figure 7: Time-Course of the Change From Baseline in ADAS-cog Score for Patients Completing 26 Weeks of Treatment

Figure 8 illustrates the cumulative percentages of patients from each of the three treatment groups who had attained at least the measure of improvement in ADAS-cog score shown on the X-axis. Three change scores (10-point, 7-point and 4-point reductions) and no change in score from baseline have been identified for illustrative purposes, and the percent of patients in each group achieving that result is shown in the inset table.

The curves demonstrate that both patients assigned to galantamine and placebo have a wide range of responses, but that the galantamine groups are more likely to show the greater improvements.

Figure 8: Cumulative Percentage of Patients Completing 26 Weeks of Double-Blind Treatment With Specified Changes From Baseline in ADAS-cog Scores. The Percentages of Randomized Patients Who Completed the Study Were: Placebo 87%, 24 mg/day 80%, and 32 mg/day 75%.




Figure 8: Cumulative Percentage of Patients Completing 26 Weeks of Double-Blind Treatment With Specified Changes From Baseline in ADAS-cog Scores. The Percentages of Randomized Patients Who Completed the Study Were: Placebo 87%, 24 mg/day 80%, and 32 mg/day 75%.

Treatment	Change in ADAS-cog			
	-10	-7	-4	-0
Placebo	1.2%	5.8%	15.2%	39.8%
24 mg/day	4.5%	15.4%	30.8%	65.4%
32 mg/day	7.9%	19.7%	34.9%	63.8%

Effects on the CIBIC-plus: Figure 9 is a histogram of the percentage distribution of CIBIC-plus scores attained by patients assigned to each of the three treatment groups who completed 26 weeks of treatment. The mean galantamine-placebo differences for these groups of patients in the mean rating of change from baseline were 0.34 and 0.47 for 24 mg and 32 mg/day of galantamine respectively. The mean ratings for the galantamine groups were statistically significantly superior to placebo, but were not significantly different from each other.

Figure 9: Distribution of CIBIC-plus Rating at Week 26




Figure 9: Distribution of CIBIC-plus Rating at Week 26

International Thirteen-Week Flexible-Dose Study: In a study of 13 weeks duration, 386 patients were randomized to either a flexible dose of 24 mg to 32 mg/day of galantamine or to placebo, each given in two divided doses. The 13-week study was divided into a 3-week dose titration phase and a 10-week maintenance phase. The patients in the active treatment arm of the study were maintained at either 24 mg/day or 32 mg/day at the discretion of the investigator.

Effects on the ADAS-cog: Figure 10 illustrates the time course for the change from baseline in ADAS-cog scores for both dose groups over the 13 weeks of the study. At 13 weeks of treatment, the mean difference in the ADAS-cog change scores for the treated patients compared to the patients on placebo was 1.9. Galantamine at a dose of 24 mg to 32 mg/day was statistically significantly superior to placebo.

Figure 10: Time-Course of the Change From Baseline in ADAS-cog Score for Patients Completing 13 Weeks of Treatment




Figure 10: Time-Course of the Change From Baseline in ADAS-cog Score for Patients Completing 13 Weeks of Treatment

Figure 11 illustrates the cumulative percentages of patients from each of the two treatment groups who had attained at least the measure of improvement in ADAS-cog score shown on the X-axis. Three change scores (10-point, 7-point and 4-point reductions) and no change in score from baseline have been identified for illustrative purposes, and the percent of patients in each group achieving that result is shown in the inset table.

The curves demonstrate that both patients assigned to galantamine and placebo have a wide range of responses, but that the galantamine group is more likely to show the greater improvement.

Figure 11: Cumulative Percentage of Patients Completing 13 Weeks of Double-Blind Treatment With Specified Changes from Baseline in ADAS-cog Scores. The Percentages of Randomized Patients Who Completed the Study Were: Placebo 90%, 24 mg to 32 mg/day 67%.




Figure 11: Cumulative Percentage of Patients Completing 13 Weeks of Double-Blind Treatment With Specified Changes from Baseline in ADAS-cog Scores. The Percentages of Randomized Patients Who Completed the Study Were: Placebo 90%, 24 mg to 32 mg/day 67%.

Treatment	Change in ADAS-cog			
	-10	-7	-4	-0
Placebo	1.9%	5.6%	19.4%	50.0%
24 mg or 32 mg/day	7.1%	18.8%	32.9%	65.3%

Effects on the CIBIC-plus: Figure 12 is a histogram of the percentage distribution of CIBIC-plus scores attained by patients assigned to each of the two treatment groups who completed 13 weeks of treatment. The mean galantamine-placebo differences for the group of patients in the mean rating of change from baseline were 0.37 units. The mean rating for the 24 mg to 32 mg/day group was statistically significantly superior to placebo.

Figure 12: Distribution of CIBIC-plus Ratings at Week 13




Figure 12: Distribution of CIBIC-plus Ratings at Week 13

Age, Gender and Race: Patient's age, gender, or race did not predict clinical outcome of treatment.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Galantamine Oral Solution, USP

4 mg/mL oral solution is supplied in a 100 mL bottle as a clear, colorless to pale yellow solution with a calibrated (in milliliters) syringe. The minimum calibrated volume is 0.5 mL, while the maximum calibrated volume is 5 mL.

NDC 0054-0137-49: Bottle of 100 mL

16.2 Storage and Handling

Galantamine Oral Solution, USP should be stored at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.] **DO NOT FREEZE.**

Keep this and all drugs out of the reach of children.

17 PATIENT COUNSELING INFORMATION

Serious Skin Reactions:

Advise patients and caregivers to discontinue galantamine and seek immediate medical attention at the first appearance of skin rash [see *Warnings and Precautions (5.1)*].

General Dosing Guidance:

Instruct caregivers about the recommended dosage and administration of galantamine. Galantamine oral solution should be administered twice per day, preferably with the morning and evening meals. Dose escalation (dose increases) should follow a minimum of four weeks at prior dose. If therapy has been interrupted for more than three days, the patient should be restarted with the lowest dose and then re-titrated to an appropriate dosage [see *Dosage and Administration (2)*].

Advise patients and caregivers to ensure adequate fluid intake during treatment [see *Dosage and Administration (2)*].

Advise patients and caregivers that the most frequent adverse events associated with use of the drug can be minimized by following the recommended dosage and administration.

Oral Solution Instruction Sheet:

Instruct caregivers in the correct procedure for administering galantamine oral solution. In addition, inform them of the existence of an Instruction Sheet (included with the product) describing how the solution is to be administered. Urge caregivers to read this sheet prior to administering galantamine oral solution, and to direct questions about the administration of the solution to either their physician or pharmacist.

Distributed by:

Hikma Pharmaceuticals USA Inc.

Berkeley Heights, NJ 07922

C50000575/03

Revised October 2023

PATIENT INFORMATION INSTRUCTION SHEET

**Galantamine
Oral Solution, USP
(ga lan' ta meen)**

Rx only

USING YOUR GALANTAMINE DISPENSING-SYRINGE AND BOTTLE

Follow the directions below to use your galantamine dispensing-syringe and bottle, unless your doctor gave you different directions.

IMPORTANT: Read these instructions before using Galantamine Oral Solution.

To open the bottle and use the syringe (plastic tube):

The bottle comes with a child-proof cap. Here is how to open it:

1. Push the plastic cap on the bottle down while turning the cap counter-clockwise (to the left) (Figure 1). Remove the unscrewed cap.




Figure 1

2. Pull the syringe out of the carton. The syringe is a tube that you use to measure your dose of galantamine. Place the syringe fully into the bottle of galantamine (Figure 2).




Figure 2

3. While holding the bottom ring of the syringe, invert bottle and pull the syringe plunger up to the level that equals the dose prescribed by your doctor. Use the markings on the syringe to guide you. This will draw the medicine into the syringe (Figures 3a-3b).




Figure 3a




Figure 3b

4. Be careful not to push the plunger in during this step. Hold the bottom ring of the syringe. Turn bottle upright and remove the entire syringe from the bottle (Figures 4a-4b).



Figure 4a



Figure 4b

5. Empty all the medicine in the syringe into 3 to 4 ounces (100 mL) of any non-alcoholic drink. To do this, push the plunger all the way in (Figures 5a-5b).



Figure 5a




Figure 5b

6. Stir the drink well (Figure 6). Drink all of the mixture right away.



Figure 6

7. Replace the plastic cap on the bottle by turning it clock-wise (to the right) (Figure 7).



Figure 7

8. Rinse the empty syringe by inserting the open end of the syringe into a glass of water, pulling the plunger out, and pushing the plunger in to remove the water (Figures 8a-8b).




Figure 8a




Figure 8b

For more information about galantamine, see the leaflet that came with the package.

Storage

Galantamine Oral Solution, USP should be stored at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.] DO NOT FREEZE.

Keep out of reach of children.

Galantamine Oral Solution USP, 4 mg/mL

Each 1 mL contains the equivalent of 4 mg galantamine in an aqueous solution.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

This Patient Information has been approved by the U.S. Food and Drug Administration.


Distributed by:

Hikma Pharmaceuticals USA Inc.

Berkeley Heights, NJ 07922

C50000575/03**Revised October 2023****PRINCIPAL DISPLAY PANEL - 100 mL Bottle Label****NDC 0054-0137-49** 100 mL**Galantamine**
Oral Solution, USP
4 mg/mL

Rx only

**GALANTAMINE HYDROBROMIDE**

galantamine hydrobromide solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0054-0137
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GALANTAMINE HYDROBROMIDE (UNII: MJ4PTD2VWW) (GALANTAMINE - UNII:0D3Q044KCA)	GALANTAMINE	4 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	

SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
WATER (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0054-0137-49	100 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package	08/14/2009	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA078185	08/14/2009	

Labeler - Hikma Pharmaceuticals USA Inc. (080189610)

Establishment

Name	Address	ID/FEI	Business Operations
West-Ward Columbus Inc.		058839929	MANUFACTURE(0054-0137)

Revised: 10/2023

Hikma Pharmaceuticals USA Inc.