

**KIRKLAND SIGNATURE ACID CONTROLLER- famotidine tablet, film coated**  
**Costco Wholesale Company**

-----  
**Costco Wholesale Corp. Acid Controller Drug Facts**

**Active ingredient (in each tablet)**

Famotidine 20 mg

**Purpose**

Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

**Other information**

- read the directions and warnings before use
- store at 20°-25°C (68°-77°F)
- protect from moisture

**Inactive ingredients**

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

**Questions or comments?**

**1-800-719-9260**

**Principal Display Panel**

KIRKLAND Signature

COMPARE TO MAXIMUM STRENGTH PEPCID® AC active ingredient

ACID CONTROLLER

Famotidine Tablets USP, 20 mg

Acid Reducer

- Maximum Strength

- Just One Tablet Prevents & Relieves Heartburn Due to Acid Indigestion

Actual Size

2 x 125 COUNT BOTTLES TOTAL

250 TABLETS

**KIRKLAND**  
*Signature*

**COMPARE TO  
MAXIMUM STRENGTH  
PEPCID® AC**  
active ingredient\*

NDC 63951-194-85  
ITM / ART. 1086592

# ACID CONTROLLER

Famotidine Tablets USP, 20 mg

**Acid Reducer**

**Maximum Strength**

**Just One Tablet Prevents  
& Relieves Heartburn Due  
to Acid Indigestion**



Actual Size

**2 x 125 COUNT  
BOTTLES TOTAL  
250 TABLETS**

### Drug Facts

#### Active ingredient (in each tablet)

Famotidine 20 mg

For use as  
Acid reducer

#### Use

- Relieves heartburn associated with acid indigestion and sour stomach
- Prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

#### Warnings

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

#### Do not use

- If you have trouble or pain swallowing food, vomiting with blood, or black stools. There may be signs of a serious condition. See your doctor
- With other acid reducers

Ask a doctor before use if you have

- Heartburn over 3 months. This may be a sign of a more serious condition
- Heartburn with lightheadedness, sweating, or dizziness
- Heart pain or chest pain with shortness of breath, sweating, pain spreading to arms, neck or jawline, or lightheadedness
- Frequency of heartburn
- Frequency of heartburn, particularly with sour or unexplained weight loss
- Nausea or vomiting
- Stomach pain
- Kidney disease

Ask a doctor or pharmacist before use. If you are taking a prescription drug, Acid Reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- Your heartburn continues or worsens
- You need to take this product for more than 14 days

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away. (1-800-322-1222)

#### Directions

- Adults and children 12 years and over:  
Take one or two tablets, swallow 1 tablet with a glass of water. Do not chew.
- To prevent symptoms, swallow 1 tablet with a glass of water at the same time for

**JUST ONE TABLET** prevents and relieves heartburn due to acid indigestion brought on by eating or drinking certain foods and beverages.

**DO NOT USE IF PRINTED FOIL LINER CAP IS BROKEN OR MISSING**

#### Tips For Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid lying in bed
- If overweight, lose weight
- Quit smoking



## KIRKLAND SIGNATURE ACID CONTROLLER

famotidine tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63981-194
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>FAMOTIDINE</b> (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	L194

**Contains****Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63981-194-96	2 in 1 PACKAGE	05/24/2007	08/01/2018
1		85 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:63981-194-85	2 in 1 CARTON	11/08/2016	
2		125 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:63981-194-01	85 in 1 BOTTLE; Type 0: Not a Combination Product	05/24/2007	08/01/2014
4	NDC:63981-194-05	125 in 1 BOTTLE; Type 0: Not a Combination Product	01/25/2017	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077351	05/24/2007	

**Labeler** - Costco Wholesale Company (103391843)

Revised: 2/2025

Costco Wholesale Company