

IMAGE SKINCARE PREVENTION PLUS DAILY HYDRATING MOISTURIZER- zinc oxide cream Natural Technology, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Zinc Oxide 17.1%. Purpose: Sunscreen





170g. NDC: 79458-001-00

Purpose: Sunscreen

| Active Ingredient Zinc Oxide 17.1% | Sunscreen |
|--|-----------|
| Uses Helps prevent sunburn If used as directed with other sun protection decreases the risk of skin cancer and early | |
| Warnings For external use only | |
| Do not use | |

wnen using this product

· Keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if

rash occurs.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions For Sunscreen Use:

- Apply generously 15 minutes before sun exposure
- Children under 6 months of age: ask a doctor.
- Reapply at least every 2 hours.
- Use a water resistant sunscreen if swimming or sweating:
 - Immediately after towel drying.
 - at least every 2 hours.
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 a.m.-2 p.m.
 - · Wear long-sleeved shirts, pants, hats and sunglasses.

Other Information

Protect the product in this container from excessive heat and direct sun.



Helps prevent sunburn.

If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun.

| Active Ingredient Zinc Oxide 17.1% | Purpose Sunscreen |
|--|---|
| Uses Helps prevent sunburn If used as directed with other sun protection decreases the risk of skin cancer and early sl | |
| Warnings • For external use only | |
| Do not use • On damaged or broken skin | |
| When using this product Keep out of eyes. Rinse with water to remove. | |
| Stop use and ask a doctor if • rash occurs. | |
| Keep out of reach of children. If swallowed, get medical help or contact a Poison | Control Center right away. |
| Directions For Sunscreen Use: Apply generously 15 minutes before sun exposure Children under 6 months of age: ask a doctor. Reapply at least every 2 hours. Use a water resistant sunscreen if swimming or sv Immediately after towel drying. at least every 2 hours. Sun Protection Measures. Spending time in the scancer and early skin aging. To decrease this risk, Broad Spectrum SPF value of 15 or higher and oth including: Limit time in the sun, especially from 10 a.m2 Wear long-sleeved shirts, pants, hats and sunglated to the sungle state of the sungle state st | sun increases your risk of skin regularly use a sunscreen with a er sun protection measures p.m. |

Inactive ingredients: ALOE BARBADENSIS LEAF JUICE, CYCLOPENTASILOXANE, DICAPRYLYL CARBONATE, GLYCERIN, AQUA/WATER/EAU, ALCOHOL DENAT. (SD ALCOHOL 40-B), LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE, HYDRATED SILICA, DIMETHICONE, POLYGLYCERYL-3 POLYDIMETHYLSILOXYETHYL DIMETHICONE, ISOHEXADECANE MAGNESIUM SUI FATE POLYHYDROXYSTEABIC ACID. TRIFTHOXYSII YI FTHYI



170 g. NDC: 79458-000-07

For external use only

| Active Ingredient Zinc Oxide 17.1% | Purpose Sunscreen |
|--|----------------------|
| Uses Helps prevent sunburn If used as directed with other sun protection decreases the risk of skin cancer and early s | |
| <i>Warnings</i> • For external use only | |
| Do not use • On damaged or broken skin | |
| • | |
| When using this product Keep out of eyes. Rinse with water to remove. | |

Directions For Sunscreen Use:

- Apply generously 15 minutes before sun exposure
- Children under 6 months of age: ask a doctor.
- Reapply at least every 2 hours.
- Use a water resistant sunscreen if swimming or sweating:
 - Immediately after towel drying.
 - at least every 2 hours.
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 a.m.-2 p.m.
 - Wear long-sleeved shirts, pants, hats and sunglasses.

Other Information

Protect the product in this container from excessive heat and direct sun.

Inactive ingredients: ALOE BARBADENSIS LEAF JUICE, CYCLOPENTASILOXANE, DICAPRYLYL CARBONATE, GLYCERIN, AQUA/WATER/EAU, ALCOHOL DENAT. (SD ALCOHOL 40-B), LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE, HYDRATED SILICA, DIMETHICONE, POLYGLYCERYL-3 POLYDIMETHYLSILOXYETHYL DIMETHICONE, ISOHEXADECANE, MAGNESIUM SULFATE, POLYHYDROXYSTEARIC ACID, TRIETHOXYSILYLETHYL POLYDIMETHYLSILOXYETHYL HEXYL DIMETHICONE, DIMETHICONE/PEG-10/15 CROSSPOLYMER, RICINUS COMMUNIS (CASTOR) SEED OIL, LIMONENE, CAPRYLIC ACID, CAPRYLIC/CAPRIC TRIGLYCERIDE, CITRUS AURANTIUM DULCIS (ORANGE) PEEL EXTRACT, CITRUS GRANDIS (GRAPEFRUIT) PEEL EXTRACT, LECITHIN, PROPANEDIOL, ORBIGNYA PHALERATA SEED POWDER, LAURIC ACID, CALENDULA OFFICINALIS FLOWER EXTRACT, CITRUS LIMON (LEMON) PEEL EXTRACT, CITRUS TANGERINA (TANGERINE) PEEL EXTRACT, POTASSIUM SORBATE, HYDROXYETHYL ACRYLATE/SODIUM ACRYLOYLDIMETHYL TAURATE COPOLYMER, CAMELLIA OLEIFERA LEAF EXTRACT, DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER, ന TETRAHEXYLDECYL ASCORBATE, LYCIUM CHINENSE FRUIT EXTRACT, PYRUS MALUS (APPLE) FRUIT EXTRACT, ERYTHRITOL, PUNICA GRANATUM (POMEGRANATE) FRUIT EXTRACT, PHENOXYETHANOL, TOCOPHEROL, CITRAL, GLUCOSE, HYDROLYZED PEA PROTEIN, SODIUM CHLORIDE, PLANKTON EXTRACT, SODIUM BENZOATE, HOMARINE HCL, SODIUM SUCCINATE, ARABIDOPSIS THALIANA EXTRACT, MICROCOCCUS LYSATE, ERGOTHIONEINE. Questions Call 800-796-7546

Manufactured for IMAGE Skincare, Lantana, FL 33462 USA Manufactured in USA from globally sourced ingredients and components www.ImageSkincare.com

BB-402-V2

N

0

8

Area

170 g. NDC: 79458-001-00

Drug Facts

Do not use on damaged or broken skin.

| Active Ingredient Zinc Oxide 17.1% | Sunscreen |
|--|---|
| Uses Helps prevent sunburn If used as directed with other sun protection measure decreases the risk of skin cancer and early skin aging | |
| Warnings • For external use only | |
| Do not use • On damaged or broken skin | |
| When using this product Keep out of eyes. Rinse with water to remove. | |
| Stop use and ask a doctor if • rash occurs. | |
| Keep out of reach of children.If swallowed, get medical help or contact a Poison Control | Center right away. |
| Directions For Sunscreen Use: Apply generously 15 minutes before sun exposure Children under 6 months of age: ask a doctor. Reapply at least every 2 hours. Use a water resistant sunscreen if swimming or sweating: Immediately after towel drying. at least every 2 hours. Sun Protection Measures. Spending time in the sun increancer and early skin aging. To decrease this risk, regular Broad Spectrum SPF value of 15 or higher and other sun protection for the sun increase the sum of the sun increase the sum of t | eases your risk of skin rly use a sunscreen with a |

- Limit time in the sun, especially from 10 a.m.-2 p.m.
- · Wear long-sleeved shirts, pants, hats and sunglasses.

Other Information

· Protect the product in this container from excessive heat and direct sun.

Inactive ingredients: ALOE BARBADENSIS LEAF JUICE, CYCLOPENTASILOXANE, DICAPRYLYL CARBONATE, GLYCERIN, AQUA/WATER/EAU, ALCOHOL DENAT. (SD ALCOHOL 40-B), LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE, HYDRATED SILICA, DIMETHICONE, POLYGLYCERYL-3 POLYDIMETHYLSILOXYETHYL DIMETHICONE, ISOHEXADECANE, MAGNESIUM SULFATE, POLYHYDROXYSTEARIC ACID, TRIETHOXYSILYLETHYL POLYDIMETHYLSILOXYETHYL HEXYL DIMETHICONE, DIMETHICONE/PEG-10/15 CROSSPOLYMER, RICINUS COMMUNIS (CASTOR) SEED OIL, LIMONENE, CAPRYLIC ACID, CAPRYLIC/CAPRIC TRIGLYCERIDE, CITRUS AURANTIUM DULCIS (ORANGE) PEEL EXTRACT, CITRUS GRANDIS (GRAPEFRUIT) PEEL EXTRACT, LECITHIN, PROPANEDIOL, ORBIGNYA PHALERATA SEED POWDER, LAURIC ACID, CALENDULA OFFICINALIS FLOWER EXTRACT, CITRUS LIMON (LEMON) PEEL EXTRACT, CITRUS TANGERINA (TANGERINE) PEEL EXTRACT, POTASSIUM SORBATE, HYDROXYETHYL ACRYLATE/SODIUM ACRYLOYLDIMETHYL TAURATE COPOLYMER, CAMELLIA OLEIFERA LEAF EXTRACT, DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER, TETRAHEXYLDECYL ASCORBATE, LYCIUM CHINENSE FRUIT EXTRACT, PYRUS MALUS (APPLE) FRUIT EXTRACT, ERYTHRITOL, PUNICA GRANATUM (POMEGRANATE) FRUIT EXTRACT, PHENOXYETHANOL, TOCOPHEROL, CITRAL, GLUCOSE, HYDROLYZED PEA PROTEIN, SODIUM

| CHLORIDE, PLANKTON EXTRACT, SODIUM BENZOATE, HOMARINE HCL, SODIUM SU ARABIDOPSIS THALIANA EXTRACT, MICROCOCCUS LYSATE, ERGOTHIONEINE. Questions Call 800-796-7546 | CCINATE, | 9984 01 |
|--|-----------|---------|
| Manufactured for IMAGE Skincare, Lantana, FL 33462 USA Manufactured in USA from globally sourced ingredients and components www.ImageSkincare.com | BB-402-V2 | |
| Area | | |
| 70 g. NDC: 79458-001-00 | | |

When using this product keep out of eyes. Rinse with water to remove.

| Active Ingredient Zinc Oxide 17.1% | Purpose Sunscreen |
|---|------------------------------|
| Uses Helps prevent sunburn If used as directed with other sun protection decreases the risk of skin cancer and early successes the risk of skin cancer and early successes | |
| Warnings • For external use only | |
| Do not use • On damaged or broken skin | |
| When using this product Keep out of eyes. Rinse with water to remove. | |
| Stop use and ask a doctor if • rash occurs. | |
| Keep out of reach of children. If swallowed, get medical help or contact a Poison | n Control Center right away. |
| Directions For Sunscreen Use: Apply generously 15 minutes before sun exposure Children under 6 months of age: ask a doctor. Reapply at least every 2 hours. Use a water resistant sunscreen if swimming or s Immediately after towel drying. at least every 2 hours. Sun Protection Measures. Spending time in the | weating: |

cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:

- Limit time in the sun, especially from 10 a.m.-2 p.m.
- · Wear long-sleeved shirts, pants, hats and sunglasses.

Other Information

Protect the product in this container from excessive heat and direct sun.

Inactive ingredients: ALOE BARBADENSIS LEAF JUICE, CYCLOPENTASILOXANE, DICAPRYLYL CARBONATE, GLYCERIN, AQUA/WATER/EAU, ALCOHOL DENAT. (SD ALCOHOL 40-B), LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE, HYDRATED SILICA, DIMETHICONE, POLYGLYCERYL-3 POLYDIMETHYLSILOXYETHYL DIMETHICONE, ISOHEXADECANE, MAGNESIUM SULFATE, POLYHYDROXYSTEARIC ACID, TRIETHOXYSILYLETHYL POLYDIMETHYLSILOXYETHYL HEXYL DIMETHICONE, DIMETHICONE/PEG-10/15 CROSSPOLYMER, RICINUS COMMUNIS (CASTOR) SEED OIL, LIMONENE, CAPRYLIC ACID, CAPRYLIC/CAPRIC TRIGLYCERIDE, CITRUS AURANTIUM DULCIS (ORANGE) PEEL EXTRACT, CITRUS GRANDIS (GRAPEFRUIT) PEEL EXTRACT, LECITHIN, PROPANEDIOL, ORBIGNYA PHALERATA SEED POWDER, LAURIC ACID, CALENDULA OFFICINALIS FLOWER EXTRACT, CITRUS LIMON (LEMON) PEEL EXTRACT, CITRUS TANGERINA (TANGERINE) PEEL EXTRACT, POTASSIUM SORBATE, N HYDROXYETHYL ACRYLATE/SODIUM ACRYLOYLDIMETHYL TAURATE COPOLYMER, CAMELLIA OLEIFERA LEAF EXTRACT, DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER, ∞ TETRAHEXYLDECYL ASCORBATE, LYCIUM CHINENSE FRUIT EXTRACT, PYRUS MALUS (APPLE) FRUIT EXTRACT, ERYTHRITOL, PUNICA GRANATUM (POMEGRANATE) FRUIT EXTRACT, PHENOXYETHANOL, TOCOPHEROL, CITRAL, GLUCOSE, HYDROLYZED PEA PROTEIN, SODIUM CHLORIDE, PLANKTON EXTRACT, SODIUM BENZOATE, HOMARINE HCL, SODIUM SUCCINATE, ARABIDOPSIS THALIANA EXTRACT, MICROCOCCUS LYSATE, ERGOTHIONEINE. Questions တ Call 800-796-7546 5

Manufactured for IMAGE Skincare, Lantana, FL 33462 USA Manufactured in USA from globally sourced ingredients and components www.ImageSkincare.com


BB-402-V2

8

Area

170 g. NDC: 79458-001-00

Stop use and ask a doctor if rash occurs.



Warnings

For external use only

Do not use

• On damaged or broken skin

When using this product

Keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if

rash occurs.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions For Sunscreen Use:

- Apply generously 15 minutes before sun exposure
- Children under 6 months of age: ask a doctor.
- · Reapply at least every 2 hours.
- Use a water resistant sunscreen if swimming or sweating:
 - Immediately after towel drying.
 - at least every 2 hours.
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 a.m.-2 p.m.
 - Wear long-sleeved shirts, pants, hats and sunglasses.


Other Information

Protect the product in this container from excessive heat and direct sun.

Inactive ingredients: ALOE BARBADENSIS LEAF JUICE, CYCLOPENTASILOXANE, DICAPRYLYL CARBONATE, GLYCERIN, AQUA/WATER/EAU, ALCOHOL DENAT. (SD ALCOHOL 40-B), LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE, HYDRATED SILICA, DIMETHICONE, POLYGLYCERYL-3 POLYDIMETHYLSILOXYETHYL DIMETHICONE, ISOHEXADECANE, MAGNESIUM SULFATE, POLYHYDROXYSTEARIC ACID, TRIETHOXYSILYLETHYL POLYDIMETHYLSILOXYETHYL HEXYL DIMETHICONE, DIMETHICONE/PEG-10/15 CROSSPOLYMER, RICINUS COMMUNIS (CASTOR) SEED OIL, LIMONENE, CAPRYLIC ACID, CAPRYLIC/CAPRIC TRIGLYCERIDE, CITRUS AURANTIUM DULCIS (ORANGE) PEEL EXTRACT, CITRUS GRANDIS (GRAPEFRUIT) PEEL EXTRACT, LECITHIN, PROPANEDIOL, ORBIGNYA PHALERATA SEED

POWDER, LAURIC ACID, CALENDULA OFFICINALIS FLOWER EXTRACT, CITRUS LIMON (LEMON) PEEL EXTRACT, CITRUS TANGERINA (TANGERINE) PEEL EXTRACT, POTASSIUM SORBATE, HYDROXYETHYL ACRYLATE/SODIUM ACRYLOYLDIMETHYL TAURATE COPOLYMER, CAMELLIA OLEIFERA LEAF EXTRACT, DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER, TETRAHEXYLDECYL ASCORBATE, LYCIUM CHINENSE FRUIT EXTRACT, PYRUS MALUS (APPLE) FRUIT EXTRACT, ERYTHRITOL, PUNICA GRANATUM (POMEGRANATE) FRUIT EXTRACT, PHENOXYETHANOL, TOCOPHEROL, CITRAL, GLUCOSE, HYDROLYZED PEA PROTEIN, SODIUM CHLORIDE, PLANKTON EXTRACT, SODIUM BENZOATE, HOMARINE HCL, SODIUM SUCCINATE, ARABIDOPSIS THALIANA EXTRACT, MICROCOCCUS LYSATE, ERGOTHIONEINE.

Questions Call 800-796-7546





170 g. NDC: 79458-001-00

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

| Drug Facts | |
|--|---|
| Active Ingredient Zinc Oxide 17.1% | Purpose Sunscreen |
| Uses Helps prevent sunburn If used as directed with other sun protection decreases the risk of skin cancer and early | |
| <i>Warnings</i> • For external use only | |
| Do not use On damaged or broken skin | |
| When using this product Keep out of eyes. Rinse with water to remove. | |
| Stop use and ask a doctor if • rash occurs. | |
| Keep out of reach of children. If swallowed, get medical help or contact a Poiso | on Control Center right away. |
| Directions For Sunscreen Use: Apply generously 15 minutes before sun exposur Children under 6 months of age: ask a doctor. Reapply at least every 2 hours. Use a water resistant sunscreen if swimming or a lmmediately after towel drying. at least every 2 hours. Sun Protection Measures. Spending time in the cancer and early skin aging. To decrease this rise Broad Spectrum SPF value of 15 or higher and o including: Limit time in the sun, especially from 10 a.m Wear long-sleeved shirts, pants, hats and sun | sweating: e sun increases your risk of skin sk, regularly use a sunscreen with a ther sun protection measures •2 p.m. |



170 g. NDC: 79458-001-00

Apply generously 15 minutes before sun exposure. Children under 6 months of age: ask a doctor. Reapply at least every 2 hours. Use a water resistant sunscreen if swimming or sweating: Immediately after towel drying. At least every 2 hours. Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including: Limit time in the sun, especially from 10am to 2pm. Wear long-sleeved shirts, pants, hats and sunglasses.

| Active Ingredient Zinc Oxide 17.1% | Sunscreen |
|--|-----------|
| Uses Helps prevent sunburn If used as directed with other sun protection decreases the risk of skin cancer and early s | |

· For external use only

Do not use

On damaged or broken skin

When using this product

Keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if

rash occurs.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions For Sunscreen Use:

- Apply generously 15 minutes before sun exposure
- Children under 6 months of age: ask a doctor.
- Reapply at least every 2 hours.
- Use a water resistant sunscreen if swimming or sweating:
 - Immediately after towel drying.
 - at least every 2 hours.
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 a.m.-2 p.m.
 - Wear long-sleeved shirts, pants, hats and sunglasses.

Other Information

Protect the product in this container from excessive heat and direct sun.


Inactive ingredients: ALOE BARBADENSIS LEAF JUICE, CYCLOPENTASILOXANE, DICAPRYLYL CARBONATE, GLYCERIN, AQUA/WATER/EAU, ALCOHOL DENAT. (SD ALCOHOL 40-B), LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE, HYDRATED SILICA, DIMETHICONE, POLYGLYCERYL-3 POLYDIMETHYLSILOXYETHYL DIMETHICONE, ISOHEXADECANE, MAGNESIUM SULFATE, POLYHYDROXYSTEARIC ACID, TRIETHOXYSILYLETHYL POLYDIMETHYLSILOXYETHYL HEXYL DIMETHICONE, DIMETHICONE/PEG-10/15 CROSSPOLYMER, RICINUS COMMUNIS (CASTOR) SEED OIL, LIMONENE, CAPRYLIC ACID, CAPRYLIC/CAPRIC TRIGLYCERIDE, CITRUS AURANTIUM DULCIS (ORANGE)

PEEL EXTRACT, CITRUS GRANDIS (GRAPEFRUIT) PEEL EXTRACT, LECITHIN, PROPANEDIOL, ORBIGNYA PHALERATA SEED

POWDER, LAURIC ACID, CALENDULA OFFICINALIS FLOWER EXTRACT, CITRUS LIMON (LEMON) PEEL EXTRACT, CITRUS TANGERINA (TANGERINE) PEEL EXTRACT, POTASSIUM SORBATE, HYDROXYETHYL ACRYLATE/SODIUM ACRYLOYLDIMETHYL TAURATE COPOLYMER, CAMELLIA OLEIFERA LEAF EXTRACT, DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER, TETRAHEXYLDECYL ASCORBATE, LYCIUM CHINENSE FRUIT EXTRACT, PYRUS MALUS (APPLE) FRUIT EXTRACT, ERYTHRITOL, PUNICA GRANATUM (POMEGRANATE) FRUIT EXTRACT, PHENOXYETHANOL, TOCOPHEROL, CITRAL, GLUCOSE, HYDROLYZED PEA PROTEIN, SODIUM CHLORIDE, PLANKTON EXTRACT, SODIUM BENZOATE, HOMARINE HCL, SODIUM SUCCINATE, ARABIDOPSIS THALIANA EXTRACT, MICROCOCCUS LYSATE, ERGOTHIONEINE.

Questions Call 800-796-7546

Manufactured for IMAGE Skincare, Lantana, FL 33462 USA Manufactured in USA from globally sourced ingredients and components www.ImageSkincare.com



BB-402-V2

170 g. NDC: 79458-001-00

Protect the product in this container from excessive heat and direct sun.

| Active Ingredient Zinc Oxide 17.1% | Purpose Sunscreen |
|--|----------------------|
| Uses | |
| Helps prevent sunburn If used as directed with other sun protection measures (se decreases the risk of skin cancer and early skin aging call | |
| Warnings • For external use only | |
| Do not use On damaged or broken skin | |
| When using this product Keep out of eyes. Rinse with water to remove. | |
| Stop use and ask a doctor if • rash occurs. | |
| Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Cente | r right away. |
| Directions For Sunscreen Use: Apply generously 15 minutes before sun exposure | |
| Children under 6 months of age: ask a doctor. | |
| Reapply at least every 2 hours.Use a water resistant sunscreen if swimming or sweating: | |
| Immediately after towel drying. at least every 2 hours. | |
| Sun Protection Measures. Spending time in the sun increases cancer and early skin aging. To decrease this risk, regularly use Broad Spectrum SPF value of 15 or higher and other sun protect | a sunscreen with a |
| Including: Limit time in the sun, especially from 10 a.m2 p.m. Wear long-sleeved shirts, pants, hats and sunglasses. | |

Inactive ingredients: ALOE BARBADENSIS LEAF JUICE, CYCLOPENTASILOXANE, DICAPRYLYL CARBONATE,




170 g. NDC: 79458-000-07

Aloe Barbadensis Leaf Juice, Cyclopentasiloxane, Dicaprylyl Carbonate, Glycerin, Agua/Water/Eau, Alcohol Denat. (SD Alcohol 40-B), Lauryl PEG-9 Polydimethylsiloxyethyl Dimethicone, Hydrated Silica, Dimethicone, Polyglyceryl-3 Polydimethylsiloxyethyl Dimethicone, Isohexadecane, Magnesium Sulfate, Polyhydroxystearic Acid, Triethoxysilylethyl Polydimethylsiloxyethyl Hexyl Dimethicone, Dimethicone/PEG-10/15 Crosspolymer, Ricinus Communis (Castor) Seed Oil, Limonene, Caprylic Acid, Caprylic/Capric Triglyceride, Citrus Aurantium Dulcis (Orange) Peel Extract, Citrus Grandis (Grapefruit) Peel Extract, Lecithin, Propanediol, Orbignya Phalerata Seed Powder, Lauric Acid, Calendula Officinalis Flower Extract, Citrus Limon (Lemon) Peel Extract, Citrus Tangerina (Tangerine) Peel Extract, Potassium Sorbate, Hydroxyethyl Acrylate/Sodium Acryloydimethyl Taurate Copolymer, Camellia Oleifera Leaf Extract, Dimethicone/Vinyl Dimethicone Crosspolymer, Tetrahexyldecyl Ascorbate, Lycium Chinense Fruit Extract, Pyrus Malus (Apple) Fruit Extract, Erythritol, Punica Granatum (Pomegranate) Fruit Extract, Phenoxyethanol, Tocopherol, Citral, Glucose, Hydrolyzed Pea Protein, Sodium Chloride, Plankton Extract, Sodium Benzoate, Homarine HCL, Sodium Succinate, Arabidopsis Thaliana Extract, Micrococcus Lysate, **Ergothioneine**

| Zinc Oxide 17.1% | Sunscreen |
|---|---|
| Uses Helps prevent sunburn If used as directed with other sun protection n decreases the risk of skin cancer and early skin | |
| Warnings • For external use only | |
| Do not use On damaged or broken skin | |
| When using this product Keep out of eyes. Rinse with water to remove. | |
| Stop use and ask a doctor if • rash occurs. | |
| Keep out of reach of children. If swallowed, get medical help or contact a Poison (| Control Center right away. |
| Directions For Sunscreen Use: Apply generously 15 minutes before sun exposure Children under 6 months of age: ask a doctor. Reapply at least every 2 hours. Use a water resistant sunscreen if swimming or sw Immediately after towel drying. at least every 2 hours. Sun Protection Measures. Spending time in the su cancer and early skin aging. To decrease this risk, Broad Spectrum SPF value of 15 or higher and other including: Limit time in the sun, especially from 10 a.m2 performance. Wear long-sleeved shirts, pants, hats and sunglated of the sungle of | un increases your risk of skin regularly use a sunscreen with a er sun protection measures p.m. |
| Protect the product in this container from excessive here | at and direct sun. |
| Inactive ingredients: ALOE BARBADENSIS LEAF JUICE, CYCLOPEN GLYCERIN, AQUA/WATER/EAU, ALCOHOL DENAT. (SD ALCOHOL 40-B), LAU DIMETHICONE, HYDRATED SILICA, DIMETHICONE, POLYGLYCERYL-3 POLYD ISOHEXADECANE, MAGNESIUM SULFATE, POLYHYDROXYSTEARIC ACID, TR POLYDIMETHYLSILOXYETHYL HEXYL DIMETHICONE, DIMETHICONE/PEG-10, (CASTOR) SEED OIL, LIMONENE, CAPRYLIC ACID, CAPRYLIC/CAPRIC TRIGLY PEEL EXTRACT, CITRUS GRANDIS (GRAPEFRUIT) PEEL EXTRACT, LECITHIN, POWDER, LAURIC ACID, CALENDULA OFFICINALIS FLOWER EXTRACT, CITRU | RYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHYLSILOXYETHYL DIMETHICONE, HETHOXYSILYLETHYL /15 CROSSPOLYMER, RICINUS COMMUNIS 'CERIDE, CITRUS AURANTIUM DULCIS (ORANGE) PROPANEDIOL, ORBIGNYA PHALERATA SEED |

POWDER, LAURIC ACID, CALENDULA OFFICINALIS FLOWER EXTRACT, CITRUS LIMON (LEMON) PEEL EXTRACT, CITRUS TANGERINA (TANGERINE) PEEL EXTRACT, POTASSIUM SORBATE, HYDROXYETHYL ACRYLATE/SODIUM ACRYLOYLDIMETHYL TAURATE COPOLYMER, CAMELLIA OLEIFERA LEAF EXTRACT, DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER, TETRAHEXYLDECYL ASCORBATE, LYCIUM CHINENSE FRUIT EXTRACT, PYRUS MALUS (APPLE) FRUIT EXTRACT, ERYTHRITOL, PUNICA GRANATUM (POMEGRANATE) FRUIT EXTRACT, PHENOXYETHANOL, TOCOPHEROL, CITRAL, GLUCOSE, HYDROLYZED PEA PROTEIN, SODIUM CHI ORIDE, PLANKTON EXTRACT, SODIUM BENZOATE, HOMARINE HCL, SODIUM SUCCINATE






170 g. NDC: 79458-001-00

Prevention plus daily hydrating moisturizer 30 SPF/FPS

BACKBAR US

PREV PLUS HYDRATING 30 PROFESSIONAL (PPHY-BB-001-GL).ai 9.28.18







170 g. NDC: 79458-001-00

| MOISTURIZER | | | | | |
|--|---|-----------------|-----------------------|----------|----------------------|
| zinc oxide cream | | | | | |
| Product Information | | | | | |
| Product Information | | | | | |
| Product Type | HUMAN OTC DRUG | Item Cod | e (Source) | NDC:7945 | 58-001 |
| Route of Administration | TOPICAL | | | | |
| | | | | | |
| | | | | | |
| Active Ingredient/Active | Moiety | | | | |
| Ingred | lient Name | | Basis of Strengt | h St | rength |
| | | > | | | |
| ZINC OXIDE (UNII: SOI2LOH54Z) | (ZINC OXIDE - UNII:SOI2LOH | 54Z) | ZINC OXIDE | 171 g | in 1000 g |
| ZINC OXIDE (UNII: SOI2LOH54Z) | (ZINC OXIDE - UNII:SOI2LOH | 542) | ZINC OXIDE | 171 g | in 1000 g |
| | (ZINC OXIDE - UNII:SOI2LOH | 542) | ZINC OXIDE | 171 g | in 1000 g |
| Inactive Ingredients | (ZINC OXIDE - UNII:SOI2LOH | 542) | ZINC OXIDE | 171 g | in 1000 g |
| | (ZINC OXIDE - UNII:SOI2LOH Ingredient Nan | | ZINC OXIDE | 171 g | in 1000 g Strengt |
| | | | ZINC OXIDE | 171 g | |
| Inactive Ingredients | Ingredient Nan | | ZINC OXIDE | 171 g | |
| Inactive Ingredients ALCOHOL (UNII: 3K9958V90M) | Ingredient Nan VER (UNII: P0M7O4Y7YD) | | ZINC OXIDE | 171 g | |
| Inactive Ingredients ALCOHOL (UNII: 3K9958V90M) CALENDULA OFFICINALIS FLOV POTASSIUM SORBATE (UNII: 1V CAMELLIA OLEIFERA LEAF (UNII | Ingredient Nan VER (UNII: POM7O4Y7YD) PU26JZZ4) : 5077EL0C60) | ne | | | |
| Inactive Ingredients ALCOHOL (UNII: 3K9958V90M) CALENDULA OFFICINALIS FLOW POTASSIUM SORBATE (UNII: 1V | Ingredient Nan VER (UNII: POM7O4Y7YD) PU26JZZ4) : 5077EL0C60) | ne | | | |
| Inactive Ingredients ALCOHOL (UNII: 3K9958V90M) CALENDULA OFFICINALIS FLOV POTASSIUM SORBATE (UNII: 1V CAMELLIA OLEIFERA LEAF (UNII | Ingredient Nan VER (UNII: POM7O4Y7YD) PU26JZZ4) : 5077EL0C60) IMETHYLSILOXYETHYL HE | ne | | | |
| Inactive Ingredients ALCOHOL (UNII: 3K9958V90M) CALENDULA OFFICINALIS FLOW POTASSIUM SORBATE (UNII: 1V CAMELLIA OLEIFERA LEAF (UNII TRIETHOXYSILYLETHYL POLYD | Ingredient Nam VER (UNII: P0M7O4Y7YD) PU26JZZ4) : 5077EL0C60) IMETHYLSILOXYETHYL HE G8P9) | ne | | | |
| Inactive Ingredients ALCOHOL (UNII: 3K9958V90M) CALENDULA OFFICINALIS FLOW POTASSIUM SORBATE (UNII: 1VI CAMELLIA OLEIFERA LEAF (UNII TRIETHOXYSILYLETHYL POLYD HYDRATED SILICA (UNII: Y607T4 DIMETHICONE (UNII: 92RU3N3Y1 ARABIDOPSIS THALIANA (UNII: 4 | Ingredient Nam VER (UNII: POM7O4Y7YD) PU26JZZ4) : 5077EL0C60) IMETHYLSILOXYETHYL HE G8P9) O) M3L60HQ81) | ne | | | |
| Inactive Ingredients ALCOHOL (UNII: 3K9958V90M) CALENDULA OFFICINALIS FLOW POTASSIUM SORBATE (UNII: 1V/ CAMELLIA OLEIFERA LEAF (UNII TRIETHOXYSILYLETHYL POLYD HYDRATED SILICA (UNII: Y607T4 DIMETHICONE (UNII: 92RU3N3Y1 ARABIDOPSIS THALIANA (UNII: A CITRUS MAXIMA FRUIT RIND (U | Ingredient Nam VER (UNII: P0M7O4Y7YD) PU26JZZ4) : 5077EL0C60) IMETHYLSILOXYETHYL HE G8P9) O) AI3L60HQ81) NII: 5NX3G75CA6) | ne | | | |
| Inactive Ingredients ALCOHOL (UNII: 3K9958V90M) CALENDULA OFFICINALIS FLOW POTASSIUM SORBATE (UNII: 1V) CAMELLIA OLEIFERA LEAF (UNII TRIETHOXYSILYLETHYL POLYD HYDRATED SILICA (UNII: Y607T4 DIMETHICONE (UNII: 92RU3N3Y1 ARABIDOPSIS THALIANA (UNII: 4 | Ingredient Nam VER (UNII: P0M7O4Y7YD) PU26JZZ4) : 5077EL0C60) IMETHYLSILOXYETHYL HE G8P9) O) AI3L60HQ81) NII: 5NX3G75CA6) | ne | | | |
| Inactive Ingredients ALCOHOL (UNII: 3K9958V90M) CALENDULA OFFICINALIS FLOW POTASSIUM SORBATE (UNII: 1V/ CAMELLIA OLEIFERA LEAF (UNII TRIETHOXYSILYLETHYL POLYD HYDRATED SILICA (UNII: Y607T4 DIMETHICONE (UNII: 92RU3N3Y1 ARABIDOPSIS THALIANA (UNII: A CITRUS MAXIMA FRUIT RIND (U | Ingredient Nam VER (UNII: P0M7O4Y7YD) PU26JZZ4) : 5077EL0C60) IMETHYLSILOXYETHYL HE G8P9) O) NII: 5NX3G75CA6) 057) | ופ XYL DIMET | HICONE (UNII: X75PL53 | 3TZJ) | |

| | arketing | Application Number or Mon | ograph | Marketing Start | Marketing End |
|--------|----------------------|--|----------|--------------------------|-----------------------|
| Mar | keting | Information | | | |
| | | | | | |
| 00 | | | | | |
| | :79458-001- | 170 g in 1 TUBE; Type 0: Not a Combin Product | nation | 03/17/2022 | |
| # Ite | em Code | Package Description | | Marketing Start Date | Marketing End Date |
| Pack | aging | | | | |
| | | | | | |
| | | | | | |
| | | RA96B954X6) | | | |
| | | RIGLYCERIDES (UNII: C9H2L21V7U) | | | |
| | | NII: HL2549X1EY) | | | |
| | | G-10/15 CROSSPOLYMER (UNII: 21AS) | 8B1BSS) | | |
| | | (UNII: 918X10UF1E) | | | |
| | | POLYDIMETHYLSILOXYETHYL DIMET | - | (4000 MPA.S) (UNII: RLA? | 2U05Z4O) |
| | | TE HEXAHYDRATE (UNII: U16QOD6C4E | :) | | |
| | | CHLORIDE (UNII: 8866LNG61N) | | | |
| | | E (UNII: OJ245FE5EU) | | | |
| | | NII: 56687D1Z4D) | | | |
| | • | E (UNII: 451W47IQ8X) | | | |
| | | | | | |
| | | LYDIMETHYLSILOXYETHYL DIMETHI | | III: 25G622K2RA) | |
| | | INII: GFD7C86Q1W) | | | |
| | | 1160N9NU9U) | | | |
| | | UNII: ZY81Z83H0X) | | | |
| | | LASCORBATE (UNII: 9LBV3F07AZ) | | | |
| | | II: 5965N8W85T) | , | | |
| | | EARIC ACID (2300 MW) (UNII: YXH47A | | | |
| | | ATE, UNSPECIFIED (UNII: DE08037SA | 3) | | |
| | | D5340Y2I9G) | | | |
| | | ROSE (UNII: 5SL0G7R0OK) | | | |
| | L (UNII: T7EU | 009VPP) | | | |
| PEA PR | ROTEIN (UNI | : 7Q50F46595) | | | |
| APPLE | (UNII: B423V | GH5S9) | | | |
| GLYCE | RIN (UNII: PE | DC6A3C0OX) | | | |
| ERGOT | HIONEINE (| UNII: BDZ 3DQM98W) | | | |
| PHENO | XYETHANO | L (UNII: HIE492ZZ3T) | | | |
| LYCIUM | M CHINENSE | FRUIT (UNII: TG711Q7A1Q) | | | |
| госоя | PHEROL (UN | II: R0ZB2556P8) | | | |
| ORANG | GE PEEL (UN | II: TI9T76XD44) | | | |
| DIMET | HICONE/VIN | YL DIMETHICONE CROSSPOLYMER | (SOFT PA | RTICLE) (UNII: 9E4CO0W6 | C5) |
| NATEF | R (UNII: 059Q | F0KO0R) | | | |
| ANACY | STIS NIDUL | ANS EXTRACT (UNII: UV4FTL6UAW) | | | |
| 4ICRO | COCCUS LI | JTEUS (UNII: LV6L29Z6AX) | | | |
| DICAPI | RYLYL CARB | ONATE (UNII: 609A3V1SUA) | | | |
| | | E 5 (UNII: 0THT5PCI0R) | | | |
| YCLO | METHICON | | | | |

| Category | Citation | Date | Date |
|------------------------|----------|------------|------|
| OTC monograph final pa | art352 | 03/17/2022 | |

Labeler - Natural Technology, LLC (618561906)

Revised: 3/2022

Natural Technology, LLC