

**DENTAL CITY TOPICAL ANESTHETIC- benzocaine gel**  
**ICP Inc DBA Dental City**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Active Ingredients**

Benzocaine 200mg (in each g)

**Purpose**

Oral Anesthetic

**Use**

For oral mucosal use. Use only as directed by dentist. For the temporary relief of pain due to minor dental procedures.

**Warnings**

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy Alert:** Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

**Do not use**

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

**When using this product** Avoid contact with eyes. If contact occurs, flush with water.

**Do not exceed recommended dosage.** If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

**If pregnant or breast feeding,** ask a physician before use.

**Directions**

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

**Keep out of reach of children.**

**Other Information**

Store at 59°-86°F (15°-30°C). Protect from freezing.

**Inactive Ingredients**

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.


**Questions or comments?**

1-800-353-9595 / [dentalcity.com](http://dentalcity.com)

PEEL BACK to review drug facts



Manufactured for:  
Dental City  
Green Bay, WI 54311  
dentalcity.com  
1-800-353-9595



**TOPICAL  
ANESTHETIC GEL  
BENZOCAINE 20%**

MADE IN USA

GLUTEN FREE  
1 oz (30 ml)

directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.

- for teething
- in children under 2 years of age

**When using this product**

Avoid contact with eyes. If contact occurs, flush with water.

**Do not exceed recommended dosage.**

If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

**If pregnant or breast feeding,** ask a physician before use.

**Keep out of reach of children.**

**Directions**

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

**Other information**

Store at 59° - 86°F (15 - 30°C). Protect from freezing.

**Inactive ingredients**

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

**Questions or comments?**

1-800-353-9595 | dentalcity.com

99092951 Rev 10/2018

**Drug Facts**

**Active ingredients**

Benzocaine 200 mg (in each g)

**Purpose**

Oral Anesthetic

**Use**

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

**Warnings**

**Methemoglobinemia**

**warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy Alert:**

Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

**Do not use**

- for more than 7 days unless

**DENTAL CITY TOPICAL ANESTHETIC**

benzocaine gel

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69483-006
<b>Route of Administration</b>	DENTAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>BENZOCAINE</b> (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

**Inactive Ingredients**

Ingredient Name	Strength
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	

<b>FD&amp;C YELLOW NO. 5</b> (UNII: I753WB2F1M)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>FD&amp;C RED NO. 3</b> (UNII: PN2ZH5LOQY)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C GREEN NO. 3</b> (UNII: 3P3ONR6O1S)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
<b>D&amp;C GREEN NO. 5</b> (UNII: 8J6RDU8L9X)	

### Product Characteristics

<b>Color</b>	purple	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69483-006-30	30 g in 1 JAR; Type 0: Not a Combination Product	10/01/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	04/01/2014	

**Labeler** - ICP Inc DBA Dental City (808371371)