

FAMOTIDINE - ACID CONTROLLER- famotidine tablet, film coated
WinCo Foods, LLC

Famotidine - Acid Controller

Drug Facts

Active ingredient (in each tablet)

Famotidine, USP 10 mg

Famotidine, USP 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert

Do not use if you are allergic to famotidine or other acid reducers

Do not use

Famotidine USP, 10 mg

- with other acid reducers
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Famotidine USP, 20 mg

- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor.
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

Famotidine USP, 10 mg

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Famotidine USP, 20 mg

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 20° to 25° C (68° to 77° F)
- protect from moisture
- read the directions and warnings before use
- keep the carton. It contains important information.

Inactive ingredients

colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, iron oxide red (only for Famotidine USP, 10 mg), magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, pregelatinized starch, talc, titanium dioxide

Questions?

or to report an adverse event call **800-824-1706** Monday - Friday 9am - 4pm MST

DISTRIBUTED BY: WINCO FOODS, LLC, BOISE, ID 83704

PRINCIPAL DISPLAY PANEL - 10 mg Blister Pack Carton

*Compare to the active ingredient in PEPCID AC® Original Strength

WinCo™

FOODS

ORIGINAL STRENGTH

Acid Controller

FAMOTIDINE TABLETS,

USP 10 mg/
ACID REDUCER
JUST ONE TABLET Prevents & Relieves Heartburn
Due to Acid Indigestion

30
Tablets

5113374



5113374



Drug Facts

Active Ingredient (in each tablet)
Famotidine, USP 10 mg.....Acid reducer

Uses
■ relieves heartburn associated with acid indigestion and sour stomach
■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use
■ With other acid reducers
■ If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
■ frequent chest pain
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain

Directions
■ adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
■ to prevent symptoms, swallow 1 tablet with a glass of water 60 minutes before eating food or drinking beverages that cause heartburn
■ do not use more than 2 tablets in 24 hours
■ children under 12 years: ask a doctor

Drug Facts (continued)
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ heartburn with lightheadedness, sweating, or dizziness
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).



ORIGINAL STRENGTH Acid Controller

30 TABLETS

*Compare to the active ingredient in PEPCID AC® Original Strength

WinCo FOODS

ORIGINAL STRENGTH Acid Controller

FAMOTIDINE TABLETS,
USP 10 mg/
ACID REDUCER

JUST ONE TABLET Prevents & Relieves Heartburn Due to Acid Indigestion



30
Tablets

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Expiration Date:

Batch No.

Varnish Free

TAMPER EVIDENT: DO NOT USE IF THE CARTON OR INDIVIDUAL BLISTER UNIT IS OPEN OR TORN.

DISTRIBUTED BY: WINCO FOODS, LLC, BOISE, ID 83704

SEE END PANEL FOR BATCH NUMBER AND EXPIRATION DATE.

* This product is not manufactured or distributed by MCNEIL CONSUMER PHARMACEUTICALS, distributor of Pepcid AC® Original Strength.

The Supermarket Low Price Leader®


100% SATISFACTION GUARANTEE

Comments & Questions? 800-824-1706 • www.wincofoods.com



0914

drinking certain foods and beverages.



PRINCIPAL DISPLAY PANEL - 20 mg Blister Pack Carton

*Compare to the active ingredient in PEPCID AC® Maximum Strength

WinCo™

FOODS

MAXIMUM STRENGTH

Acid Controller

FAMOTIDINE TABLETS,

USP 20 mg/

ACID REDUCER

JUST ONE TABLET Prevents & Relieves Heartburn

Due to Acid Indigestion

25

Tablets

WinCo
FOODS

MAXIMUM STRENGTH Acid Controller

25 TABLETS

*Compare to the active ingredient in PEPCID AC® Maximum Strength

WinCo
FOODS

MAXIMUM STRENGTH Acid Controller

FAMOTIDINE TABLETS,
USP 20 mg/
ACID REDUCER

JUST ONE TABLET Prevents & Relieves Heartburn
Due to Acid Indigestion



25
Tablets

Tips for Managing Heartburn

- Do not lie flat or bend over after eating ■ Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime ■ Raise the head of your bed ■ Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals ■ If overweight, lose weight ■ Quit smoking

Expiration Date:

Batch No.

Varnish Free

TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.

SEE END PANEL FOR BATCH NUMBER AND EXPIRATION DATE.

This product is not manufactured or distributed by MICHEL CONSUMER PHARMACEUTICALS, distributor of PEPCID AC® Maximum Strength.

The Supermarket Low Price Leader®

**100% SATISFACTION
GUARANTEE**

Comments & Questions?
800-824-1706 • www.wincofoods.com



09/4

Drug Facts (continued)

Other Information

- store at 20° to 25° C (68° to 77° F)
- protect from moisture
- read the directions and warnings before use
- keep the carton. It contains important information.

Inactive ingredients colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, pregelatinized starch, talc, titanium dioxide

Questions?

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting

Directions

- adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

5113376



Drug Facts (in each tablet) Famotidine, USP 20 mgAcid reducer	Drug Facts (continued) ■ stomach pain ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating or dizziness
Active Ingredient Purpose	

FAMOTIDINE - ACID CONTROLLER

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67091-242
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	10 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE (1200000 MW) (UNII: RFW2ET671P)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	035
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-242-30	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	01/08/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090283	05/01/2012	

FAMOTIDINE - ACID CONTROLLER

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67091-241
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE (1200000 MW) (UNII: RFW2ET671P)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6B30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	036
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-241-25	25 in 1 BOTTLE; Type 0: Not a Combination Product	01/08/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090283	05/01/2012	

Labeler - WinCo Foods, LLC (056098817)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		184769029	MANUFACTURE(67091-241, 67091-242)

Revised: 9/2018

WinCo Foods, LLC