

STOMACH RELIEF- bismuth subsalicylate tablet
Chain Drug Marketing Association, Inc.

Quality Choice 44-346

Active ingredient (in each caplet)

Bismuth subsalicylate 262 mg

Purpose

Upset stomach reliever/antidiarrheal

Uses

relieves:

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - belching
 - indigestion
 - gas
 - fullness
 - heartburn
 - nausea

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are:

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- diabetes
- arthritis
- anticoagulation (thinning the blood)
- gout

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- swallow with water; do not chew
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- adults and children 12 years and over: 2 caplets every 1/2 to 1 hour as needed. Do not exceed 8 doses (16 caplets) in 24 hours.
- do not use for more than 2 days unless directed by a doctor
- use until diarrhea stops, but not more than 2 days
- children under 12 years: ask a doctor

Other information

- **each caplet contains:** calcium 20 mg, salicylate 103 mg
- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- avoid excessive heat
- see end flap for expiration date and lot number

Inactive ingredients

calcium carbonate, corn starch, D&C red #27 aluminum lake, D&C red #30 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, povidone, silicon dioxide, sodium starch glycolate, stearic acid

Questions or comments?

1-800-426-9391

Principal display panel

**QC®
QUALITY
CHOICE**

NDC 83324-060-40

*Compare to the
Active Ingredient in
Pepto-Bismol®

**Stomach Relief
Bismuth Subsalicylate 262 mg
Upset Stomach Reliever/Antidiarrheal**

Relieves Nausea, Heartburn, Indigestion,
Upset Stomach, Diarrhea

40 Caplets

Actual Size

**TAMPER EVIDENT: DO NOT USE IF
IMPRINTED SAFETY SEAL UNDER
CAP IS BROKEN OR MISSING**

*This product is not manufactured or distributed by
The Procter & Gamble Company, owner of the
registered trademark Pepto-Bismol®.

50844 REV0521B34610

Distributed by CDMA, Inc.

Novi, MI 48375

www.qualitychoice.com

Questions: 800-935-2362

SATISFACTION

100%

GUARANTEED®

No Print / No Varnish
Lot no. & Exp. date



TAMPER EVIDENT: DO NOT USE IF
IMPRINTED SAFETY SEAL UNDER
CAP IS BROKEN OR MISSING

B-0220-346-10-R
REV0521B34610

Drug Facts
Active Ingredient (in each caplet)
Bismuth subsalicylate 262 mg

Purpose
..... Upset stomach reliever/antidiarrheal

Uses relieves:
 ■ travelers' diarrhea ■ diarrhea
 ■ upset stomach due to overindulgence in food and drink, including: ■ heartburn
 ■ indigestion ■ nausea ■ gas
 ■ belching ■ fullness

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Contains salicylate. Do not take if you are:
 ■ allergic to salicylates (including aspirin)
 ■ taking other salicylate products

Do not use if you have
 ■ a bleeding problem ■ an ulcer
 ■ bloody or black stool

QC QUALITY CHOICE

NDC 83324-060-40

*Compare to the Active Ingredient in Pepto-Bismol®

Stomach Relief

Bismuth Subsalicylate 262 mg

Upset Stomach Reliever/Antidiarrheal

Relieves Nausea, Heartburn, Indigestion, Upset Stomach, Diarrhea



Actual Size

40 Caplets

QC QUALITY CHOICE

Stomach Relief

Bismuth Subsalicylate 262 mg

Upset Stomach Reliever/Antidiarrheal

Drug Facts (continued)
 ■ use until diarrhea stops, but not more than 2 days
 ■ children under 12 years: ask a doctor

Other information
 ■ each caplet contains: calcium 20 mg, salicylate 103 mg
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
 ■ avoid excessive heat
 ■ see end flap for expiration date and lot number

Inactive ingredients calcium carbonate, corn starch, D&C red #27 aluminum lake, D&C red #30 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, povidone, silicon dioxide, sodium starch glycolate, stearic acid

Questions or comments?
1-800-426-9391

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Pepto-Bismol®.
50844 REV0521B34610

SATISFACTION GUARANTEED
30-DAY MONEY BACK GUARANTEE

Distributed by CDMA, Inc.
Novi, MI 48476
www.cdma.com
Questions: 800-935-2362

Drug Facts (continued)
 Ask a doctor before use if you have
 ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for:
 ■ diabetes ■ arthritis ■ gout ■ anticoagulation (thinning the blood)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if
 ■ symptoms get worse
 ■ ringing in the ears or loss of hearing occurs
 ■ diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ do not take more than directed
 ■ swallow with water; do not chew
 ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea
 ■ adults and children 12 years and over: 2 caplets every ½ to 1 hour as needed. Do not exceed 8 doses (16 caplets) in 24 hours.
 ■ do not use for more than 2 days unless directed by a doctor

Quality Choice 44-346

STOMACH RELIEF

bismuth subsalicylate tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83324-060
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ, BISMUTH CATION - UNII:ZS9CD1I8YE)	BISMUTH SUBSALICYLATE	262 mg

Inactive Ingredients

Ingredient Name	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
D&C RED NO. 30 ALUMINUM LAKE (UNII: GE75M6YV5W)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	pink	Score	no score
Shape	OVAL	Size	16mm
Flavor		Imprint Code	44;346
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83324-060-40	1 in 1 CARTON	02/14/2025	
1		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	02/14/2025	

Labeler - Chain Drug Marketing Association, Inc. (011920774)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(83324-060)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(83324-060) , pack(83324-060)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(83324-060)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(83324-060)

Revised: 2/2026

Chain Drug Marketing Association, Inc.