

**KIRKLAND SIGNATURE NAPROXEN SODIUM- naproxen sodium tablet, film coated**

**Costco Wholesale Company**

-----

**Costco Wholesale Corp. Naproxen Sodium Tablets USP, 220 mg Drug Facts**

**Active ingredient (in each caplet)**

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none"> <li>• take 1 caplet every 8 to 12 hours while symptoms last</li> <li>• for the first dose you may take 2 caplets within the first hour</li> <li>• do not exceed 2 caplets in any 8- to 12-hour period</li> <li>• do not exceed 3 caplets in a 24-hour period</li> </ul>
Children under 12 years	<ul style="list-style-type: none"> <li>• ask a doctor</li> </ul>

**Other information**

- **each caplet contains:** sodium 22 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Inactive ingredients**

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

**Questions or comments?**

**1-800-719-9260**

**Principal Display Panel**

KIRKLAND Signature  
 COMPARE TO ALEVE® active ingredient  
 NAPROXEN SODIUM  
 Tablets USP, 220 mg  
 Pain Reliever/Fever Reducer  
 (NSAID)  
 Strength to Last 12 HOURS  
 Actual Size  
 400 CAPLETS†  
 †capsule-shaped tablets

DO NOT USE IF PRINTED  
 FOIL UNDER CAP IS  
 BROKEN OR MISSING

Packaged by:  
 Perrigo®, 515 Eastern Ave., Allegan, MI 49010 USA  
 1-800-719-9260  
 For: Costco Wholesale Corporation  
 999 Lake Drive, Issaquah, WA 98027 USA  
 www.costco.com

21V09073a

GLUTEN FREE



\*\*This product is not  
 manufactured or distributed  
 by Bayer HealthCare LLC,  
 distributor of Aleve®.



EXP. LOT  
 : 36679 5N F16

**KIRKLAND Signature**  
 COMPARE TO ALEVE® active ingredient\*\*  
 NDC 63981-368-79  
 ITM. / ART. 197197

# NAPROXEN SODIUM

Tablets USP, 220 mg

**Pain Reliever/Fever Reducer**  
 (NSAID)  
 Strength to Last 12 HOURS

**400 CAPLETS†**  
 †capsule-shaped tablets

Actual Size

**Drug Facts**

**Active Ingredient (in each caplet)**  
 Naproxen sodium 220 mg  
 (naproxen 200 mg) (NSAID)\*  
 \*nonsteroidal anti-inflammatory drug  
**Purposes**  
 Pain reliever/  
 fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - menstrual cramps
  - headache
  - toothache
  - the common cold
  - temporarily reduces fever

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- skin redness
- hives
- facial swelling
- asthma (wheezing)
- stomach bleeding
- stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
  - are age 60 or older
  - have had stomach ulcers or bleeding problems
  - take a blood thinning (anticoagulant) or steroid drug
  - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
  - have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

PEEL BACK HERE

**Drug Facts (continued)**

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if:

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are:

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

**When using this product:**

- take with food or milk if stomach upset occurs
- Stop use and ask a doctor if:
  - you experience any of the following signs of stomach bleeding:
    - feel faint
    - worn blood
    - have bloody or black stools
  - have stomach pain that does not get better
  - you have symptoms of heart problems or stroke:
    - chest pain
    - trouble breathing
    - weakness in one part or side of body
    - slurred speech
    - leg swelling
  - pain gets worse or lasts more than 10 days
  - tear gets worse or lasts more than 3 days
  - you have difficulty swallowing
  - It feels like the pill is stuck in your throat
  - redness or swelling is present in the painful area
  - any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than directed
  - the smallest effective dose should be used
  - drink a full glass of water with each dose
- Adults and children 12 years and older:
- take 1 caplet every 8 to 12 hours while symptoms last
  - do not exceed 2 caplets in any 8- to 12-hour period
  - do not exceed 3 caplets in a 24-hour period
- Children under 12 years:
- ask a doctor

**Other information**

each caplet contains: sodium 22 mg  
 store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).  
**Inactive ingredients:** FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polydioxane, talc, titanium dioxide

**Questions or comments?** 1-800-719-9260

naproxen sodium tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63981-368
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>NAPROXEN SODIUM</b> (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>FD&amp;C BLUE NO. 2 ALUMINUM LAKE</b> (UNII: 4AQJ3LG584)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	BLUE (Light Blue)	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	L368
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:63981-368-79	400 in 1 BOTTLE; Type 0: Not a Combination Product	06/04/2007	
2	NDC:63981-368-91	1 in 1 PACKAGE	02/03/2020	02/03/2020
2		6 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA074661	06/04/2007	

Revised: 7/2025

Costco Wholesale Company