

EZ2GO POLYETHYLENE GLYCOL 3350- polyethylene glycol 3350 powder, for solution

Valley Medical Products, LLC

ez2go Polyethylene Glycol 3350 Powder

Drug Facts

Active ingredient (in each dose)

Polyethylene Glycol 3350, 17 grams (cap filled to the indicated line)

Purpose

Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use

if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are

taking a prescription drug

When using this product

you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
- use once a day
- fill to top of white section in cap which is marked to indicate the correct dose (17g)
- stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch based thickeners used for difficulty swallowing
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- do not use more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20°-25° C (68°-77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive ingredients


none

Questions or comments?

1-800-403-9832

Package Labeling:





PAGE 12

PAGE 01

<p>Drug Facts (continued) ⑫</p> <p>"SEALED for YOUR PROTECTION" is missing, open or broken</p> <p>Inactive ingredients none</p> <p>Questions or comments? 1-800-403-9832</p>	<p>10mm Blank for Glue Area</p>	<p>This product is not manufactured or distributed by Schering-Plough Healthcare Products, Inc., the distributor of MiraLax®</p> <p>LOT: _____ EXP: _____</p> <p>Manufactured for, ©Copyright & Distributed by Valley Medical Products, LLC, Jacksonville, FL 32224 USA.</p> <p>All rights reserved. 1052422</p> <p>Code No.: KR/DRUGS/KTK/25/415/98</p> <p>SEE INSIDE FOR DRUG FACTS</p> <p>NO VARNISH ZONE</p> <p>Peel here</p>
---	---	--



PAGE 02

PAGE 11

<p>12mm Blank for Glue Area</p>	<p>Drug Facts ⑫</p> <table border="1"> <thead> <tr> <th>Active ingredient (in each dose)</th> <th>Purpose</th> </tr> </thead> <tbody> <tr> <td>Polyethylene Glycol 3350, 17 grams (cap filled to the indicated line).....</td> <td>Osmotic Laxative</td> </tr> </tbody> </table>	Active ingredient (in each dose)	Purpose	Polyethylene Glycol 3350, 17 grams (cap filled to the indicated line).....	Osmotic Laxative	<p>Drug Facts (continued) ⑪</p> <ul style="list-style-type: none"> do not use more than 7 days children 16 years of age or under: ask a doctor <p>Other information</p> <ul style="list-style-type: none"> store at 20° - 25°C (68° - 77°F) tamper-evident: do not use if foil seal under cap, printed with
Active ingredient (in each dose)	Purpose					
Polyethylene Glycol 3350, 17 grams (cap filled to the indicated line).....	Osmotic Laxative					



PAGE 10

PAGE 03

<p>Drug Facts (continued) ⑫</p>	<p>Drug Facts (continued) ⑪</p>
--	--



<p>Drug Facts (continued) ④</p> <p>hot or room temperature) then drink</p> <ul style="list-style-type: none"> do not combine with starch-based thickeners used for difficulty swallowing ensure that the powder is fully dissolved before drinking do not drink if there are any clumps 	<p>Drug Facts (continued) ⑤</p> <p>Use</p> <ul style="list-style-type: none"> relieves occasional constipation (irregularity) generally produces a bowel movement in 1 to 3 days <p>Warnings</p> <p>Allergy alert: Do not use if you</p>
---	--

3

78mm

PAGE 04

PAGE 09

<p>Drug Facts (continued) ④</p> <p>are allergic to polyethylene glycol</p> <p>Do not use if you have kidney disease, except under the advice and supervision of a doctor</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> nausea, vomiting or abdominal pain 	<p>Drug Facts (continued) ⑤</p> <ul style="list-style-type: none"> adults and children 17 years of age and older: <ul style="list-style-type: none"> use once a day fill to top of white section in cap which is marked to indicate the correct dose (17g) stir and dissolve in any 4 to 8 ounces of beverage (cold,
---	--

3

78mm

PAGE 08

PAGE 05

<p>Drug Facts (continued) ⑧</p> <p>Directions</p> <ul style="list-style-type: none"> do not take more than directed unless advised by your doctor the bottle top is a measuring cap marked to contain 17grams of powder when filled to the indicated line (white section in cap) 	<p>Drug Facts (continued) ⑤</p> <ul style="list-style-type: none"> a sudden change in bowel habits that lasts over 2 weeks irritable bowel syndrome <p>Ask a doctor or pharmacist before use if you are taking a prescription drug</p> <p>When using this product you may</p>
--	--

3

← 78mm →


PAGE 06

PAGE 07

<p>Drug Facts (continued) ⑥</p> <p>have loose, watery, more frequent stools</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> • you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. • you get diarrhea ▶ 	<p>Drug Facts (continued) ⑦</p> <ul style="list-style-type: none"> • you need to use a laxative for longer than 1 week <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶</p>
--	---

↑ 3 ↓

← 78mm →



EZ2GO POLYETHYLENE GLYCOL 3350

polyethylene glycol 3350 powder, for solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:76470-050
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (PEG-75 - UNII:G2M7P15E5P)		POLYETHYLENE GLYCOL 3350	17 g in 17 g	
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76470-050-50	119 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/01/2026	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA203928	04/01/2026		

Labeler - Valley Medical Products, LLC (969389407)

Registrant - Valley Medical Products, LLC (969389407)

Revised: 2/2026

Valley Medical Products, LLC