

**SODIUM SULFACETAMIDE- sodium sulfacetamide liquid**  
**Method Pharmaceuticals, LLC**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----  
**Sodium Sulfacetamide 10%**

**Inactive Ingredients**

Butylated Hydroxytoluene, Citric Acid, Cetyl Alcohol, Cocamidopropyl Betaine, Disodium EDTA, Glycerin, Glyceryl Stearate SE, PEG-100 Stearate, Phenoxyethanol, Purified Water, Sodium Laureth Sulfate, Sodium Thiosulfate, Stearyl Alcohol, Triacetin, Xanthan Gum.

**Description**

Each gram contains 100 mg of sodium sulfacetamide in a vehicle consisting of: Butylated Hydroxytoluene, Citric Acid, Cetyl Alcohol, Cocamidopropyl Betaine, Disodium EDTA, Glycerin, Glyceryl Stearate SE, PEG-100 Stearate, Phenoxyethanol, Purified Water, Sodium Laureth Sulfate, Sodium Thiosulfate, Stearyl Alcohol, Triacetin, Xanthan Gum.

**Indications**

This product is intended for topical application in the following scaling dermatoses: seborrheic dermatitis and seborrhea sicca (dandruff). It also is indicated for the treatment of secondary bacterial infections of the skin due to organisms susceptible to sulfonamides.

**Contraindications**

This product is contraindicated in persons with known or suspected hypersensitivity to any of the ingredients of the product. This product is not to be used by patients with kidney disease.

**Dosage and Administration**

*Seborrheic dermatitis including seborrhea sicca* - Wash affected areas twice daily (morning and evening) or as directed by your physician. Avoid contact with eyes or mucous membranes. Wet skin and liberally apply to areas to be cleansed, massage gently into skin working into a full lather, rinse thoroughly, pat dry and repeat after 10 to 20 seconds. Rinsing with plain water will remove any excess medication. Repeat application as described above for 8 to 10 days or as directed by your physician. If skin dryness occurs it may be controlled by rinsing cleanser off sooner or using less frequently. Regular shampooing following the use of this product is not necessary, but the hair should be shampooed at least once a week. As the condition subsides, the interval between

applications may be lengthened. Applications once or twice weekly or every other week may prevent recurrence. Should the condition recur after stopping therapy, the application of this product should be reinitiated as at the beginning of treatment.

*Secondary cutaneous bacterial infections* - Wash affected areas twice daily (morning and evening) or as directed by your physician. Avoid contact with eyes or mucous membranes. Wet skin and liberally apply to areas to be cleansed, massage gently into skin for 10 to 20 seconds working into a full lather, rinse thoroughly and pat dry. Rinsing with plain water will remove any excess medication. Repeat application as described above for 8 to 10 days or as directed by your physician. If skin dryness occurs it may be controlled by rinsing cleanser off sooner or using less frequently. See package insert for full prescribing information.

## **Warnings**

WARNING: FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.  
KEEP OUT OF REACH OF CHILDREN.

Avoid contact with eyes, lips and mucous membranes.

See label booklet for Full prescribing information.

## **Storage**

Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F). Protect from freezing and excessive heat. Keep bottle tightly closed.

This bottle is not filled to the top but does contain 6 fl oz of product as identified on the front panel of the bottle.

To report a serious adverse event or obtain product information, call (877) 250-3427.

## **Package**



NDC 58657-477-06

# Sodium Sulfacetamide

(Sodium Sulfacetamide 10%)

10%  
Wash

For External Use Only.  
Not For Ophthalmic Use.

Rx Only  
Net Wt. 6 oz (177 g)

#### DESCRIPTION:

Each gram contains 100 mg of sodium sulfacetamide in a vehicle consisting of: Butylated Hydroxytoluene, Citric Acid, Cetyl Alcohol, Cocamidopropyl Betaine, Disodium EDTA, Glycerin, Glyceryl Stearate SE, PEG-100 Stearate, Phenoxethanol, Purified Water, Sodium Laureth Sulfate, Sodium Thiosulfate, Stearyl Alcohol, Triacetin, Xanthan Gum.

#### INDICATIONS:

This product is intended for topical application in the following scaling dermatoses: seborrheic dermatitis and seborrhea sicca (dandruff). It also is indicated for the treatment of secondary bacterial infections of the skin due to organisms susceptible to sulfonamides.

#### CONTRAINDICATIONS:

This product is contraindicated in persons with known or suspected hypersensitivity to any of the ingredients of the product. This product is not to be used by patients with kidney disease.

#### DOSAGE AND ADMINISTRATION:

**Seborrheic dermatitis including seborrhea sicca** - Wash affected areas twice daily (morning and evening) or as directed by your physician. Avoid contact with eyes or mucous membranes. Wet skin and liberally apply to areas to be cleansed, massage gently into skin working into a full lather, rinse thoroughly, pat dry and repeat after 10 to 20 seconds. Rinsing with plain water will remove any excess medication. Repeat application as described above for 8 to 10 days or as directed by your physician. If skin dryness occurs it may be controlled by rinsing cleanser off sooner or using less frequently. Regular shampooing following the use of this product is not necessary, but the hair should be shampooed at least once a week. As the condition subsides, the interval between applications may be lengthened. Applications once or twice weekly or every other week may prevent recurrence. Should the condition recur after stopping therapy, the application of this product should be reinstated as at the beginning of treatment.

**Secondary cutaneous bacterial infections** - Wash affected areas twice daily (morning and evening) or as directed by your physician. Avoid contact with eyes or mucous membranes. Wet skin and liberally apply to areas to be cleansed, massage gently into skin for 10 to 20 seconds working into a full lather, rinse thoroughly and pat dry. Rinsing with plain water will remove any excess medication. Repeat application as described above for 8 to 10 days or as directed by your physician. If skin dryness occurs it may be controlled by rinsing cleanser off sooner or using less frequently.

See package insert for full prescribing information.

#### WARNING: FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

**KEEP OUT OF REACH OF CHILDREN.** Avoid contact with eyes, lips and mucous membranes.

See label booklet for Full prescribing information.

**STORAGE:** Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F). Protect from freezing and excessive heat. Keep bottle tightly closed.

This bottle is not filled to the top but does contain 6 fl oz of product as identified on the front panel of the bottle.

To report a serious adverse event or obtain product information, call (877) 250-3427.

Manufactured in the U.S.A. for  
Method Pharmaceuticals, LLC  
Southlake, Texas 76092

Rev 09/21



Manufactured in the U.S.A. for  
Method Pharmaceuticals, LLC  
Southlake, Texas 76092  
Rev 09/21

**Package 12oz**



NDC 58657-477-12

# Sodium Sulfacetamide

(Sodium Sulfacetamide 10%)

**10% Wash**

**For External Use Only.  
Not For Ophthalmic Use.**

**Rx Only  
Net Wt. 12 oz. (355 g)**

**DESCRIPTION:**

Each gram contains 100 mg of sodium sulfacetamide in a vehicle consisting of: Butylated Hydroxytoluene, Citric Acid, Cetyl Alcohol, Cocamidopropyl Betaine, Disodium EDTA, Glycerin, Glyceryl Stearate SE, PEG-100 Stearate, Phenoxyethanol, Purified Water, Sodium Laureth Sulfate, Sodium Thiosulfate, Stearyl Alcohol, Triacetin, Xanthan Gum.

**INDICATIONS:**

This product is intended for topical application in the following scaling dermatoses: seborrheic dermatitis and seborrhea sicca (dandruff). It also is indicated for the treatment of secondary bacterial infections of the skin due to organisms susceptible to sulfonamides.

**CONTRAINDICATIONS:**

This product is contraindicated in persons with known or suspected hypersensitivity to any of the ingredients of the product. This product is not to be used by patients with kidney disease.

**DOSAGE AND ADMINISTRATION:**

**Seborrheic dermatitis including seborrhea sicca** - Wash affected areas twice daily (morning and evening) or as directed by your physician. Avoid contact with eyes or mucous membranes. Wet skin and liberally apply to areas to be dosed, massage gently into skin working into a full lather, rinse thoroughly, pat dry and repeat after 10 to 20 seconds. Rinsing with plain water will remove any excess medication. Repeat application as described above for 8 to 10 days or as directed by your physician. If skin dryness occurs it may be controlled by rinsing cleanser off sooner or using less frequently. Regular shampooing following the use of this product is not necessary, but the hair should be shampooed at least once a week. As the condition subsides, the interval between applications may be lengthened. Applications once or twice weekly or every other week may prevent recurrence. Should the condition recur after stopping therapy, the application of this product should be reinstated as at the beginning of treatment.

**Secondary cutaneous bacterial infections** - Wash affected areas twice daily (morning and evening) or as directed by your physician. Avoid contact with eyes or mucous membranes. Wet skin and liberally apply to areas to be dosed, massage gently into skin for 10 to 20 seconds working into a full lather, rinse thoroughly and pat dry. Rinsing with plain water will remove any excess medication. Repeat application as described above for 8 to 10 days or as directed by your physician. If skin dryness occurs it may be controlled by rinsing cleanser off sooner or using less frequently.

See package insert for full prescribing information.

**WARNING: FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE. KEEP OUT OF REACH OF CHILDREN.** Avoid contact with eyes, lips and mucous membranes.

See label booklet for Full prescribing information.

**STORAGE:** Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F). Protect from freezing and excessive heat. Keep bottle tightly closed.

To report a serious adverse event or obtain product information, call (877) 250-3427.

Manufactured in the U.S.A. for Method Pharmaceuticals, LLC Southlake, Texas 76092

Rev 09/21



3 58657 47712 7

Package 16 oz



NDC 58657-477-16

# Sodium Sulfacetamide

(Sodium Sulfacetamide 10%)

10% Wash

For External Use Only.  
Not For Ophthalmic Use.

Rx Only  
Net Wt. 16 oz. (480 g)

**DESCRIPTION:**

Each gram contains 100 mg of sodium sulfacetamide in a vehicle consisting of: Butylated Hydroxytoluene, Citric Acid, Cetyl Alcohol, Cocamidopropyl Betaine, Disodium EDTA, Glycerin, Glyceryl Stearate SE, PEG-100 Stearate, Phenoxyethanol, Purified Water, Sodium Laureth Sulfate, Sodium Thiosulfate, Stearyl Alcohol, Tricetin, Xanthan Gum.

**INDICATIONS:**

This product is intended for topical application in the following scaling dermatoses: seborrheic dermatitis and seborrhea sicca (dandruff), it also is indicated for the treatment of secondary bacterial infections of the skin due to organisms susceptible to sulfonamides.

**CONTRAINDICATIONS:**

This product is contraindicated in persons with known or suspected hypersensitivity to any of the ingredients of the product. This product is not to be used by patients with kidney disease.

**DOSEAGE AND ADMINISTRATION:**

**Seborrheic dermatitis including seborrhea sicca** - Wash affected areas twice daily (morning and evening) or as directed by your physician. Avoid contact with eyes or mucous membranes. Wet skin and liberally apply to areas to be cleansed, massage gently into skin working into a full lather, rinse thoroughly, pat dry and repeat after 10 to 20 seconds. Rinsing with plain water will remove any excess medication. Repeat application as described above for 8 to 10 days or as directed by your physician. If skin dryness occurs it may be controlled by rinsing cleanser off sooner or using less frequently. Regular shampooing following the use of this product is not necessary, but the hair should be shampooed at least once a week. As the condition subsides, the interval between applications may be lengthened. Applications once or twice weekly or every other week may prevent recurrence. Should the condition recur after stopping therapy, the application of this product should be reinstated as at the beginning of treatment.

**Secondary cutaneous bacterial infections** - Wash affected areas twice daily (morning and evening) or as directed by your physician. Avoid contact with eyes or mucous membranes. Wet skin and liberally apply to areas to be cleansed, massage gently into skin for 10 to 20 seconds working into a full lather, rinse thoroughly and pat dry. Rinsing with plain water will remove any excess medication. Repeat application as described above for 8 to 10 days or as directed by your physician. If skin dryness occurs it may be controlled by rinsing cleanser off sooner or using less frequently.

See label booklet for Full prescribing information.

**WARNING: FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.**

**KEEP OUT OF REACH OF CHILDREN.** Avoid contact with eyes, lips and mucous membranes.

**STORAGE:** Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F). Protect from freezing and excessive heat. Keep bottle tightly closed.

To report a serious adverse event or obtain product information, call (877) 250-3427.

Manufactured in the U.S.A. for  
Method Pharmaceuticals, LLC  
Southlake, Texas 76092

Rev 09/21



## SODIUM SULFACETAMIDE

sodium sulfacetamide liquid

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:58657-477
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>SULFACETAMIDE SODIUM</b> (UNII: 4NRT660KJQ) (SULFACETAMIDE - UNII:4965G3J0F5)	SULFACETAMIDE SODIUM	100 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>GLYCERYL MONOSTEARATE</b> (UNII: 230OU9XXE4)	
<b>PEG-100 STEARATE</b> (UNII: YD01N1999R)	

<b>PHENOXYETHANOL</b> (UNII: HIE492ZZ3T)	
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>SODIUM THIOSULFATE</b> (UNII: HX1032V43M)	
<b>STEARYL ALCOHOL</b> (UNII: 2KR89I4H1Y)	
<b>CETYL ALCOHOL</b> (UNII: 936JST6JCN)	
<b>COCAMIDOPROPYL BETAINE</b> (UNII: 5OCF3O11KX)	
<b>EDETATE DISODIUM ANHYDROUS</b> (UNII: 8NLQ36F6MM)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM LAURETH SULFATE</b> (UNII: BPV390UAP0)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58657-477-12	355 g in 1 BOTTLE; Type 0: Not a Combination Product	01/15/2022	
2	NDC:58657-477-16	480 g in 1 BOTTLE; Type 0: Not a Combination Product	01/15/2022	
3	NDC:58657-477-06	177 g in 1 BOTTLE; Type 0: Not a Combination Product	01/15/2022	

<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/15/2022	

**Labeler** - Method Pharmaceuticals, LLC (060216698)