


**SUCRALFATE- sucralfate tablet**  
**NCS HealthCare of KY, LLC dba Vanguard Labs**

-----  
**SUCRALFATE TABLETS, USP**  
**2210**  
**Rx only**

**DESCRIPTION**

Sucralfate, USP is an  $\alpha$ -D-glucopyranoside,  $\beta$ -D-fructofuranosyl-, octakis(hydrogen sulfate), aluminum complex.



R =  $SO_3Al(OH)_2$

Tablets for oral administration contain 1 g of sucralfate, USP and the following inactive ingredients: corn starch, magnesium stearate, and microcrystalline cellulose.

**Therapeutic Category**

antiulcer

**CLINICAL PHARMACOLOGY**

Sucralfate is only minimally absorbed from the gastrointestinal tract. The small amounts of the sulfated disaccharide that are absorbed are excreted primarily in the urine.

Although the mechanism of sucralfate's ability to accelerate healing of duodenal ulcers remains to be fully defined, it is known that it exerts its effect through a local, rather than systemic, action. The following observations also appear pertinent:

1. Studies in human subjects and with animal models of ulcer disease have shown that sucralfate forms an ulcer-adherent complex with proteinaceous exudate at the ulcer site.
2. *In vitro*, a sucralfate-albumin film provides a barrier to diffusion of hydrogen ions.
3. In human subjects, sucralfate given in doses recommended for ulcer therapy inhibits

pepsin activity in gastric juice by 32%.

4. *In vitro*, sucralfate adsorbs bile salts.

These observations suggest that sucralfate's antiulcer activity is the result of formation of an ulcer-adherent complex that covers the ulcer site and protects it against further attack by acid, pepsin, and bile salts. There are approximately 14 to 16 mEq of acid-neutralizing capacity per 1 g dose of sucralfate.

## CLINICAL TRIALS

### Acute Duodenal Ulcer

Over 600 patients have participated in well-controlled clinical trials worldwide. Multicenter trials conducted in the United States, both of them placebo-controlled studies with endoscopic evaluation at 2 and 4 weeks, showed:

#### STUDY 1

Treatment Groups	Ulcer Healing/No. Patients	
	2 wk	4 wk (Overall)
Sucralfate	37/105 (35.2%)	82/109 (75.2%)
Placebo	26/106 (24.5%)	68/107 (63.6%)

#### STUDY 2

Treatment Groups	Ulcer Healing/No. Patients	
	2 wk	4 wk (Overall)
Sucralfate	8/24 (33%)	22/24 (92%)
Placebo	4/31 (13%)	18/31 (58%)

The sucralfate-placebo differences were statistically significant in both studies at 4 weeks but not at 2 weeks. The poorer result in the first study may have occurred because sucralfate was given 2 hours after meals and at bedtime rather than 1 hour before meals and at bedtime, the regimen used in international studies and in the second United States study. In addition, in the first study liquid antacid was utilized as needed, whereas in the second study antacid tablets were used.

### Maintenance Therapy After Healing of Duodenal Ulcer

Two double-blind randomized placebo-controlled U.S. multicenter trials have demonstrated that sucralfate (1 g bid) is effective as maintenance therapy following healing of duodenal ulcers.

In one study, endoscopies were performed monthly for 4 months. Of the 254 patients who enrolled, 239 were analyzed in the intention-to-treat life table analysis presented below.

### Duodenal Ulcer Recurrence Rate (%)

<b>Months of Therapy</b>					
<b>Drug</b>	<b>n</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
Sucralfate	122	20*	30*	38†	42†
Placebo	117	33	46	55	63

\* p < 0.05

† p < 0.01

In this study, prn antacids were not permitted.

In the other study, scheduled endoscopies were performed at 6 and 12 months, but for-cause endoscopies were permitted as symptoms dictated. Median symptom scores between the sucralfate and placebo groups were not significantly different. A life table intention-to-treat analysis for the 94 patients enrolled in the trial had the following results:

#### **Duodenal Ulcer Recurrence Rate (%)**

<b>Drug</b>	<b>n</b>	<b>6 months</b>	<b>12 months</b>
Sucralfate	48	19*	27*
Placebo	46	54	65

\* p < 0.002

In this study, prn antacids were permitted.

Data from placebo-controlled studies longer than 1 year are not available.

## **INDICATIONS AND USAGE**

Sucralfate tablets, USP are indicated in:

- Short-term treatment (up to 8 weeks) of active duodenal ulcer. While healing with sucralfate may occur during the first week or two, treatment should be continued for 4 to 8 weeks unless healing has been demonstrated by x-ray or endoscopic examination.
- Maintenance therapy for duodenal ulcer patients at reduced dosage after healing of acute ulcers.

## **CONTRAINDICATIONS**

Sucralfate tablets are contraindicated in patients with known hypersensitivity reactions to the active substance or to any of the excipients.

## **PRECAUTIONS**

The physician should read the **PRECAUTIONS** section when considering the use of this drug in pregnant or pediatric patients, or patients of childbearing potential.

Duodenal ulcer is a chronic, recurrent disease. While short-term treatment with sucralfate can result in complete healing of the ulcer, a successful course of treatment with sucralfate should not be expected to alter the post healing frequency or severity of

duodenal ulceration.

Isolated reports of sucralfate tablet aspiration with accompanying respiratory complications have been received. Therefore, sucralfate tablets should be used with caution by patients who have known conditions that may impair swallowing, such as recent or prolonged intubation, tracheostomy, prior history of aspiration, dysphagia, or any other conditions that may alter gag and cough reflexes, or diminish oropharyngeal coordination or motility.

## **Special Populations**

### **Chronic Renal Failure and Dialysis Patients**

When sucralfate is administered orally, small amounts of aluminum are absorbed from the gastrointestinal tract. Concomitant use of sucralfate with other products that contain aluminum, such as aluminum-containing antacids, may increase the total body burden of aluminum. Patients with normal renal function receiving the recommended doses of sucralfate and aluminum-containing products adequately excrete aluminum in the urine. Patients with chronic renal failure or those receiving dialysis have impaired excretion of absorbed aluminum. In addition, aluminum does not cross dialysis membranes because it is bound to albumin and transferrin plasma proteins. Aluminum accumulation and toxicity (aluminum osteodystrophy, osteomalacia, encephalopathy) have been described in patients with renal impairment. Sucralfate should be used with caution in patients with chronic renal failure.

## **Drug Interactions**

Some studies have shown that simultaneous sucralfate administration in healthy volunteers reduced the extent of absorption (bioavailability) of single doses of the following: cimetidine, digoxin, fluoroquinolone antibiotics, ketoconazole, l-thyroxine, phenytoin, quinidine, ranitidine, tetracycline, and theophylline. Subtherapeutic prothrombin times with concomitant warfarin and sucralfate therapy have been reported in spontaneous and published case reports. However, two clinical studies have demonstrated no change in either serum warfarin concentration or prothrombin time with the addition of sucralfate to chronic warfarin therapy.

The mechanism of these interactions appears to be nonsystemic in nature, presumably resulting from sucralfate binding to the concomitant agent in the gastrointestinal tract. In all case studies to date (cimetidine, ciprofloxacin, digoxin, norfloxacin, ofloxacin, and ranitidine), dosing the concomitant medication 2 hours before sucralfate eliminated the interaction. Because of the potential of sucralfate to alter the absorption of some drugs, sucralfate should be administered separately from other drugs when alterations in bioavailability are felt to be critical. In these cases, patients should be monitored appropriately.

## **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Chronic oral toxicity studies of 24 months' duration were conducted in mice and rats at doses up to 1 g/kg (12 times the human dose).

There was no evidence of drug-related tumorigenicity. A reproduction study in rats at doses up to 38 times the human dose did not reveal any indication of fertility impairment. Mutagenicity studies were not conducted.

## **Pregnancy**

### Teratogenic Effects

#### *Pregnancy category B*

Teratogenicity studies have been performed in mice, rats, and rabbits at doses up to 50 times the human dose and have revealed no evidence of harm to the fetus due to sucralfate. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

## **Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when sucralfate is administered to a nursing woman.

## **Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

## **Geriatric Use**

Clinical studies of sucralfate tablets did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy (see **DOSAGE AND ADMINISTRATION**).

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function (see **PRECAUTIONS, Special Populations, Chronic Renal Failure and Dialysis Patients**). Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

## **ADVERSE REACTIONS**

Adverse reactions to sucralfate in clinical trials were minor and only rarely led to discontinuation of the drug. In studies involving over 2700 patients treated with sucralfate tablets, adverse effects were reported in 129 (4.7%).

Constipation was the most frequent complaint (2%). Other adverse effects reported in less than 0.5% of the patients are listed below by body system:

### **Gastrointestinal**

diarrhea, nausea, vomiting, gastric discomfort, indigestion, flatulence, dry mouth

### **Dermatological**

pruritus, rash

## **Nervous System**

dizziness, insomnia, sleepiness, vertigo

## **Other**

back pain, headache

Postmarketing cases of hypersensitivity have been reported with the use of sucralfate tablets, including dyspnea, lip swelling, pruritus, rash, and urticaria.

Cases of anaphylactic reactions, bronchospasm, laryngeal edema, edema of the mouth, pharyngeal edema, respiratory tract edema and swelling of the face have been reported with an unknown oral formulation of sucralfate.

Bezoars have been reported in patients treated with sucralfate. The majority of patients had underlying medical conditions that may predispose to bezoar formation (such as delayed gastric emptying) or were receiving concomitant enteral tube feedings.

Inadvertent injection of insoluble sucralfate and its insoluble excipients has led to fatal complications, including pulmonary and cerebral emboli. Sucralfate is **not** intended for intravenous administration.

## **OVERDOSAGE**

Due to limited experience in humans with overdosage of sucralfate, no specific treatment recommendations can be given. Acute oral toxicity studies in animals, however, using doses up to 12 g/kg body weight, could not find a lethal dose. Sucralfate is only minimally absorbed from the gastrointestinal tract. Risks associated with acute overdosage should, therefore, be minimal. In rare reports describing sucralfate overdose, most patients remained asymptomatic. Those few reports where adverse events were described included symptoms of dyspepsia, abdominal pain, nausea, and vomiting.

## **DOSAGE AND ADMINISTRATION**

### **Active Duodenal Ulcer**

The recommended adult oral dosage for duodenal ulcer is 1 g four times per day on an empty stomach.

Antacids may be prescribed as needed for relief of pain but should not be taken within one-half hour before or after sucralfate.

While healing with sucralfate may occur during the first week or two, treatment should be continued for 4 to 8 weeks unless healing has been demonstrated by x-ray or endoscopic examination.

### **Maintenance Therapy**

The recommended adult oral dosage is 1 g twice a day.

## **Elderly**

In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy (see **PRECAUTIONS, Geriatric Use**).

Call your doctor for medical advice about side effects. You may report side effects to TEVA USA, PHARMACOVIGILANCE at 1-866-832-8537 or [drug.safety@tevapharm.com](mailto:drug.safety@tevapharm.com); or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

## **HOW SUPPLIED**

Sucralfate tablets, USP are available as white, capsule-shaped, biconvex, scored tablets, debossed "TEVA" on one side, and "22" and "10" on the scored side, containing 1 gram of sucralfate USP, packaged in blistercards of 30 (NDC 0615-4517-39), and unit-dose boxes of 30 (NDC 0615-4517-30) tablets.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required).

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Manufactured In Croatia By:

**PLIVA HRVATSKA d.o.o.**

Zagreb, Croatia

Manufactured For:

**TEVA PHARMACEUTICALS USA, INC.**

North Wales, PA 19454

Rev. L 9/2015

**Package/Label Display Panel - BINGO CARD**



VLI NDC 0615-4517-39  
Sucralfate  
Tablet USP 1 gram

(Teva NDC 0093-2210-01)  
**Sucralfate**  
**Tablets USP 1 gram**



LOT 4517-  
EXP

Rx Only

4517-AB-39 v01

QTY  
30

Mfg By Pliva HRVATSKA  
for Teva USA  
(NDC 0093-2210-01)  
PKG BY VANGARD  
GLASGOW, KY 42141

			16	8
30	23	15	7	
29	22	14	6	
28	21	13	5	
27	20	12	4	
26	19	11	3	
25	18	10	2	
24	17	9	1	



The overall configuration of this package is a trademark of Omnicare, Inc.

**Received:** \_\_\_\_\_

31	24	16	8
30	23	15	7
29	22	14	6
28	21	13	5
27	20	12	4
26	19	11	3
25	18	10	2
24	17	9	1

Start Date \_\_\_\_\_ Start Time \_\_\_\_\_

Each tablet contains  
1 gram sucralfate.

Store at 20° to 25°C (68° to 77°F) (See  
USP Controlled Room Temperature).

Usual Dosage: See package insert for full  
prescribing information.

**FOR INSTITUTIONAL USE ONLY**

<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>4517-AB-B-v03 Vanguard Labs Glasgow, KY 42141</p> <p>LOT 4517- </p>
<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>
<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>
<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>
<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>
<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>
<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>	<p>Pkg by Vanguard, Glasgow, KY 42141 <b>Sucralfate Tablet</b> USP 1 gram</p> <p>LOT 4517- </p>

**Package/Label Display Panel - UNIT-DOSE**

4517-AB-JD1-v01

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

Pkg By Vanguard, Glasgow, KY 42141  
**Sucralfate**  
 Tab USP 1 gram  
 LOT 4517-  
  
 0093221001

<b>VG NDC 0615-4517-30</b> <b>Sucralfate Tablets USP 1 gram</b> QTY 30 Rx Only Each tablet contains 1 gram sucralfate. Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].  LOT: EXP:	<b>VG NDC 0615-4517-30</b> <b>Sucralfate Tablets USP 1 gram</b> Rx Only FOR INSTITUTIONAL USE ONLY QTY 30 LOT: EXP: Manufactured By <b>Pilva HRVA TSKA</b> or <b>Teva USA</b> NDC 0093-2210-01 Usual Dosage: See package insert for full prescribing information.  009322100130
---	---

# SUCRALFATE

sucralfate tablet

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0615-4517(NDC:0093-2210)
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>SUCRALFATE</b> (UNII: XX73205DH5) (SUCRALFATE - UNII:XX73205DH5)	SUCRALFATE	1 g

## Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL (capsule-shaped)	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	TEVA;22;10
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0615-4517-39	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	01/08/2010	
2	NDC:0615-4517-30	6 in 1 BOX, UNIT-DOSE	01/08/2010	01/31/2026
2		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA070848	11/11/1996	

**Labeler** - NCS HealthCare of KY, LLC dba Vanguard Labs (050052943)

### Establishment

Name	Address	ID/FEI	Business Operations
NCS HealthCare of KY, LLC dba Vanguard Labs		050052943	repack(0615-4517)

Revised: 6/2025

NCS HealthCare of KY, LLC dba Vanguard Labs