

NICOTINE- nicotine polacrilex gum, chewing Walgreens

Drug Facts

Active ingredient (in each chewing piece)

Nicotine polacrilex (equal to 4 mg nicotine)

Purpose

Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding,

only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum according to the following 12 week schedule

| Weeks 1 to 6 | Weeks 7 to 9 | Weeks 10 to 12 |
|----------------------------|----------------------------|----------------------------|
| 1 piece every 1 to 2 hours | 1 piece every 2 to 4 hours | 1 piece every 4 to 8 hours |

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat the process until most of the tingle is gone (about 30 minutes)
- do not eat or drink 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- It is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information

- **each piece contains:** calcium 110 mg, sodium 11 mg
- store at 20-25°C (68-77°F)
- protect from light

Inactive ingredients

acesulfame potassium, butylated hydroxytoluene, carnauba wax, D&C yellow #10 lake, FD&C blue #2 lake, FD&C red #40, FD&C Yellow #6 lake, flavors, gum base, sodium bicarbonate, sodium carbonate, sorbitol, talc

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredient in Nicorette® Gum††

UNCOATED

Nicotine Gum

NICOTINE POLACRILEX GUM 4 mg (NICOTINE)

STOP SMOKING AID

Sugar Free

Original flavor

PIECES

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette **MORE THAN 30 MINUTES** after waking up, use Nicotine Polacrilex Gum USP, 2 mg

TO INCREASE YOUR SUCCESS IN QUITTING:

- 1.** You must be motivated to quit.
- 2.** Use Enough - Chew at least **9** pieces of Nicotine Polacrilex Gum per day during the first six weeks.
- 3.** Use Long Enough - Use Nicotine Polacrilex Gum for the full 12 weeks.
- 4. Use With a Support Program** as directed in the enclosed User's Guide.

To remove the gum, tear off single unit.

Peel off backing starting at corner with loose edge.

Push gum through foil

- **NOT FOR SALE TO THOSE UNDER 18 YEARS OF AGE**
- **PROOF OF AGE REQUIRED**
- **NOT FOR SALE IN VENDING MACHINES OR FROM ANY SOURCE WHERE PROOF OF AGE CANNOT BE VERIFIED**

††This product is not manufactured or distributed by Haleon US LP, owner of the registered trademark Nicorette®.

See inside for more details or visit www.nitthehabit.com

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN

DISTRIBUTED BY: **WALGREEN CO**

DEERFIELD, IL 60015

Package Label

TO INCREASE YOUR SUCCESS IN QUITTING:

- 1. You must be motivated to quit.
- 2. Use Enough - Chew at least 8 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
- 3. Use Long Enough - Use Nicotine Polacrilex Gum for the full 12 weeks.
- 4. Use with a support program as directed in the enclosed User's Guide.



To remove the gum, peel off
single unit.
Roll off backing
starting at corner
with loose edge.
Push gum
through foil.

For tools, resources and motivation, please visit
Walgreens.com/QuitSmoking

NCTC 0368-0308-92

Walgreens



Compare to the active
ingredient in Nicorette® Gum*

UNCOATED
Nicotine Gum

NICOTINE POLACRILEX GUM 4 mg (NICOTINE)
STOP SMOKING AID

Sugar Free

300
PIECES

4
mg



KTML182
Original flavor

4 mg EACH

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE AT THE BEGINNING OF WORKING UP.
If you smoke your first cigarette **SOONER THAN 30 MINUTES** after waking up, use Nicotine Polacrilex Gum, 4 mg.

© 2013 Walgreens. All rights reserved. 31423

100% SATISFACTION
GUARANTEE

Walgreens.com ©2013 Walgreen Co.

ITEM 609601 WT104Z 104Z-4

PRODUCT OF IRELAND
WALGREENS - RDV-1004

Walgreens is committed to providing the highest quality products and services to our customers. We strive to be the most convenient and most helpful place to shop. Our commitment to customer service is reflected in our 100% Satisfaction Guarantee. If you are not completely satisfied with your purchase, we will refund the purchase price of the product, with or without a receipt.

Manufacturing Information: Nicotine Polacrilex Gum, 4 mg, is manufactured by Nicorette, Inc., located at 10050 West Valley Parkway, Minneapolis, MN 55426. The gum is made from polybutadiene, polyethylene glycol, and other ingredients. Each piece of gum contains 4 mg of nicotine.

Directions for Use: Chew one piece of gum slowly and continuously for 30 minutes, up to 8 pieces per day. Do not swallow the gum. Do not eat, drink, or use tobacco during the 30-minute chewing period. Use the gum as directed in the enclosed User's Guide.

Warnings: Do not use if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you have a history of heart disease, high blood pressure, or diabetes. Do not use if you are taking other medications, as they may interact with the nicotine. Do not use if you are taking other products containing nicotine.

Directions for Disposal: Do not dispose of the gum in fire. Dispose of the gum in the trash.



WALGREENS Sugar Free Uncoated Nicotine Gum Original Flavor

NICOTINE

nicotine polacrilex gum, chewing

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:0363-9350 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------|
| NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R) | NICOTINE | 4 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| ACESULFAME POTASSIUM (UNII: 23OV73Q5G9) | |
| BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| SODIUM BICARBONATE (UNII: 8MDF5V39QO) | |
| SODIUM CARBONATE (UNII: 45P3261C7T) | |
| TALC (UNII: 7SEV7J4R1U) | |
| SORBITOL (UNII: 506T60A25R) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| FD&C BLUE NO. 2 (UNII: L06K8R7DQK) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |

Product Characteristics

| | | | |
|-----------------|--------|---------------------|----------|
| Color | white | Score | no score |
| Shape | SQUARE | Size | 14mm |
| Flavor | | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:0363-9350-30 | 300 in 1 CARTON | 12/27/2019 | |
| 1 | | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA074707 | 12/27/2019 | |

Labeler - Walgreens (008965063)

Revised: 10/2024

Walgreens