

MIDOL COMPLETE- acetaminophen, caffeine, and pyrilamine maleate tablet
MIDOL COMPLETE, TRAVEL BASIX- acetaminophen, caffeine, and pyrilamine maleate tablet
Lil' Drug Store Products, Inc

Midol ® Complete

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 500 mg	Pain reliever
Caffeine 60 mg	Diuretic
Pyrilamine maleate 15 mg	Antihistamine

Uses

for the temporary relief of these symptoms associated with menstrual periods:

- cramps
- bloating
- water-weight gain
- headache
- backache
- muscle aches
- fatigue

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- adults and children 12 years and older:
 - take 2 caplets with water
 - repeat every 6 hours, as needed
 - do not exceed 6 caplets per day
- children under 12 years: consult a doctor

Other information

- Store at room temperature.

Avoid excessive heat above 40°C (104°F).

Inactive ingredients

carnauba wax, croscarmellose sodium, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, propylene glycol, shellac, titanium dioxide, triacetin

Questions or comments?

1-800-331-4536 (Mon-Fri 9AM-5PM EST)

Midol[®] Complete 6ct, Lil' Drug Store[®] - PDP/Package

Midol[®]

Complete

Acetaminophen/Caffeine/Pyrilamine Maleate

Pain Reliever/Diuretic/Antihistamine

MULTI-SYMPTOM RELIEF OF

CRAMPS • BLOATING • FATIGUE

HEADACHE • BACKACHE

[Bayer cross]

[caplet image]

6

Caplets

[Lil' Drug Store[®] logo]

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have

- liver disease • glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- you may get drowsy • avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

Stop use and ask a doctor if

- new symptoms occur • redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

If pregnant or breast-feeding, ask a health professional before use.

DO NOT USE IF INNER POUCH OR CARTON IS OPEN OR DAMAGED!

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 500 mg	Pain reliever
Caffeine 60 mg	Diuretic
Pyrilamine maleate 15 mg	Antihistamine

Purpose

Uses for the temporary relief of these symptoms associated with menstrual periods:

- cramps • bloating
- water-weight gain • headache
- backache • muscle aches
- fatigue

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount for this product.
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening • blisters • rash • hives
- facial swelling • asthma (wheezing) • shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

LIFT PANEL FOR CONTINUED DRUG FACTS

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have

- liver disease • glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- you may get drowsy • avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

Stop use and ask a doctor if

- new symptoms occur • redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

If pregnant or breast-feeding, ask a health professional before use.

Midol[®] Complete 4ct, TRAVEL BASIX - PDP/Package

Midol[®]

COMPLETE

Acetaminophen/Caffeine/Pyrilamine Maleate

Pain Reliever/Diuretic/Antihistamine

MULTI-SYMPTOM RELIEF OF

CRAMPS • BLOATING • FATIGUE

HEADACHE • BACKACHE

[BAYER cross]

[caplets image]

4

Caplets

[TRAVEL BASIX logo]



Midol[®] Complete , 25ct Dispens-It - PDP/Package

Midol[®]

Complete

[Bayer cross]

**MULTI-SYMPTOM RELIEF OF
CRAMPS · BLOATING
FATIGUE · HEADACHE · BACKACHE**

[caplets (2) image]

25 POUCHES

2 CAPLETS EACH

[Lil' Drug Store logo]

Do not use if pouch is open or torn.

Drug Facts
Active ingredients (in each caplet)

Acetaminophen 160 mg	Pain reliever
Caffeine 65 mg	Diuretic
Pyrilamine maleate 15 mg	Antihistamine

Uses
 For the temporary relief of these symptoms associated with menstrual periods:
 • cramps • bloating • water-weight gain
 • headache • backache • muscle aches • fatigue

Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
 • more than 6 caplets in 24 hours, which is the maximum daily amount for this product
 • with other drugs containing acetaminophen
 • 3 or more alcoholic drinks every day while using this product

Other information
 Allergy alert: acetaminophen may cause severe or severe allergic reactions. Symptoms may include:
 • skin rash • hives • facial swelling • asthma (wheezing) • shock
 If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Do not use
 • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 • if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
 • liver disease • gallstones • difficulty in urination due to enlargement of the prostate gland
 • a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are
 • taking the blood thinning drug warfarin
 • taking sedatives or tranquilizers

Drug Facts (continued)
When using this product
 • you may get drowsy • avoid alcoholic drinks
 • excitability may occur, especially in children
 • alcohol, sedatives, and tranquilizers may increase drowsiness
 • be careful when driving a motor vehicle or operating machinery
 • limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

Stop use and ask a doctor if
 • new symptoms occur
 • redness or swelling is present
 • pain gets worse or lasts more than 10 days
 • fever gets worse or lasts more than 3 days
 • if pregnant or breast-feeding, ask a health professional before use.
 • Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
 • Do not take more than the recommended dose
 • adults and children 12 years and older: take 2 caplets with water. Repeat every 6 hours, as needed. Do not exceed 6 caplets per day
 • children under 12 years: consult a doctor

Other information
 • Store at room temperature. Avoid excessive heat above 40°C (104°F)

Inactive ingredients
 • carmellose sodium, croscarmellose sodium, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, polyethylene glycol, shellac, titanium dioxide, triacetin

Questions or comments?
 1-800-351-4338 (Mon-Fri 9AM-5PM EST)

Bayer Bayer HealthCare LLC, Whippany, NJ 07981

This product is packaged by L.P. Drug Store Products, Inc., 2000 Central Express Lane, Cedar Rapids, IA 52404 1-577-577-5535, Mon-Fri 9 AM-4:30 PM CT. Item is a product manufactured by Bayer.

Bayer, the Bayer Cross and Midol are registered trademarks of Bayer, used with permission.


M

Midol[®]
COMPLETE

Acetaminophen/Caffeine/Pyrimilamine Maleate
 Pain Reliever/Diuretic/Antihistamine

MULTI-SYMPTOM RELIEF OF
CRAMPS • BLOATING
FATIGUE • HEADACHE • BACKACHE

34670-05-08-22




0 94781 27964 9


M

Midol[®]
COMPLETE


Acetaminophen/Caffeine/Pyrimilamine Maleate
 Pain Reliever/Diuretic/Antihistamine

MULTI-SYMPTOM RELIEF OF
CRAMPS • BLOATING
FATIGUE • HEADACHE • BACKACHE





25 POUCHES
2 CAPLETS EACH





M

Midol[®]
COMPLETE


Acetaminophen/Caffeine/Pyrimilamine Maleate
 Pain Reliever/Diuretic/Antihistamine

MULTI-SYMPTOM RELIEF OF
CRAMPS • BLOATING
FATIGUE • HEADACHE • BACKACHE





25 POUCHES
2 CAPLETS EACH



Midol[®] Complete 30ct Dispens-It Carton - PDP/Package

Midol®

Complete

Acetaminophen/Caffeine/Pyrimidine Maleate

Pain Reliever/Diuretic/Antihistamine

[Bayer cross]


**MULTI-SYMPTOM RELIEF OF
CRAMPS · BLOATING
FATIGUE · HEADACHE · BACKACHE**

[caplets (2) image]

30 POUCHES

2 CAPLETS EACH

[Lil' Drug Store logo]



MIDOL COMPLETE

acetaminophen, caffeine, and pyrilamine maleate tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-9745
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	500 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)		CAFFEINE	60 mg
PYRILAMINE MALEATE (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL)		PYRILAMINE MALEATE	15 mg

Inactive Ingredients	
Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B710)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	

Product Characteristics			
Color	white	Score	no score

Shape	OVAL	Size	17mm	
Flavor		Imprint Code	Midol	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-9745-2	2 in 1 CARTON	12/01/2011	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M013	12/01/2011	

MIDOL COMPLETE, TRAVEL BASIX			
acetaminophen, caffeine, and pyrilamine maleate tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-6445
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	500 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)		CAFFEINE	60 mg
PYRILAMINE MALEATE (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL)		PYRILAMINE MALEATE	15 mg
Inactive Ingredients			
Ingredient Name			Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)			
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)			
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
STARCH, CORN (UNII: O8232NY3SJ)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
SHELLAC (UNII: 46N107B71O)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
TRIACETIN (UNII: XHX3C3X673)			

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	Midol
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-6445-2	2 in 1 CARTON	11/29/2022	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:66715-6445-7	25 in 1 CARTON	04/19/2022	
2		2 in 1 POUCH; Type 0: Not a Combination Product		
3	NDC:66715-6445-8	30 in 1 CARTON	04/19/2022	
3		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/19/2022	

Labeler - Lil' Drug Store Products, Inc (093103646)

Revised: 7/2024

Lil' Drug Store Products, Inc