

**COCCIAID FOR CALVES- amprolium solution**  
**Aurora Pharmaceutical, Inc**

-----  
**CocciAid® for Calves**

**(amprolium)**

**9.6% Oral Solution Coccidiostat**

**INDICATIONS FOR USE:** An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

**DOSAGE AND ADMINISTRATION:**

<b>Protocol</b>	<b>5-Day Treatment Protocol</b>	<b>21-Day Prevention</b>
	Daily Dosage: 10 mg amprolium/kg (10 mg per 2.2 lb body weight)	Daily Dosage: 5 mg amprolium/kg (5 mg per 2.2 lb body weight)

**USE DIRECTIONS:** There are **96 mg** of amprolium in every **1 mL** of CocciAid for Calves 9.6% Solution.

1 fl oz = 29.57 mL

8 fl oz = 1/2 pint (236.56 mL)

1 fl oz = 2 measuring tablespoons

16 fl oz = 1 pint (473.12 mL)

**IN DRINKING WATER**

Mix CocciAid for Calves daily with fresh drinking water.

**5-DAY TREATMENT:** Add CocciAid for Calves 9.6% Oral Solution to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add CocciAid for Calves 9.6% Oral Solution to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

**CocciAid® for Calves is a registered trademark of Aurora Pharmaceutical, Inc.**

**AS A DRENCH**

CocciAid for Calves drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

**5-DAY TREATMENT:** Add 3 fl oz CocciAid for Calves 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Use on a herd basis only; when one or more calves

show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 fl oz of CocciAid for Calves 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

Use as the sole source of amprolium.

**WITHDRAWAL PERIODS AND RESIDUE WARNINGS:** Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**USER SAFETY WARNINGS: NOT FOR HUMAN USE. Keep this and all drugs out of the reach of children.**

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Aurora Pharmaceutical at 1-888-215-1256 or [www.aurorapharmaceutical.com](http://www.aurorapharmaceutical.com). For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at [www.fda.gov/reportanimalae](http://www.fda.gov/reportanimalae).

**PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION.** For irritations, flush with plenty of water; get medical attention.

**STORAGE:** Store between 5°-25°C (41°- 7°F) with brief excursions to 40°C.

**Benzoic acid 0.1% added as preservative.**

Restricted Drug (California) - Use Only as Directed

**CocciAid® for Calves**

**(amprolium)**

**9.6% Oral Solution Coccidiostat**

**Net Contents:** 1 gal. (128 fl. oz) (3,785 ml)

**Active Ingredient:** amprolium.....9.6%

Approved by FDA under ANADA # 200-780

Distributed by:

Aurora Pharmaceutical, Inc.

Northfield, MN 55057

IN 50-1644 rev 03/2024

**REORDER NO: 24004**

Aurora Pharmaceutical, Inc.  
NORTHFIELD, MN 55057  
1-888-215-1256  
www.aurorapharmaceutical.com  
IN 50-1644 03/2024

TAKE TIME  
OBSERVE LABEL  
DIRECTIONS



3 51072 11000 4

Lot  
Exp

**NDC 51072-110-00**



**CocciAid® for Calves  
(amprolium)  
9.6% Oral Solution Coccidiostat**

Net Contents: 1 gal. (128 fl. oz) (3,785 mL)  
Active Ingredient: amprolium.....9.6%  
Approved by FDA under ANADA # 200-780

Manufactured by:  
Aurora Pharmaceutical, Inc.  
Northfield, MN 55057

**INDICATIONS FOR USE:** An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

**DOSEAGE AND ADMINISTRATION:**

**5-Day Treatment Protocol**  
Daily Dosage: 10 mg amprolium/kg  
(10 mg per 2.2 lb body weight)

**21-Day Prevention Protocol**  
Daily Dosage: 5 mg amprolium/kg  
(5 mg per 2.2 lb body weight)

**USE DIRECTIONS:** There are 96 mg of amprolium in every 1 mL of CocciAid for Calves 9.6% Solution.  
1 fl oz = 29.57 mL  
1 fl oz = 2 measuring tablespoons  
8 fl oz = 1/2 pint (236.56 mL)  
16 fl oz = 1 pint (473.12 mL)

**IN DRINKING WATER**  
Mix CocciAid for Calves daily with fresh drinking water.

**5-DAY TREATMENT:** Add CocciAid for Calves 9.6% Oral Solution to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only: when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add CocciAid for Calves 9.6% Oral Solution to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

CocciAid® for Calves is a registered trademark of Aurora Pharmaceutical, Inc.

**AS A DRENCH**  
CocciAid for Calves drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

**5-DAY TREATMENT:** Add 3 fl oz CocciAid for Calves 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 fl oz of CocciAid for Calves 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days. Use as the sole source of amprolium.

**WITHDRAWAL PERIODS AND RESIDUE WARNINGS:** Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**USER SAFETY WARNINGS: NOT FOR HUMAN USE.** Keep this and all drugs out of the reach of children. To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Aurora Pharmaceutical at 1-888-215-1256 or www.aurorapharmaceutical.com. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-1088 or online at www.fda.gov/reportmalae.

**PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY, MAY CAUSE EYE IRRITATION.** For irritations, flush with plenty of water; get medical attention.

**STORAGE:** Store between 5°-25° C (41°-77° F) with brief excursions to 40° C.

**Benzic acid 0.1% added as preservative.**  
Restricted Drug (California) - Use Only as Directed

# COCCIAID FOR CALVES

amprolium solution

## Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:51072-110
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>AMPROLIUM</b> (UNII: 95CO6N199Q) (AMPROLIUM ION - UNII:H2T307KMZR)	AMPROLIUM	96 mg in 1 mL

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51072-110-00	3785 mL in 1 BOTTLE, PLASTIC		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200780	05/08/2024	

**Labeler** - Aurora Pharmaceutical, Inc (832848639)

## Establishment

Name	Address	ID/FEI	Business Operations
Aurora Pharmaceutical, Inc		832848639	manufacture

## Establishment

Name	Address	ID/FEI	Business Operations
Zhejiang Dayang Biotech Group Co., Ltd.		420600785	api manufacture

